




## ORIGINAL ARTICLE

# Curative anticoagulation prevents endothelial lesion in COVID-19 patients

Lina Khider<sup>1</sup> | Nicolas Gendron<sup>2,3</sup>  | Guillaume Goudot<sup>1</sup> | Richard Chocron<sup>4,5</sup> | Caroline Hauw-Berlemont<sup>6</sup> | Charles Cheng<sup>1</sup> | Nadia Rivet<sup>2,3</sup> | Helene Pere<sup>4,7</sup> | Ariel Roffe<sup>1</sup> | Sébastien Clerc<sup>8</sup> | David Lebeaux<sup>9</sup> | Benjamin Debuc<sup>10</sup> | David Veyer<sup>11,7</sup> | Bastien Rance<sup>12</sup> | Pascale Gaussem<sup>2,13</sup> | Sébastien Bertil<sup>2,13</sup> | Cécile Badoual<sup>4,14</sup> | Philippe Juvin<sup>15</sup> | Benjamin Planquette<sup>2,16</sup> | Emmanuel Messas<sup>1</sup> | Olivier Sanchez<sup>2,16</sup> | Jean-Sébastien Hulot<sup>4,17</sup> | Jean-Luc Diehl<sup>2,18</sup> | Tristan Mirault<sup>1</sup> | David M. Smadja<sup>2,3</sup>  

<sup>1</sup>Vascular Medicine Department and Biosurgical Research Lab (Carpentier Foundation), AP-HP, Georges Pompidou European Hospital, Université de Paris, Paris, France

<sup>2</sup>Innovative Therapies in Haemostasis, INSERM, Université de Paris, Paris, France

<sup>3</sup>Hematology Department and Biosurgical Research Lab (Carpentier Foundation), AP-HP, Georges Pompidou European Hospital, Paris, France

<sup>4</sup>PARCC, INSERM, Université de Paris, Paris, France

<sup>5</sup>Emergency Department, AP-HP, Georges Pompidou European Hospital, Paris, France

<sup>6</sup>Intensive Care Unit, AP-HP, Georges Pompidou European Hospital, Université de Paris, Paris, France

<sup>7</sup>Virology Department, AP-HP, Georges Pompidou European Hospital, Paris, France

<sup>8</sup>Respiratory Medicine Department, AP-HP, Georges Pompidou European Hospital, Université de Paris, Paris, France

<sup>9</sup>Infectious Disease Department, AP-HP, Georges Pompidou European Hospital, Université de Paris, Paris, France

<sup>10</sup>Plastic Surgery Department, AP-HP, Georges Pompidou European Hospital,

## Abstract

**Background:** Coronavirus disease-2019 (COVID-19) has been associated with cardiovascular complications and coagulation disorders.

**Objectives:** To explore the coagulopathy and endothelial dysfunction in COVID-19 patients.

**Methods:** The study analyzed clinical and biological profiles of patients with suspected COVID-19 infection at admission, including hemostasis tests and quantification of circulating endothelial cells (CECs).

**Results:** Among 96 consecutive COVID-19-suspected patients fulfilling criteria for hospitalization, 66 were tested positive for SARS-CoV-2. COVID-19-positive patients were more likely to present with fever ( $P = .02$ ), cough ( $P = .03$ ), and pneumonia at computed tomography (CT) scan ( $P = .002$ ) at admission. Prevalence of D-dimer  $>500$  ng/mL was higher in COVID-19-positive patients (74.2% versus 43.3%;  $P = .007$ ). No sign of disseminated intravascular coagulation were identified. Adding D-dimers  $>500$  ng/mL to gender and pneumonia at CT scan in receiver operating characteristic curve analysis significantly increased area under the curve for COVID-19 diagnosis. COVID-19-positive patients had significantly more CECs at admission ( $P = .008$ ) than COVID-19-negative ones. COVID-19-positive patients treated with curative anticoagulant prior to admission had fewer CECs ( $P = .02$ ) than those without. Interestingly, patients

Lina Khider and Nicolas Gendron contributed equally to this work.

Manuscript handled by: Alan Mast

Final decision: Alan Mast, 08 June 2020

© 2020 International Society on Thrombosis and Haemostasis

Université de Paris, Paris, France

<sup>11</sup>Centre de Recherche des Cordeliers, Functional Genomics of Solid Tumors, INSERM, Université de Paris, Paris, France

<sup>12</sup>Department of Medical Informatics, AP-HP, Georges Pompidou European Hospital, Université de Paris, Paris, France

<sup>13</sup>Hematology Department, AH-HP, Georges Pompidou European Hospital, Paris, France

<sup>14</sup>Pathology Department and PRB (Plateforme de ressources biologiques), AP-HP, Georges Pompidou European Hospital, Paris, France

<sup>15</sup>Emergency Department, AP-HP, Georges Pompidou European Hospital, Université de Paris, Paris, France

<sup>16</sup>Respiratory Medicine Department and Biosurgical Research Lab (Carpentier Foundation), AH-HP, Georges Pompidou European Hospital, Paris, France

<sup>17</sup>Clinical Center of Investigation, AP-HP, Georges Pompidou European Hospital, Paris, France

<sup>18</sup>Intensive Care Unit and Biosurgical Research Lab (Carpentier Foundation), AH-HP, Georges Pompidou European Hospital, Paris, France

#### Correspondence

David M. Smadja, Hematology Department and Biosurgical Research Lab (Carpentier Foundation), AH-HP, Georges Pompidou European Hospital, 20 rue Leblanc, 75015 Paris, France.  
Email: david.smadja@aphp.fr

#### Funding Information

This work is supported by grants from ANR SARCODO (Fondation de France) and GHU APHP.CUP with Appel d'offre mécénat "collecte crise covid".

treated with curative anticoagulation and angiotensin-converting-enzyme inhibitors or angiotensin receptor blockers had even fewer CECs ( $P = .007$ ).

**Conclusion:** Curative anticoagulation could prevent COVID-19-associated coagulopathy and endothelial lesion.

#### KEYWORDS

circulating endothelial cells, coagulopathy, COVID-19, D-dimers, SARS-CoV-2

## 1 | INTRODUCTION

In December 2019, an epidemic pneumonia caused by a new coronavirus occurred in Wuhan (Hubei Province) and spread rapidly throughout China, evolving into a global pandemic.<sup>1</sup> Originally called new coronavirus 2019 (2019-nCoV), this virus was then officially named severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) by the World Health Organization (WHO). On January 30, 2020, the WHO declared the SARS-CoV-2 epidemic (COVID-19) a public health emergency of international concern. Compared to SARS-CoV, which caused a severe acute respiratory syndrome (SARS) epidemic in 2003, SARS-CoV-2 has a higher transmission capacity with a higher mortality.<sup>2</sup> With a rapid increase in confirmed cases, there is an unmet need of prevention and therapeutic strategies of COVID-19. Although COVID-19's clinical manifestations are dominated by respiratory symptoms, some patients have severe cardiovascular damage<sup>3,4</sup> and kidney disease contributing to multiple

#### ESSENTIALS

- Coronavirus disease-2019 (COVID-19) has been associated with coagulopathy and endotheliitis.
- Coagulopathy and endothelial dysfunction in COVID-19 patients at admission need to be precisely determined.
- Adding D-dimers to gender and pneumonia at computed tomography scan significantly increased specificity of COVID-19 diagnosis.
- Curative anticoagulation could prevent COVID-19-associated endothelial lesion.

organ failure. Patients with cardiovascular comorbidities may also be at higher risk of death.<sup>4</sup> For example, COVID-19 patients with hypertension have an increased mortality and morbidity with hazard ratio

ranging from 1.7 to 3.05, depending on the study.<sup>3,5</sup> Based on this, rising concerns have emerged regarding angiotensin-converting-enzyme inhibitors (ACEi) and angiotensin receptor blockers (ARBs). Indeed, angiotensin-converting enzyme 2 (ACE2) has been shown to be a co-receptor for viral entry for SARS-CoV-2, and it has been demonstrated that ACEi and ARBs could enhance ACE2 expression,<sup>6</sup> which may therefore jeopardize patient susceptibility to viral host cell entry and dissemination.

During the epidemic in China, coagulopathy was reported in severe COVID-19 patients. D-dimer levels above 1000 ng/mL were an independent risk factor of in-hospital death.<sup>5</sup> Coagulopathy was also found in fatal cases of COVID-19 patients, including a significantly higher proportion of patients with D-dimers above 500 ng/mL and prolonged prothrombin time (PT) in non-survivors.<sup>7</sup> This study was replicated in a second Chinese population in which D-dimers were still associated with in-hospital mortality.<sup>8</sup> The hypothesis of micro-thrombi in kidneys was also suggested in COVID-19 patients because high creatinine level was correlated with D-dimers above 500 ng/mL.<sup>9</sup> Finally, endothelial dysfunction might also play a role in the incidence of severe respiratory symptoms and viral systemic dissemination.<sup>10</sup> Indeed, the SARS-CoV-2 receptor ACE2 is strongly expressed on endothelial cells.<sup>11</sup> We hypothesized that endothelial cell infection may induce endothelium damage and dysfunction/activation that triggers coagulation activation.<sup>12</sup> Circulating endothelial cells (CECs) are considered relevant markers of endothelial lesion or dysfunction<sup>13</sup> and were used to explore the potential vascular dysfunction in COVID-19 patients.

The aim of our study was to identify biological markers related to COVID-19 diagnosis and severity. We also aimed at better characterizing subpopulations of patients at risk of coagulopathy and/or endothelial dysfunction to target COVID-19 patients at risk for the worst outcomes.

## 2 | METHODS

### 2.1 | Study design and population

This is a single-center, prospective observational cohort study conducted in a university hospital in Paris (France). From March 14, 2020 to March 20, 2020, all consecutive patients aged over 18 years, presenting to the emergency room of the Georges Pompidou European hospital and fulfilling hospitalization criteria or direct in-patient referral, with an infectious syndrome suspect of COVID-19 were included. Hospitalization criteria were based on local guidelines as described in Table 1. Suspicion of COVID-19 was defined by the presence of at least one of the following: fever, headache, myalgia, cough, dyspnea, rhinorrhea, or digestive symptoms. All COVID-19-suspected patients had a clinical evaluation, blood tests, and computed tomography (CT) scan, and were tested for SARS-CoV-2 infection by nasopharyngeal swab before they were transferred to dedicated hospitalization units: medical department or intensive care unit (ICU). The study was performed

**TABLE 1** Hospitalization criteria for COVID-19-suspected patients

Hospitalization criteria for COVID-19 suspected-patients

Co-morbidities other than respiratory failure **AND** requiring supplemental oxygen <3L/min to obtain SpO<sub>2</sub> >96%.

Respiratory failure **OR** dyspnea **OR** patients requiring supplemental oxygen >3L/min to obtain SpO<sub>2</sub> >96%

Predominant **AND/OR** decompensated cardiovascular comorbidities

Abbreviation: SpO<sub>2</sub>, oxygen saturation.

in accordance with the Declaration of Helsinki. All patients provided written informed consent before enrollment (CPP 2020-04-048/ 2020-A01048-31/ 20.04.21.49318). For all patients, baseline characteristics (demographic, treatment, clinical, cardiovascular risk factors, and body mass index), biological data, and CT scan results were retrieved from the medical records using standardized data collection.

### 2.2 | Laboratory confirmation of SARS-CoV-2 infection

Nasopharyngeal swabs were collected in universal transport medium (Xpert® nasopharyngeal sample collection kit). SARS-CoV-2 was detected using Allplex™ 2019-nCoV Assay (Seegene), a multiplex real-time polymerase chain reaction (PCR) assay that detects three target genes (E gene, RdRP gene, and N gene) in a single tube. Data were automatically analyzed using Seegene viewer software. Only qualitative data were available.

### 2.3 | Routine blood examinations

All samples were collected on ethylenediaminetetraacetic acid (EDTA), sodium heparin, and 0.129 M trisodium citrate tubes (9NC BD Vacutainer, Plymouth, UK). Routine lab tests were complete blood count, creatinine, C-reactive protein (CRP), and high-sensitive troponin I (Hs-TnI). Coagulation tests were PT ratio, fibrinogen, soluble fibrin monomer (STA®-Liatest FM; Stago), and antithrombin levels (Stachrom® AT III 6, Stago) explored on a STA-R® Max (Stago) coagulometer as previously described.<sup>14</sup> D-dimers concentrations were determined using the Vidas D-Dimer assay (BioMérieux) according to the manufacturer's instructions.

### 2.4 | CECs quantification

Peripheral venous blood samples were collected on EDTA after having always discarded the first milliliter of blood to avoid presence of endothelial cells dislodged by puncture. CECs were isolated by immunomagnetic separation with mAb CD146-coated beads and staining with the fluorescent probe acridin orange as previously described.<sup>13,15-17</sup>

## 2.5 | Statistical analysis

Continuous data were expressed as median [interquartile range: (IQR)] and categorical data as proportion. In the univariate analysis, we determined differences in median using the unpaired t-test (Mann-Whitney U test) for continuous variables and in proportions we were using the Chi-square test or Fischer exact test if necessary. We generated receiver operating characteristics (ROC) curve with the regression models that included variables with a significant difference in the univariate analysis.<sup>18,19</sup> The model included gender, pneumonia at CT scan, and D-dimers above 500 ng/mL. This model helped assess the extent to which the level of D-dimers influenced the predictability of COVID-19 diagnosis. We compared area under the curve (AUC) of each ROC curve using the Delong test. All analyses were two-sided and a *P*-value of *P* < .05 was considered statistically significant. Statistical analysis was performed using R studio software (R Foundation for Statistical Computing).

## 3 | RESULTS

### 3.1 | High level of D-dimers is a discriminant factor during COVID-19 suspicion

Among the 96 COVID-19-suspected patients included, 66 were positive for SARS-CoV-2. COVID-19-positive patients were more likely to be males (*N* = 44, 66.7%) than COVID-19-negative patients (*N* = 13, 43.3%; *P* = .05). Otherwise, the two populations were strictly comparable in terms of time from illness onset to hospital admission, age, body mass index, cardiovascular risk factors, medical history, and treatments (Table 2). Considering clinical features at admission, COVID-19-positive patients were more likely to have fever (*P* = .02), cough (*P* = .03), and interstitial pneumonia at CT scan (*P* = .002). In terms of biological features (Table 3) COVID-19-positive patients had a significantly lower white blood cell count, including neutrophil count (respectively, *P* = .008 and 0.02). Regarding hemostasis, the proportion of COVID-19-positive patients with D-dimers above 500 ng/mL was significantly higher (74.2% versus 43.3%; *P* = .007). No difference in PT ratio or fibrin monomers was observed between groups. In the context of COVID-19-associated coagulopathy with high levels of D-dimers, fibrinogen, and CRP at admission, the low level of fibrin monomers and normal antithrombin levels allowed us to exclude a disseminated intravascular coagulation (DIC). When adding D-dimers above 500 ng/mL to gender and pneumonia at CT scan, ROC curve area (Figure 1) significantly increased from AUC 0.73 (95% confidence interval [CI] 0.61-0.85) to AUC 0.82 (95% CI 0.69-0.95; *P* = .02), and confirm the relevance of D-dimers in diagnosis of COVID-19 at admission in hospital. The ROC curve with D-dimers above 500 ng/mL yielded a high sensitivity of 98.1% (95% CI 50.0-100.0), a low specificity of 28.6% (95% CI 21.0-85.8), a high positive predictive value of 77.9% (95% CI 65.3-90.0), and a high negative predictive value of 85.7% (95% CI 55.0-100.0).

### 3.2 | Anticoagulated COVID-19-positive patients have a significant lower CECs count

CECs were quantified as markers of endothelial lesion, using the reference method<sup>15</sup> as detailed in the Method section.<sup>13</sup> Using this assay, the upper limit of normal range at 10 CECs per mL of whole blood was previously determined and confirmed in several studies, including ours.<sup>13,20-22</sup> Among COVID-19-positive patients, 64% were above this threshold, suggesting a SARS-CoV-2-induced endothelial lesion. For comparison, COVID-19 negative population had fewer CECs (*P* = .008; Table 3) with only 27% above the normal range (*P* = .012). Because the coagulopathy observed in COVID-19 patients could be related to this endothelial lesion, we analyzed whether anticoagulation could impact CEC level. Patients treated with curative anticoagulation prior to admission for any medical reason (83% for atrial fibrillation; 17% for venous thromboembolism) had indeed a lower CEC level than those without curative anticoagulation: 9 [8, 17] versus 24 [14, 42] CECs per mL (*P* = .02), respectively (Figure 2). Interestingly, in patients treated with ACEi or ARBs the effect of curative anticoagulation on the CEC level was more pronounced with 10 [5, 14] versus 30 [17, 43] CECs per mL (*P* = .007) in those without curative anticoagulation (Figure 2).

## 4 | DISCUSSION

The originality of this study was to evidence an endothelial lesion during SARS-CoV-2 infection, as witnessed by increased levels of CECs. Second, we show that this endothelial damage is thwarted by curative anticoagulation.

Several Chinese studies found that increased D-dimer level correlated with in-hospital mortality,<sup>5,7,8</sup> suggesting a COVID-19-associated DIC.<sup>7,8,23</sup> However, in our population, no overt DIC was diagnosed at admission. Indeed, patients had no significant thrombocytopenia, a normal PT ratio, and high fibrinogen levels. This was confirmed by a low level of fibrin monomers, which are early markers of DIC. DIC might nevertheless be involved in patient worsening and in particular for those with acute respiratory distress syndrome. Therefore, the increase of D-dimers largely reported in COVID-19 patients is probably not related to DIC but might reflect the microthrombi formation. Indeed, histopathological observations and imaging features of pulmonary lesions associated with SARS-CoV-2 revealed intra-alveolar fibrin deposit in pulmonary samples.<sup>24</sup> Moreover, a recent study using standardized protective ventilation settings in COVID-19 patients confirmed that oxygenation was severely compromised with a moderate alteration in the respiratory system compliance. Hypercapnia high prevalence led to the hypothesis of a large amount of ventilated/not perfused alveoli, which could reflect diffuse microthrombi in the pulmonary microvascular bed.<sup>25</sup> In renal disease associated to COVID-19, thrombotic lesions were proposed, because high D-dimers were more commonly observed in patients with elevated baseline serum creatinine.<sup>9</sup> In this context, the International Society on Thrombosis and Haemostasis (ISTH) has

**TABLE 2** Demographic, clinical, and treatment characteristics of patients on admission according to COVID-19 viral status

	COVID-19 negative n = 30	COVID-19 positive n = 66	P- value
Male sex, n (%)	13 (43.3)	44 (66.7)	.05
Age, years, median [IQR]	63.0 [55.3, 75.8]	66.0 [54.3, 79.8]	.690
BMI, kg/m <sup>2</sup> , median [IQR]	24.8 [22.6, 26.3]	26.5 [24.7, 29.1]	.110
Time from illness onset to hospital admission, days, median [IQR]	4.47 (4.57)	5.45 (3.71)	.260
CV risk factors, n (%)			
Hypertension	16 (53.3)	31 (47.0)	.720
Dyslipidemia	6 (20.0)	21 (31.8)	.340
Diabetes	4 (13.3)	12 (18.2)	.760
Sedentarity	6 (20.0)	6 (9.1)	.260
Chronic kidney disease	4 (13.3)	8 (12.1)	1.000
Medical history, n (%)			
Cancer	6 (20.0)	6 (9.1)	.240
Coronary heart disease	3 (10.0)	7 (10.6)	.920
Stroke	0 (0.0)	4 (6.1)	.400
treatments, n (%)			
Statins	3 (10.0)	13 (19.7)	.220
Oral antidiabetic agents	2 (6.7)	9 (13.6)	.510
Insulin	2 (6.7)	5 (7.6)	1.000
β-blockers	5 (16.7)	8 (12.1)	.770
Calcium channel blockers	7 (23.3)	13 (19.7)	.890
ACEi or ARBs	9 (30.0)	21 (31.8)	1.000
Diuretics	3 (10.0)	6 (9.1)	1.000
Central acting agent	1 (3.3)	0 (0.0)	.680
Curative anticoagulation	3 (10.0)	12 (18.2)	.470
Clinical features, n (%)			
Fever	22 (73.3)	61 (92.4)	.020
Headache	1 (3.3)	12 (18.2)	.090
Cough	14 (46.7)	47 (71.2)	.030
Productive cough	3 (10.0)	8 (12.1)	1.000
Dyspnea	9 (30.0)	31 (47.0)	.210
Myalgia	5 (16.7)	21 (31.8)	.190
Diarrhea	2 (6.7)	7 (10.6)	.640
Pneumonia at CT scan	11 (36.7)	48 (72.7)	.002
ARDS	1 (3.3)	9 (13.6)	.240
SpO <sub>2</sub> %, median [IQR]	96.0 [92.0, 98.0]	95.0 [91.0, 96.0]	.050
Respiratory rate, breaths per minute, median [IQR]	20.0 [16.5, 25.0]	19.0 [16.0, 22.8]	.550
Pulse, beats per min, median [IQR]	87.0 [74.0, 100.0]	87.0 [74.5, 103.5]	.850

Abbreviations:: ACEi, angiotensin conversion enzyme inhibitor; ARBs, angiotensin 2 receptor blocker; ARDS, acute respiratory distress syndrome; BMI, body mass index; CV, cardiovascular; IQR, interquartile range; SpO<sub>2</sub>, oxygen saturation.

recently recommended measuring D-dimers, PT ratio, and platelet count in all COVID-19 patients to help stratify those who may benefit from hospitalization and a close monitoring.<sup>26</sup> Therefore, ISTH, the American College of Cardiology, and the French Society for Vascular Medicine suggested the use of prophylactic anticoagulation with low weight molecular heparin (LMWH) for COVID-19 patients, in the absence of any contraindications.<sup>26-28</sup> Indeed, preventive LMWH


treatment could be associated with better prognosis in severe COVID-19 patients meeting sepsis-induced coagulopathy criteria.<sup>29</sup>

We further hypothesized that COVID-19-induced coagulopathy could be a consequence of endothelial injury, based on the rationale that SARS-CoV-2 has an endothelial tropism linked to ACE2 expression. Moreover, recently an endotheliitis has been described in SARS-CoV-2 infection and could be at the origin of impaired

	COVID-19 negative n = 30	COVID-19 positive n = 66	P- value
White blood cells, $\times 10^9$ per L, median [IQR]	7.8 [6.1, 11.4]	6.0 [4.6, 7.4]	.008
Hemoglobin, g/L, median [IQR]	126.0 [110.3, 141.0]	130.5 [112.0, 144.0]	.490
Platelet count, $\times 10^9$ per L, median [IQR]	217.5 [157.0, 278.3]	167.5 [146.3, 223.0]	.090
Neutrophils, $\times 10^9$ per L, median [IQR]	5.7 [4.2, 9.0]	4.0 [3.0, 5.9]	.020
Lymphocytes, $\times 10^9$ per L, median [IQR]	1.0 [0.8, 1.7]	0.9 [0.7, 1.3]	.170
Monocytes, $\times 10^9$ per L, median [IQR]	0.6 [0.4, 0.8]	0.4 [0.3, 0.6]	.300
CRP, mg/L, median [IQR]	55.6 [3.3, 127.2]	74.0 [22.7, 126.3]	.210
Plasma creatinine level, $\mu\text{mol/L}$ , median [IQR]	99.0 [55.0, 112.0]	79.0 [62.7, 108.7]	.970
Hs-TNI, pg/mL, median [IQR]	10.5 [3.5, 32.2]	9.5 [5.1, 22.9]	.740
PT ratio, median [IQR]	0.94 [0.70, 1.10]	0.95 [0.86, 1.00]	.320
Fibrinogen, g/L, median [IQR]	5.1 [4.3, 5.8]	5.3 [4.7, 6.2]	.420
D-dimers > 500 ng/mL, n (%)	13 (43.3)	49 (74.2)	.007
Fibrin monomers, $\mu\text{g/mL}$ , median [IQR]	7.0 [7.0, 7.0]	7.0 [7.0, 7.0]	.550
Antithrombin, %, median [IQR]	99.0 [85.3, 102.8]	101.0 [86.5, 106.5]	.500
CECs per mL, median [IQR]	9 [6, 18]	19 [10, 39]	.008
CECs $\geq 10$ per mL, n (%)	8 (27)	42 (64)	.012

**TABLE 3** Biological parameters of patients on admission according to COVID-19 viral status


Abbreviations: CECs, circulating endothelial cells; CRP, C-reactive protein; Hs-Tni, high-sensitive troponin I; IQR, interquartile range; PT, thromboplastin time.



**FIGURE 1** Potential exclusion criteria for COVID-19 diagnosis. The receiver operating characteristic (ROC) curve including gender and pneumonia with (red line) or without D-dimers threshold at 500 ng/mL (blue line). ROC curve analysis identified the association of female gender, absence of pneumonia at computed tomography scan, and D-dimers below or equal 500 ng/mL as potential exclusion criteria for COVID-19 diagnosis (area under the curve 81.9% (95% IC 68.7%–95.1%))

microcirculatory function affecting particularly the lungs and kidneys.<sup>30</sup> Thus, we explored CECs as a recognized non-invasive marker for endothelial lesion, as demonstrated in acute cardiovascular

conditions such as acute coronary syndrome<sup>31</sup> and pulmonary arterial hypertension.<sup>13,16,20</sup> In agreement with the consensus protocol from ISTH, we used the reference method for CEC quantification and the threshold of 10 CECs per mL as upper normal value.<sup>15</sup> We found that more than 60% of COVID-19-positive patients were above this threshold. Interestingly, patients enrolled while they were treated with curative anticoagulation had a significantly lower level of CECs, especially in the hypertensive population treated with ACEi or ARBs. Increased mortality and/or morbidity of COVID-19 in patients with hypertension has been described in China.<sup>3</sup> One of the most important concerns is the association between hypertension and treatment with ACEi or ARBs. Indeed, because ACE2 is a receptor for viral entry of SARS-CoV-2, a link between ACEi or ARBs was considered. ACEi or ARBs were described to increase ACE2 expression in the heart, brain, and even in urine after treatment.<sup>6,32</sup> However, the main scientific societies of cardiology and more specifically of hypertension took the position not to withdraw ACEi or ARB therapy in COVID-19. Hence, the Council on Hypertension of the European Society of Cardiology recommended that "patients should continue treatment with their usual anti-hypertensive therapy because there is no clinical or scientific evidence to suggest that treatment with ACEi or ARBs should be discontinued because of the COVID-19 infection."<sup>33</sup> Indeed, recent studies did not find any association between ACEi or ARBs therapy and worsening in COVID-19 patients.<sup>34</sup>




**FIGURE 2** Effect of curative anticoagulation on circulating endothelial cell (CEC) levels in COVID-19. Quantification of CECs in COVID-19-positive patients at admission. CEC level according to presence or the absence of curative anticoagulation and/or the presence or the absence of angiotensin-converting enzyme inhibitors (ACEi) or angiotensin receptor blockers (ARBs). Red dotted line shows the upper limit of reference values for CECs (<10 CECs per mL)

In our population patients treated with curative anticoagulation had a lower level of CECs, so we hypothesized that curative anticoagulation in COVID-19 patients could decrease thrombotic risk and subsequent mortality. To our knowledge, this is the first time that a potential protective effect of anticoagulant therapy on endothelial dysfunction is described (Figure 3). Besides the coagulopathy associated to endothelial lesion, the effect of anticoagulation may act directly on SARS-CoV-2 entrance in endothelial cells. Indeed, cell entry of SARS-CoV-2 depends on the binding of the viral spike (S) proteins to cellular receptors and on S protein priming by host cell proteases. It has been demonstrated that serine protease TMPRSS2 is necessary for S protein priming.<sup>35</sup> Thus, a TMPRSS2 inhibitor has been proposed as a treatment option. If the virus's entrance into cells is dependent on a serine protease action, anticoagulation by inhibiting thrombin activity and serine proteases of the coagulation cascade could directly limit virus entrance in endothelial but also in other cells. This hypothesis needs to be tested in preclinical models of infection.

#### 4.1 | Study limitations

Our study has several limitations. First, we are aware that false COVID-19-negative patients may exist in our study population due

Endothelial injury in relation with SARS-CoV-2 infection



**FIGURE 3** Curative anticoagulant treatment could be part of COVID-19 management in order to limit associated endothelial dysfunction

to the imperfect sensitivity of the diagnostic test currently used.<sup>36</sup> Second, the study population size was necessarily small given the emergency of understanding COVID-19; indeed we decided to investigate the endothelial dysfunction associated to COVID-19 in order to consider new therapeutic alternatives. In addition, due to this small population, our results concerning anticoagulant treatment and ACEi or ARBs therapy generated hypotheses that need to be validated in larger cohorts.

In conclusion, it therefore seems consistent to open the way to curative anticoagulant treatment as part of the management of COVID-19 patients in order to limit associated coagulopathy and endothelial dysfunction (Figure 3). Anticoagulation may not only modify COVID-19-associated coagulopathy but also pathophysiology of SARS-CoV-2 systemic dissemination. Curative anticoagulation could decrease mortality observed in COVID-19 patients. Further studies should evaluate safety and efficacy of curative anticoagulation in COVID-19 patients to prevent worsening of disease and reduction of admittance in ICUs.

#### ACKNOWLEDGMENTS

We would like to acknowledge all nurses, technicians, and physicians involved in the vascular medicine, internal medicine, respiratory medicine, intensive care, clinical investigation center, and hematology departments of the George Pompidou European Hospital and Cochin Hospital for their help in taking care of patients and including them in the study. We thank AP-HP for promotion of the SARCODO Project. We thank the unit of clinical research URC HEGP CIC-EC1418 (Natacha Nohile, Pauline Jouany and Dr Juliette

Djadi-Prat) and Helene Cart-Grandjean from AP-HP for their involvement in the SARCODO Project. We would also like to acknowledge Laurent Garcia, Florence Desvard, Yann Burnel, Julie Bricet, and Nadège Ochat for specific technical organization in the hematology department.

## AUTHOR CONTRIBUTIONS

D. M. Smadja, T. Mirault, and J.-L. Diehl interpreted data, and conceived and supervised the study. L. Khider and N. Gendron interpreted data and drafted the manuscript. R. Chocron analyzed the data and supervised statistical analysis. L. Khider, N. Gendron, G. Goudot, R. Chocron, and B. Debuc analyzed the data and reviewed all patients' characteristics. All authors interpreted data, drafted and revised the manuscript, and approved the final version.

## CONFLICTS OF INTEREST

All the authors have nothing to disclose.

## ORCID

Nicolas Gendron  <https://orcid.org/0000-0003-4852-4738>

David M. Smadja  <https://orcid.org/0000-0001-7731-9202>

## TWITTER

David M. Smadja  @DavidMSmadja

## REFERENCES

- Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. 2020. <https://doi.org/10.1056/NEJMoa2002032>.
- van Doremalen N, Bushmaker T, Morris DH, et al. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *N Engl J Med*. 2020. <https://doi.org/10.1056/NEJMc2004973>.
- Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China. *JAMA*. 2020;323(13):1239.
- Zheng YY, Ma YT, Zhang JY, Xie X. COVID-19 and the cardiovascular system. *Nat Rev Cardiol*. 2020;17(5):259-260.
- Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395(10229):1054-1062.
- Ferrario CM, Jessup J, Chappell MC, et al. Effect of angiotensin-converting enzyme inhibition and angiotensin II receptor blockers on cardiac angiotensin-converting enzyme 2. *Circulation*. 2005;111(20):2605-2610.
- Tang N, Li D, Wang X, Sun Z. Abnormal Coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost*. 2020;18(4):844-847.
- Han H, Yang L, Liu R, et al. Prominent changes in blood coagulation of patients with SARS-CoV-2 infection. *Clin Chem Lab Med*. 2020;58(7):1116-1120.
- Cheng Y, Luo R, Wang K, et al. Kidney disease is associated with in-hospital death of patients with COVID-19. *Kidney Int*. 2020. In press. <https://doi.org/10.1016/j.kint.2020.03.005>.
- Smadja DM, Guerin CL, Chocron R, et al. Angiopietin-2 as a Marker of Endothelial Activation Is a Good Predictor Factor for Intensive Care Unit Admission of COVID-19 Patients. *Angiogenesis*. 2020;1-10. <https://doi.org/10.1007/s10456-020-09730-0>.
- Hamming I, Timens W, Bulthuis ML, Lely AT, Navis G, van Goor H. Tissue distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS pathogenesis. *J Pathol*. 2004;203(2):631-637.
- Goon PK, Boos CJ, Lip GY. Circulating endothelial cells: markers of vascular dysfunction. *Clin Lab*. 2005;51:531-538.
- Smadja DM, Gaussem P, Mauge L, et al. Circulating endothelial cells: a new candidate biomarker of irreversible pulmonary hypertension secondary to congenital heart disease. *Circulation*. 2009;119(3):374-381.
- Latremouille C, Carpentier A, Leprince P, et al. A bioprosthetic total artificial heart for end-stage heart failure: results from a pilot study. *J Heart Lung Transplant*. 2018;37(1):33-37.
- Woywodt A, Blann AD, Kirsch T, et al. Isolation and enumeration of circulating endothelial cells by immunomagnetic isolation: proposal of a definition and a consensus protocol. *J Thromb Haemost*. 2006;4(3):671-677.
- Levy M, Bonnet D, Mauge L, Celermajer DS, Gaussem P, Smadja DM. Circulating endothelial cells in refractory pulmonary hypertension in children: markers of treatment efficacy and clinical worsening. *PLoS One*. 2013;8:e65114.
- Smadja DM, Mauge L, Nunes H, et al. Imbalance of circulating endothelial cells and progenitors in idiopathic pulmonary fibrosis. *Angiogenesis*. 2013;16(1):147-157.
- Hajian-Tilaki K. Receiver operating characteristic (ROC) curve analysis for medical diagnostic test evaluation. *Caspian J Intern Med*. 2013;4:627-635.
- Park SH, Goo JM, Jo CH. Receiver operating characteristic (ROC) curve: practical review for radiologists. *Korean J Radiol*. 2004;5:11-18.
- Smadja DM, Mauge L, Sanchez O, et al. Distinct patterns of circulating endothelial cells in pulmonary hypertension. *Eur Respir J*. 2010;36:1284-1293.
- Sabatier F, Camoin-Jau L, Anfosso F, Sampol J, Dignat-George F. Circulating endothelial cells, microparticles and progenitors: key players towards the definition of vascular competence. *J Cell Mol Med*. 2009;13:454-471.
- Mauge L, Sabatier F, Boutouyrie P, et al. Forearm ischemia decreases endothelial colony-forming cell angiogenic potential. *Cytotherapy*. 2014;16:213-224.
- Lillicrap D. Disseminated intravascular coagulation in patients with 2019-nCoV pneumonia. *J Thromb Haemost*. 2020;18:786-787.
- Zhang H, Zhou P, Wei Y, et al. Histopathologic changes and SARS-CoV-2 immunostaining in the lung of a patient with COVID-19. *Ann Intern Med*. 2020;172(9):629-632.
- Liu X, Liu X, Xu Y, et al. Ventilatory ratio in hypercapnic mechanically ventilated patients with COVID-19 associated ARDS. *Am J Respir Crit Care Med*. 2020. doi:10.1164/rccm.202002-0373LE.
- Thachil J, Tang N, Gando S, et al. ISTH interim guidance on recognition and management of coagulopathy in COVID-19. *J Thromb Haemost*. 2020;18(5):1023-1026.
- Bikdeli B, Madhavan MV, Jimenez D, et al. COVID-19 and thrombotic or thromboembolic disease: implications for prevention, antithrombotic therapy, and follow-up. *J Am Coll Cardiol*. 2020;75(23):2950-2973.
- Khider L, Soudet S, Laneelle D, et al. Proposal of the French Society of Vascular Medicine for the prevention, diagnosis and treatment of venous thromboembolic disease in outpatients with COVID-19. *J Méd Vascu*. 2020;45(4):210-213.
- Tang N, Bai H, Chen X, Gong J, Li D, Sun Z. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. *J Thromb Haemost*. 2020;18(5):1094-1099.
- Varga Z, Flammer AJ, Steiger P, et al. Endothelial cell infection and endotheliitis in COVID-19. *Lancet*. 2020;395(10234):1417-1418.



31. Quilici J, Banzet N, Paule P, et al. Circulating endothelial cell count as a diagnostic marker for non-ST-elevation acute coronary syndromes. *Circulation*. 2004;110(12):1586-1591.
32. Furuhashi M, Moniwa N, Mita T, et al. Urinary angiotensin-converting enzyme 2 in hypertensive patients may be increased by olmesartan, an angiotensin II receptor blocker. *Am J Hypertens*. 2015;28:15-21.
33. ESC. Position statement of the European Society of Cardiology (ESC) Council on Hypertension on ACE-inhibitors and angiotensin receptor blockers. [https://www.escardio.org/Councils/Council-on-Hypertension-\(CHT\)/News/position-statement-of-the-esc-council-on-hypertension-on-ace-inhibitors-and-ang](https://www.escardio.org/Councils/Council-on-Hypertension-(CHT)/News/position-statement-of-the-esc-council-on-hypertension-on-ace-inhibitors-and-ang) 2020; Published March 13, 2020. Accessed March 20
34. Fosbol EL. Association of Angiotensin-Converting Enzyme Inhibitor or Angiotensin Receptor Blocker Use With COVID-19 Diagnosis and Mortality. *JAMA*. 2020;e2011301. doi:10.1001/jama.2020.11301.
35. Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell*. 2020;181(2):271-280.
36. Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in different types of clinical specimens. *JAMA*. 2020.

**How to cite this article:** Khider L, Gendron N, Goudot G, et al. Curative anticoagulation prevents endothelial lesion in COVID-19 patients. *J Thromb Haemost*. 2020;18:2391-2399. <https://doi.org/10.1111/jth.14968>