

PIANO COVID-19 - Effect of organizational measures to prevent and control COVID-19 infection in nursing homes on the risk of death of residents during and after the epidemic period - PIANO COVID-19

Head :ROLLAND Yves

Last update : 04/23/2021 | Version : 1 | ID : 73934

General

Identification

Detailed name Effect of organizational measures to prevent and control COVID-19 infection in nursing homes on the risk of death of residents during and after the epidemic period - PIANO COVID-19

Sign or acronym PIANO COVID-19

General Aspects

Medical area Geriatrics

Study in connection with Covid-19 Yes

Scientific investigator(s) (Contact)

Name of the director ROLLAND

Surname Yves

Address Gérontopôle de Toulouse,
Cité de la Santé,
Place Lange
31059 Toulouse

Phone +33 (0)5 61 77 21 00

Organization GERIATRIC UNIT TOULOUSE UNIVERSITY HOSPITAL

Collaborations

Participation in projects, networks and consortia Yes

Funding

Funding status Public

Details DGOS (General Directorate of Healthcare Supply) funding

Governance of the database

Sponsor(s) or organisation(s) responsible Toulouse University Hospital

Organisation status Public

Presence of scientific or steering committees No

Additional contact

Name of the contact PENNETIER

Surname Delphine

Address Gérontopôle de Toulouse,
Cité de la Santé,
Place Lange
31059 Toulouse

Phone 05 61 77 70 11

Email pennetier.d@chu-toulouse.fr

Organization GERIATRIC UNIT - TOULOUSE UNIVERSITY HOSPITAL

Main features

Type of database

Type of database Study databases

Study databases (details) Cohort study

Database recruitment is carried out by an intermediary A selection of health institutions and services

Database recruitment is carried out as part of an interventional study No

Database objective

Main objective The main objective is to compare the occurrence of global death of residents during a 1-year period (from January 2020 to December 2020) in NH and LTCU with a high level of implementation of the recommendation/guidance to prevent and control COVID-19 NH/LTCU residents' infection and NH and LTCU with a low level of implementation of the recommendation/guidance to prevent and control COVID-19 NH/LTCU.

The secondary objectives are:

- To compare the rate of death related to confirmed or suspected COVID cases in NH/LTCU with a high level of implementation of the recommendation/guidance to prevent and control COVID-19 NH/LTCU residents' infection and NH/LTCU with a low level of implementation of the recommendation/guidance to prevent and control COVID-19 NH/LTCU.
- To compare the rate of death not related to COVID cases in NH/LTCU with a high level of implementation of the recommendation/guidance to prevent and control COVID-19 NH/LTCU residents' infection and NH/LTCU with a low level of implementation of the recommendation/guidance to prevent and control COVID-19 NH/LTCU.
- To analyze the incidence of serious health events (deaths, hospitalizations) and the occurrence of COVID in residents (vaccinated and unvaccinated), in NHs and LTCUs in France during a 12-month (2021) observation period.
- To analyze the occurrence of COVID in residents (vaccinated and unvaccinated), in NHs and LTCUs in France according to the rate of vaccination against COVID among the NH's/LTCU's staff during the same observational period.

From an economic perspective:

- To assess the economic impact of the implementation of prevention measures applied by nursing homes staff, from healthcare system and the NH perspectives, 6, and 12 months before and 6, 12 months to 24 months after the implementation of the preventive measures.
- To assess the economic impact of the vaccination among residents and NH's staff, from healthcare system and the NH perspectives, 6 and 12 months before and after the vaccination campaign. These data will also provide indirect economic information about the tolerance of the vaccination.
- To assess the efficiency of the high level implementation of the recommendation to prevent and control COVID-19 in comparison with the low level implementation of recommendation to prevent and control COVID-19, using a cost-effectiveness analysis at 12 and 24 months.
- To assess the efficiency of the vaccination, at 6 and 12 months, among residents and healthcare professionals in comparison with no vaccination, using a cost-effectiveness analysis

Inclusion criteria

NHs and LTCU volunteers to participate from different region of France (from low impacted to

highly impacted region) will be welcomed.
NON-INCLUSION CRITERIA
NHs or LTCUs that refuse to participate.

Population type

Age
Elderly (65 to 79 years)
Great age (80 years and more)

Population covered
General population

Pathology

Gender
Male
Woman

Geography area
National

Data collection

Dates

Date of first collection (YYYY or MM/YYYY)
2020

Date of last collection (YYYY or MM/YYYY)
2021

Size of the database

Size of the database (number of individuals)
[500-1000[individuals

Details of the number of individuals
9,600 persons from 120 nursing homes (coordinating nurses or coordinating physicians).

Data

Type of data collected
Clinical data
Cost data

Clinical data (detail)
Direct physical measures

Cost data (detail)
healthcare consumption data collected via the health insurance scheme

Presence of a biobank
No

Health parameters studied
Health event/mortality
Health care consumption and services

Care consumption (detail)
Hospitalization

Medical/paramedical consultation

Procedures

Data collection method

Excel file; CRF

Participant monitoring

Yes

Details on monitoring of participants

in 2021: vital status, GIR, and COVID vaccination data of participants will be collected

Promotion and access

Promotion

Access