

# EFICARE - TRANSVERSAL STUDY ON MARKERS AND RISK FACTORS ASSOCIATED WITH CARDIAC INSUFFICIENCY IN PATIENTS AGED OVER 70 YEARS FOLLOWED IN AMBULATORY MODE

Last update : 11/01/2019

Head :  
Hanon Olivier

Responsible organization :  
MENARINI France

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Describe the markers and risk factors associated with cardiac insufficiency in patients aged over 70 years followed in ambulatory mode. Assess the cognitive functions using Memory Impairment Score and assess the Senni prognostic index.

## Inclusion criteria

"Male or Female aged 70 years or older  
Having Chronic cardiac insufficiency  
Hospitalized for cardiac decompensation in the last 12 months"

# ELSA - TOULOUSE - LONGITUDINAL STUDY MONITORING ALZHEIMER'S DISEASE PATIENTS - TOULOUSE

Last update : 09/05/2017

Head :  
Andrieu Sandrine

Responsible organization :  
Centre Hospitalier Universitaire Toulouse

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

To the describe the risks of rapid weight loss in individuals with Alzheimer disease.

### **Inclusion criteria**

Men or women - between 45 and 89 years old - presenting symptoms of dementia - 5 kg weight loss in 6 months

## **SNAC - STUDY OF NORPREGNANES COAGULATION**

Last update : 09/05/2017

Head :

Responsible organization :

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Impact of different types of HRT on activated protein C resistance and haemostatic parameters.

### **Inclusion criteria**

- Women - between 45 and 70 years of age - menopausal

## **PRALIMAP - PROMOTION OF HEALTHY EATING AND PHYSICAL ACTIVITY**

Last update : 12/01/2019

Head :

Briançon Serge, ECOLE DE SANTÉ PUBLIQUE - NANCY UNIVERSITÉ LABORATOIRE EA4003

Responsible organization :

CONSERVATOIRE NATIONAL DES ARTS ET MÉTIERS (CNAM)

### **Type of database**

Study databases

### **Study databases (details)**

Cohort study

### **Main objective**

General objective: The general objective is to evaluate the impact of three intervention strategies, to develop individual skills, to identify and support those at risk, to create a favourable environment, alone or by association, regarding awareness, attitudes and behaviour of adolescents towards food and physical

activity and to measure corpulence using anthropometric markers. Secondary objectives: - To assess the feasibility of implementing each strategy in secondary schools; - To analyse the psychological and sociological factors involved in adherence and the success or failure of the implemented actions.

### **Inclusion criteria**

Students in administrative secondary schools in Meurthe-et-Moselle, Meuse, Moselle and Vosges.

## **GERICOTS - RELATIONSHIP BETWEEN REDUCTION IN WORKING TIME, WORK LIFE AND IMPACT ON HEALTH IN THE NORD-PAS-DE-CALAIS AND PICARDIE REGIONS**

Last update : 09/05/2017

Head :

Edme Jean-Louis, LABORATOIRE DE MÉDECINE DU TRAVAIL FACULTÉ DE MÉDECINE 1 PLACE DE VERDUN 59045 LILLE CEDEX UNIVERSITÉ DE LILLE 2

Frimat Paul, LABORATOIRE DE MÉDECINE DU TRAVAIL FACULTÉ DE MÉDECINE 1 PLACE DE VERDUN 59045 LILLE CEDEX UNIVERSITÉ DE LILLE 2

Responsible organization :

Université de Lille

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

To analyse the relationship between organisational factors, working hours and employee health in the Nord-Pas-de-Calais/Picardie regions. Secondary objectives: - Regional reference on stress levels at work; - Relationship between occupational stress and health; - Relationship between incidences of stress and health problems.

### **Inclusion criteria**

Employees with a permanent employment contract who understand French and are living in the Nord-Pas-de-Calais or Picardie region.

## **UTILE - INTÉRÊT ET USAGES DES ANTI-TNF ALPHA DANS LA PRISE EN CHARGE DU PSORIASIS DE PLAQUES**

Last update : 01/01/2019

Head :

Thibout Emmanuel, Laboratoire Abbott

Responsible organization :

Abbott France

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Know the profile of the patients followed for psoriasis

Specify the therapeutic schemas used and in particular the place of biotherapies

## Inclusion criteria

patients suffering from psoriasis

# THDA/TPL - ÉTUDE TRANSVERSALE SUR LES TROUBLES D'HYPERACTIVITÉ AVEC DÉFICIT DE L'ATTENTION (THDA) CHEZ LES ADOLESCENTS SOUFFRANT D'UN TROUBLE LIMITE DE LA PERSONNALITÉ (TPL)

Last update : 12/01/2017

Head :

Falissard Bruno, Inserm U1018 - Centre de Recherche en Épidémiologie et Santé des Populations Équipe:

Genre, Santé sexuelle et Reproductive

Revah-Levy Anne, Centre de Soins Psychothérapeutiques de Transition pour Adolescents

Speranza Mario , Service de Pédopsychiatrie.

Cortese Samuele

Pham-Scottez Alexandra

Corcos Maurice , Département de Psychiatrie de l'Adolescent et du Jeune Adulte

Responsible organization :

INSERM - Institut National de la Santé et de la Recherche Médicale

## Type of database

Bases de données issues d'enquêtes

## Study databases (details)

Etudes transversales non répétées (hors enquêtes cas-témoins)

## Main objective

1) évaluer la prévalence passée et actuelle de la comorbidité du diagnostic THDA/TPL et son impact sur la présentation clinique de trouble de la personnalité limite chez les adolescents

2) déterminer le type d'impulsivité qui caractérise spécifiquement les adolescents avec THDA/TPL.

## Inclusion criteria

Patients de cinq centres universitaires psychiatriques en France, en Belgique et en Suisse âgés de 15 à 19 ans avec un diagnostic de trouble limite de la personnalité selon les critères du DSM-IV.

Les critères d'exclusion: diagnostic de schizophrénie ou de maladie chronique et/ou médicale grave

impliquant le pronostic vital.

## **PERFECTA - INTRAVENOUS ANALGESICS FOR POST-OPERATIVE PAIN: PATTERNS OF USE IN HIGH-RISK POPULATIONS**

Last update : 01/01/2018

Head :

Boccard Éric, Bristol-Myers Squibb

Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :

Bristol-Myers Squibb France (BMS)

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Describe the post-surgical drug treatment for pain after a surgical intervention, in target groups of patients having specific risks of developing complications during the use of certain classes of analgesics

### **Inclusion criteria**

M/F patients,  $\geq 18$  years, have an ASA score of level 3 or 4 and having undergone a surgical intervention, including coeloscopy, excluding cardiac surgery and neurosurgery, requiring analgic treatment through IV in order to relieve postoperative pain.

Patients having at least one of the following risk factors: aged  $\geq 65$  years, OCPD including sleep apnea, high blood pressure, kidney, heart or liver failure.

## **I-SYPRES - OBSERVATIONAL STUDY ON THE DIAGNOSTIC ELEMENTS OF HEART FAILURE WITH PRESERVED SYSTOLIC FUNCTION (HF-PSF) IN PRIVATE PRACTICE**

Last update : 01/01/2019

Head :

Vignal Franck, Sanofi Aventis

Responsible organization :

Sanofi-aventis France

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

## **Main objective**

Describe in private cardiology practice the diagnostic elements of Heart Failure with Preserved Systolic Function (HF-PSF) for patients:

- for whom the private cardiologist makes an initial diagnosis of HF-PSF,
- monitored through appointments and for whom the HF-PSF diagnosis was made during hospitalization

## **Inclusion criteria**

Aged over 18; Presenting HF-PSF that has been diagnosed in the past 6 months either by a private cardiologist or during hospitalization; Agreeing to take part in the study

# **EPCV - CONTINUOUS SURVEY ON HOUSEHOLD LIVING CONDITIONS**

Last update : 09/05/2017

Head :

Accardo Jerome

Responsible organization :

INSEE - Institut national de la statistique et des études économiques

## **Type of database**

Study databases

## **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

## **Main objective**

To investigate actions and health practices of individuals in their daily life.

## **Inclusion criteria**

Households in France (random selection of one household member)

# **ETNA - FIELD STUDY OF INNOVATIVE THERAPIES IN ONCOLOGY: BEVACIZUMAB (AVASTIN®), AN ANTI-ANGIOGENIC AGENT**

Last update : 02/07/2014

Head :

Fourrier-Reglat Annie, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen  
Moore Nicholas, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Responsible organization :

Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

## **Type of database**

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

### Main objective

The main objectives of the study were to describe the population of patients with metastatic colorectal cancer and treated in real-life with first-line bevacizumab, describe the conditions of use of this drug, evaluate the safety of treatments and effectiveness in terms of response and overall and progression free survival at 12 and 24 months follow up.

### Inclusion criteria

Patient with metastatic colorectal cancer who initiated bevacizumab in first-line palliative treatment regardless of the associated cancer treatment between 1 January 2006 and 31 December 2007 (whether or not the treatment is continued); Interval between adjuvant chemotherapy for primary cancer and the initiation of bevacizumab  $\geq$  6 months; Absence of chemotherapy for metastases before initiation of bevacizumab; Patient had not previously treated with bevacizumab, including during a clinical trial or Temporary Authorisation of Use; Patient with a prescribing physician or their head of department who agreed to participate in the study; Patient not participating in a clinical trial (Huriet-Sérusclat), unless it has a standard treatment (control arm) in an open-label Phase III study (bevacizumab known); Patient not objecting to the data collection.

## VENUS - THYROID CANCER

Last update : 09/05/2017

Head :

De Vathaire Florent, Equipe 3 Épidémiologie des cancers : radiocarcinogénèse et effets iatrogènes des traitements UMR 1018 Center for research in epidemiology and population health (CESP)

Rubino Carole, Equipe 3 Épidémiologie des cancers : radiocarcinogénèse et effets iatrogènes des traitements UMR 1018 Center for research in epidemiology and population health (CESP)

Responsible organization :

INSERM

### Type of database

Study databases

## Study databases (details)

Cohort study

### Main objective

To study long-term development and pregnancy outcomes for patients treated for thyroid cancer. Secondary objective: - To estimate the connection between <sup>131</sup>I activity received during thyroid cancer treatment and the risk of cancer or secondary leukaemia; - to determine the role of dose rate in radiation-induced cancer risk.

### Inclusion criteria

Individuals that received treatment for thyroid cancer between 1950 and 2000 in one of the centres participating in the study.

## **OPALE - SURVEY OF THE THERAPEUTIC MANAGEMENT OF RHEUMATOID ARTHRITIS IN FRANCE IN 2006**

Last update : 01/23/2015

Head :

Bregman Bruno, Bristol-Myers Squibb

Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :

Bristol-Myers Squibb (BMS)

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Describe the therapeutic practice used by French rheumatologists in patients suffering from rheumatoid arthritis patients, and describe the treatment patterns used, in a general manner and according to the patients' and disease characteristics.

### **Inclusion criteria**

Register Part: M/F  $\geq 18$  years, with a mild to severe rheumatoid arthritis currently treated or not treated with either symptomatic agents or DMARD (disease-modifying anti-rheumatism drug treatment) (biological or non-biological), seen in rheumatology clinic visit.

Detailed Study Part: patients eligible for the register part, currently treated or initiating a first DMARD (disease-modifying anti-rheumatism drug treatment) (biological or non-biological) at the time of inclusion in the study

## **INCA1, INCA2, INCA3 - FRENCH NATIONAL DIETARY SURVEY**

Last update : 09/05/2017

Head :

Volatier Jean-Luc, ANSES Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail 27-31 avenue du Général Leclerc 94701 Maisons-Alfort Cedex

Dubuisson Carine, ANSES Dietary Survey Unit Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail 27-31 avenue du Général Leclerc 94701 Maisons-Alfort Cedex

Responsible organization :

ANSES

### **Type of database**



Study databases

### **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

#### **Main objective**

Quantitatively characterize individual eating habits, nutritional intakes and dietary exposure in the population to all substances present in food, particularly those that are regulated (environmental contaminants, pesticide residues, food additives).

#### **Inclusion criteria**

INCA1: The sample is made up of 1985 adults aged 15 years old and over and 1018 children aged 3 to 14 years old. Representativeness is ensured for mainland France (excluding Corsica).

INCA2: The sample is made up of 2,624 adults aged 18 to 79 years old and 1,455 children aged 3 to 17 years old.

Representativeness is ensured for mainland France (excluding Corsica).

## **ARVD - GENETIC STUDY OF ARRHYTHMOGENIC RIGHT VENTRICULAR DYSPLASIA**

Last update : 01/01/2018

Head :

Charron Philippe, Centre de référence, CHU Pitié-Salpêtrière, Paris

Responsible organization :

Assistance Publique - Hôpitaux de Paris

### **Type of database**

Study databases

### **Study databases (details)**

Cohort study

#### **Main objective**

General objective: The aim of this project is to investigate the genetic and phenotypic aspects of the disease and determine the role of genetic testing in clinical practice: diagnostic and prognostic value. Secondary objective.

#### **Inclusion criteria**

All patients with ARVD regardless of age and family background. All first degree relations over 10 years old are included.

## **PALIERS - CROSS-SECTIONAL STUDY ON PAIN MANAGEMENT PRACTICES IN EMERGENCY DEPARTMENTS IN FRANCE IN 2010**

Last update : 01/01/2020

Head :

Boccard Éric, Bristol-Myers Squibb  
Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :

Bristol-Myers Squibb France (BMS)

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

Describe the treatment for pain in the emergency departments in France, according to the characteristics of the pain (acute, chronic or induced) - Explore the various treatments in specific sub-groups (non-communicating patients and patients aged 65 years and older).

## **Inclusion criteria**

All conscious adult patients admitted to emergency departments

# **PREVENIR 4 - PERSISTENCE OF EVIDENCE-BASED MEDICAL THERAPIES OVER TIME IN PATIENTS MONITORED BY CARDIOLOGISTS AFTER A FIRST RECENT ACS**

Last update : 09/05/2017

Head :

Amelineau Élisabeth, Bristol-Myers Squibb  
Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :

Bristol-Myers Squibb France (BMS)

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

Describe the persistence of secondary prevention therapeutics, between 3 and 18 months after hospitalization for a first acute coronary syndrome (ACS), in patients monitored by private or hospital cardiologists

## **Inclusion criteria**

M/F  $\geq$ 18 years, outpatient seen in cardiology consultation, hospitalized in 2005 for a first ACS

---

# PHOENIX - COHORT ON THE SCALABLE PROCESS OF POST TRAUMATIC STRESS DISORDERS : THE INVOLVEMENT OF STRESS REGULATION SYSTEMS AND THE ROLE OF ALLOSTATIC LOAD

Last update : 06/01/2020

Head :

Chaudieu Isabelle, Inserm U1061

Responsible organization :

Promoteur/financeur : PHRC régional CHU de Montpellier

Coordinateur scientifique : Inserm U1061

## Type of database

Study databases

### Study databases (details)

Cohort study

### Main objective

The main goal of the project is to study the psychological consequences of violent trauma and the predictive value of biological and clinical stress markers in the development of PTSD

The specific objectives are :

- 1) To evaluate the allostatic load level and its development over time
- 2) To evaluate the prevalence of PTSD and other psychiatric co-morbidities (as depression) on subjects who did endure severe events
- 3) To determine if the allostatic load level is in link with a chronical development of a post traumatic stress disorder

### Inclusion criteria

- women and men aged 18 to 75 years
- person who have experienced a traumatic event during the previous week (0 to 7 days before)
- type of events : physical assault, sexual assault, work accident, road accident, natural disaster.

# SU.FOL.OM3 - SU.FOL.OM3 TRIAL: B-VITAMINS AND N-3 POLYUNSATURATED FATTY ACIDS SUPPLEMENTATION AND RISK OF RECURRENCE OF CARDIOVASCULAR EVENTS

Last update : 01/01/2020

Head :

Hercberg Serge, Unité de Recherche en Epidémiologie Nutritionnelle (UREN), U557 Inserm / U1125 Inra / Cnam / Université Paris 13 (Sorbonne Paris-Cité)

Galan Pilar, Unité de Recherche en EPidémiologie Nutritionnelle (UREN), U557 Inserm / U1125 Inra / Cnam / Université Paris 13 (Sorbonne Paris-Cité)

Responsible organization :

Inserm  
INRA  
CNAM  
Université Paris XIII

## **Type of database**

Study databases

## **Study databases (details)**

Cohort study

## **Main objective**

Primary objectives: check the impact of folate (and vitamin B6 or B12) and/or omega 3 supplements in preventing the recurrence of ischemic disorders in patients with a background of ischemic cardio or cerebrovascular history.

Secondary objective: assess the role of certain genetic mutations in the ability of supplements to reduce the risk of cardiovascular diseases.

## **Inclusion criteria**

Subjects aged between 45 and 80, having presented a myocardial infarction, unstable angina or stroke in the period preceding their inclusion (event occurred at least one month and at the most one year prior to inclusion).

Exclusion criterion: subjects that have to take B12, folic acid or B6 supplements, subjects under methotrexate treatment, subjects suffering from a life-threatening non-cardiovascular disease over the 5 years of the study, severe chronic kidney disease sufferers.

# **COLCHIC - ASSESSMENT OF CHEMOTHERAPY PRACTICES IN PATIENTS WITH COLORECTAL CANCER AT THE BORDEAUX UNIVERSITY HOSPITAL**

Last update : 01/01/2018

Head :

Fourrier-Reglat Annie, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen  
Moore Nicholas, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Responsible organization :

Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

## **Type of database**

Study databases

## **Study databases (details)**

Longitudinal study (except cohorts)

## **Main objective**

The main objective was to evaluate the conditions of use and monitoring of chemotherapy for the treatment of colorectal cancer clinical departments at the University Hospital of Bordeaux, and to assess the appropriateness of prescribing practices in relation to current official recommendations.

### **Inclusion criteria**

Patient with colon or rectal cancer treated with chemotherapy between 1 March 2005 and 1 March 2006;  
Patient followed at the Bordeaux University Hospital; Patient not objecting to the data collection.

## **INTERMEDE - CROSS-SECTIONAL SURVEY ON PATIENT- PHYSICIAN INTERACTION AND SOCIAL HEALTH INEQUALITIES**

Last update : 10/01/2017

Head :

Lang Thierry, Unité Inserm 558, Faculté de Médecine de Toulouse

Responsible organization :

INSERM

Université Paul Sabatier

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To clarify elements of physician-patient interaction that may explain, beyond social differences in health care sought by patients, differentiated treatment that may ultimately contribute to differences in health. - Consultation observation and semi-structured post-consultation interviews with patients and doctors independently. - Mirrored-questionnaires where patients and physicians answered the same questions post-consultation.

### **Inclusion criteria**

Patient inclusion: - at least 18 years old - not attending a general practitioner for emergency purposes - who are not attending their first consultation - not pregnant or have given birth within the last six months - who have signed the informed consent agreement

## **THEATRE - OBSERVATIONAL STUDY OF THE THERAPEUTIC STRATEGY FOR MANAGEMENT OF ACUTE EXACERBATION OF CHRONIC BRONCHITIS IN REAL-LIFE PRACTICE**

Last update : 09/05/2017

Head :

Molimard Mathieu, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Moore Nicholas, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Responsible organization :

**Type of database**

Study databases

**Study databases (details)**

Longitudinal study (except cohorts)

**Main objective**

The study objectives were to describe how the management of AECB in real-life conditions of prescription and to assess the effectiveness of the therapeutic strategies used.

**Inclusion criteria**

The study objectives were to describe how the management of AECB in real-life conditions of prescription and to assess the effectiveness of the therapeutic strategies used.

**- INFLAMMATORY PULMONARY DISEASES IN COAL MINERS**

Last update : 09/05/2017

Head :

Nadif Rachel, Recherche en Épidémiologie et Biostatistique

Responsible organization :

Institut National de Santé et Recherche Médicale

**Type of database**

Study databases

**Study databases (details)**

Cohort study

**Main objective**

The study of Lorraine Miners was set up in 1990 with 2 main objectives.

- To estimate the predictive value of chest tomodynamometry and of sensitive functional tests in the development of pneumoconioses.

- To test specific hypotheses about the physiology of emphysema and fibrosis using biological markers: proteases/antiproteases, oxidative/antioxidative and degradation/repair balances.

The study of genetics has subsequently been developed.

**Inclusion criteria**

-Male- between 34 and 50 years old - working in a coal mine

**COI - ORGANIZATIONAL CHANGES AND COMPUTERIZATION**

Last update : 09/05/2017

Head :

Amira Selma

Hamon-Cholet Sylvie, UR Dynamiques des Organisations et du Travail

Responsible organization :

DARES - Direction de l'Animation de la Recherche, des Etudes et des Statistiques

CEE - Centre d'études de l'emploi

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

C.O.I. is a matched survey employers/employees on organizational changes and computerization, enriched with administrative data from both employers and employees.

The objective is to measure the organizational changes of companies in terms of management and use of information and communication technologies and their effects on employees' work over a three-year period (2003-2006).

## **Inclusion criteria**

Staff from private companies, some central civil service administrations and establishments in the hospital sector with 10 employees and over, and employees from units with 20 employees and over in the private and hospital sector, with 10 employees and over in the civil service.

# **PARTAGE - CASE-CONTROL STUDY ON PARKINSON'S DISEASE AMONG MUTUALITÉ SOCIALE AGRICOLE AFFILIATES IN FIVE DÉPARTEMENTS**

Last update : 09/05/2017

Head :

Elbaz Alexis

Responsible organization :

Institut National de la Santé et de la Recherche Médicale

Institut National de Veille Sanitaire

Mutualité Sociale Agricole

## **Type of database**

Study databases

## **Study databases (details)**

Case control study

## **Main objective**

To determine the risks of contracting Parkinson's Disease for those affiliated with Mutualité Sociale

Agricole.

### **Inclusion criteria**

- Men or women - adult - affiliates of Mutualité Sociale Agricole

## **PLEIAD - DESCRIPTIVE EPIDEMIOLOGICAL STUDY IN NURSING HOME**

Last update : 01/01/2020

Head :  
Crochard Anne

Responsible organization :  
Lundbeck SAS

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To describe the population of nursing home residents in France and in particular to quantify transitions into or out of the nursing home and to describe the characteristics of the nursing home residents at admission and discharge.

### **Inclusion criteria**

To be eligible, the residents had to have been admitted to the nursing home for the first time (from home, hospital or another nursing home) or following a hospitalization, or had to have been temporarily or permanently discharged from the nursing home (to home, hospital or another nursing home) in the previous three months.

## **PHOTOGRAF - FACTEURS DE RISQUE DU GLAUCOME PRIMITIF À ANGLE OUVERT CHEZ LES SUJETS ATTEINTS D'HYPERTONIE OCULAIRE**

Last update : 01/01/2018

Head :  
Troy Sylvie, Pfizer

Responsible organization :  
Pfizer

### **Type of database**

Study databases



## Study databases (details)

Case control study

### Main objective

Main:

- Evaluate the differences between the cases afflicted with primary open-angle glaucoma (POAG) and the controls afflicted with isolated ocular hypertension (OHT) concerning the potential risk factors, including sociodemographic factors, systemic diseases and treatments in progress, family antecedents of glaucoma, the duration of the disease, the factors linked to the eye, nutritional factors and the exposure to environmental neurotoxic compounds.

Secondary:

- Derive a predictive function by using the significant risk factors, in such a way as to assist ophthalmologists in classifying patients having a hypertonia with a high risk of developing glaucoma

### Inclusion criteria

For the cases:

- Male or female aged 40 years or older;
- Patients with a diagnosed POAG, confirmed by a structural and function degradation;
- Patients for whom a IOP > 21 mm Hg was recorded at one time in the evolution of the disease.

For the controls

- Male or female paired with the age of the case ( $\pm$  2 years);
- Patients having an IOP > 21 mm Hg on the day of the examination and/or a hypotonisant treatment;
- Patients with a visual acuity with optimal correction higher than 5/10;
- Patients with no anomaly in the visual field characteristic of glaucoma.

## SISTOLA - OBSERVATIONAL USAGE AND EFFECTIVENESS OF CANDESARTAN IN HEART FAILURE TREATMENT IN FRANCE

Last update : 01/01/2020

Head :

Thomas-Delcourt Florence, AstraZeneca

Responsible organization :

ASTRAZENECA / TAKEDA

### Type of database

Study databases

### Study databases (details)

Longitudinal study (except cohorts)

### Main objective

To describe the way candesartan is used in heart failure treatment in France (treatment initiation and follow-up) treatment discontinuation occurrences and reasons why and patients clinical evolution

### Inclusion criteria

Patients starting or who have recently started (less than 30 days) a treatment with candesartan for heart failure

## **DISCOVER - SCREENING FOR EXTRACORONARY ATHEROTHROMBOTIC LOCALIZATIONS AFTER AN ACUTE CORONARY SYNDROME AND CLINICAL MANAGEMENT OF PATIENTS**

Last update : 09/05/2017

Head :

Herrmann Marie-Annick, Bristol-Myers Squibb

Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :

Bristol-Myers Squibb France (BMS)

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Evaluate the frequency of uninvestigated and investigated patients in the detection of extra-coronary atherothrombotic locations in a population of patients having an history of ACS in the year preceding the selection visit and followed by office based cardiologists.

### **Inclusion criteria**

M/F,  $\geq 18$  years, examined in consultation by an office based cardiologist, with an history of ACS with or without ST-segment elevation in the 12 months preceding the selection

## **ADRS (ADOLESCENT DEPRESSION RATING SCALE) - ÉTUDE TRANSVERSALE SUR DES PATIENTS ADOLESCENTS : DESCRIPTION DE L'EXPÉRIENCE DÉPRESSIVE DE L'ADOLESCENT**

Last update : 09/05/2017

Head :

Falissard Bruno, INSERM U669

Revah-Levy Anne , INSERM U669

Gasquet Isabelle

Responsible organization :

INSERM - Institut National de Santé et Recherche Médicale

### **Type of database**

Bases de données issues d'enquêtes

### **Study databases (details)**

Etudes transversales non répétées (hors enquêtes cas-témoins)

#### **Main objective**

Décrire l'expérience dépressive de l'adolescent et créer un outil de mesure de l'intensité dépressive destiné spécifiquement aux adolescents de 13 à 20 ans.

#### **Inclusion criteria**

Patients de 13 à 20 ans hospitalisés dans les 15 centres médicaux concernés (hors retards mentaux et psychoses)

## **YZATIS - EVALUATION ET ÉVOLUTION DE LA PRISE EN CHARGE THÉRAPEUTIQUE À MOYEN TERME DES PATIENTS SÉROPOSITIFS POUR LE VIH-1 TRAITÉS AVEC UNE COMBINAISON ANTIRÉTROVIRALE INCLUANT ATAZANAVIR**

Last update : 09/05/2017

Head :

Bennai Yacia, Bristol-Myers Squibb

Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :

Bristol-Myers Squibb France (BMS)

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

#### **Main objective**

Describe the arrangements for caring for patients treated by a combination of treatments including ATAZANAVIR according to the grounds for introducing the treatment (virological failure of the previous treatment or other)

#### **Inclusion criteria**

M or F, aged 18 years or older, seropositive for HIV-1, treated via a combination of ARV treatments including ATV for which the main reason for introduction of the current treatment was: - virological failure of the previous treatment OR  
- another reason.

## **APTOR - PROSPECTIVE OBSERVATIONAL STUDY OF PATIENTS WITH ACUTE CORONARY SYNDROME MANAGED BY PCI AND ANTIPLATELET TREATMENT: ANTIPLATELET TREATMENT OBSERVATIONAL REGISTRY**

Last update : 01/01/2018

Head :

Médecin pharmacoépidémiologiste

Responsible organization :

Eli Lilly

## **Type of database**

Study databases

## **Study databases (details)**

Longitudinal study (except cohorts)

## **Main objective**

Primary objective: evaluate the use of healthcare resources and cost in the 12 months following a percutaneous coronary intervention (PCI) combined with antiplatelet treatment in patients with acute coronary syndrome.

Secondary objectives: quality of life and clinical evolution.

## **Inclusion criteria**

Be diagnosed with acute coronary syndrome,

- treatment with percutaneous coronary intervention in routine care;
- start or continue antiplatelet treatment.

# **EPIDOS - RISK FACTORS FOR OSTEOPOROTIC FRACTURE**

Last update : 09/05/2017

Head :

Dargent Patricia, U 953

Bréart , U953

Responsible organization :

Institut National de la Santé et de la Recherche Médicale

## **Type of database**

Study databases

## **Study databases (details)**

Cohort study

## **Main objective**

General objective: to study the risk factors for osteoporotic fracture with a focus on femoral neck fracture

## **Inclusion criteria**

Women over 75 years old included in electoral registers with no prior femoral fracture

# **EVALHOSPITAM OR EVHAN - EVALUATION OF HOSPITALISATION FOR PATIENTS WITH ANOREXIA NERVOSA: EFFICACY OF TREATMENT AND RESEARCH PREDICTIVE OUTCOME FACTORS**

Last update : 03/01/2020

Head :

Godart Nathalie, U669 INSERM TROUBLES DES CONDUITES DE L'ADOLESCENT

Responsible organization :

INSTITUT MUTUALISTE MONSOURIS

## **Type of database**

Study databases

## **Study databases (details)**

Longitudinal study (except cohorts)

## **Main objective**

General objective: to evaluate the impact (efficacy) of five different inpatient treatment modalities for anorexia nervosa. Individual global health outcome (physical, psychiatric, nutritional and social) will be measured on a short-term basis (on discharge), as well as hospital treatment meeting or exceeding statutory charges on a medium term basis (12 months), taking into account confounding factors and recognised prognostic factors. Global health is evaluated using the Morgan and Russell scale as well as secondary criteria for deteriorating health (see evaluation criteria methodology). Secondary objectives: - to determine predictive factors for failures in inpatient treatment, duration of hospitalisation, and treatment costs using clinical epidemiology methodology. - to begin a long-term prospective study to evaluate the psychological, physical, nutritional and social outcome of a homogeneous cohort of subjects suffering from severe anorexia nervosa. - two "ancillary" biological sub-projects: - to evaluate the development of depressive and anxiety symptoms in relation to changes in clinical nutritional status and serotonin markers - to evaluate cellular immunity development in relation to changes in clinical nutrition.

## **Inclusion criteria**

- Patients between 8 and 65 years old; - patients hospitalised for anorexia nervosa (who only receive treatment by necessity of somatic condition: rapid weight loss, or significantly underweight with a BMI of less than 15, or their psychological condition), as well as parents of minor and adult patients still living at home; - obtained written consent; - patients not covered by social security health insurance.

# **EVADEF - MEDICO-ECONOMIC EVALUATION OF AUTOMATIC IMPLANTABLE DEFIBRILLATOR**

Last update : 09/05/2017

Head :

Marijon Eloi

Le Heuzey Jean-Yves, DÉPARTEMENT DE CARDIOLOGIE A, HÔPITAL EUROPÉEN GEORGES POMPIDOU, 20 RUE LEBLANC, 75908 PARIS CEDEX 15

Responsible organization :  
Société Française de Cardiologie

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

To assess the cost of automatic implantable defibrillator usage in France. Secondary objective: - Complications associated with automatic implantable defibrillators; - To investigate cause-specific mortality in patients with automatic implantable defibrillators; - The psychological impact of automatic implantable defibrillators; - Incidence of appropriate and inappropriate therapies.

### **Inclusion criteria**

All patients who received an automatic implantable defibrillator

## **EROPP - SURVEY ON REPRESENTATIONS, OPINIONS, AND PERCEPTIONS REGARDING PSYCHOACTIVE DRUGS**

Last update : 01/01/2018

Head :

Spilka Stanislas

Le Nézet Olivier

Responsible organization :

OFDT

### **Type of database**

Study databases

### **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

### **Main objective**

Current perceptions and opinions of 15-75 year-olds French people of licit and illicit psychoactive substances and the main public initiatives.

### **Inclusion criteria**

General population, individuals aged 15 to 75.

## **EPIPAGE - SURVIVAL OF VERY PRETERM INFANTS**

Last update : 01/01/2018

Head :

Larroque Béatrice, U953 - Unité de recherche épidémiologique en santé périnatale et santé des femmes et des enfants

Ancel Pierre-Yves, U953 - Unité de recherche épidémiologique en santé périnatale et santé des femmes et des enfants

Responsible organization :

INSTITUT NATIONAL DE LA SANTE ET DE LA RECHERCHE MEDICALE - INSERM

## Type of database

Study databases

### Study databases (details)

Cohort study

## Main objective

Study the evolution of severely preterm children born between 22 and 32 weeks of gestational age in 9 regions of France in 1997, and 2 comparison groups at 33-34 and 39-40 weeks.

## Inclusion criteria

Severely preterm children (birth or interruption 22-32 weeks of amenorrhea (SA) or weight between 0,5 and 1,499 Kg.

Control group: Group 1: 39-40 SA ; Groupe 2 : 33-34 SA in centers participating to the study.

# VESUVE (BETA) - VELCADE: STUDY OF REAL-LIFE USE

Last update : 09/05/2017

Head :

Fourrier-Reglat Annie, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Moore Nicholas, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Responsible organization :

Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

## Type of database

Study databases

### Study databases (details)

Longitudinal study (except cohorts)

## Main objective

The objectives are to evaluate the response and survival of patients treated by Velcade®, to describe the population of patients initiating treatment with Velcade® (socio-demographic data, previous treatments, indication), and describe the conditions of use Velcade® (dose, number and frequency of treatment cycles)

## Inclusion criteria

Patients initiating Bortezomib treatment between 1 May 2004 and April 30 2006 (whether or not treatment is continued); Patient unexposed to bortezomib, including during a clinical trial or temporary use authorization; Patient followed by a hospital physician having agreed to participate in the study; Patient not participating in a clinical trial; Patient not objecting to data collection

## **KABP - KNOWLEDGE, ATTITUDES, PRACTICES AND BELIEFS RELATED TO HIV IN ÎLE DE FRANCE**

Last update : 01/01/2020

Head :

Warszawski Josiane, Équipe: Épidémiologie de VIH et des IST  
Beltzer Nathalie

Responsible organization :

Institut National de la Santé et de la Recherche Médicale  
Observatoire régional de santé Île-de-France  
Agence Nationale de Recherche sur le Sida  
Institut de veille sanitaire  
Institut national de prévention et d'éducation pour la santé

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To follow the evolution of knowledge concerning HIV and AIDS and antiretroviral treatment, attitudes and opinions with regard to infected persons, the perception of being contaminated by the virus, screening tests and condom use.

### **Inclusion criteria**

- Men or women - aged 18 to 69 years old

## **PRISM - MULTIDISCIPLINARY CARE OF THE METABOLIC SYNDROME DURING SPA THERAPY**

Last update : 09/05/2017

Head :

Moore Nicholas, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Responsible organization :

Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

### **Type of database**

Study databases



## Study databases (details)

Longitudinal study (except cohorts)

### Main objective

The main objective of this study was to determine the maximal effect of multidisciplinary management initiated during spa therapy on metabolic syndrome at one year.

### Inclusion criteria

Patient presenting a metabolic syndrome according to the criteria of the NCEP-ATP III (at least three of the following criteria: waist circumference > 102 cm for men and 88 cm for women, triglycerides ? 1.5 g/l, HDL-Cholestérol < 0.4 g/l for men and 0.5 g/l for women, fasting glycaemia ? 1.10 g/l, blood pressure ? 130/85 mmHg 130/85); greater than 18 years old; patient having a BMI < 40; patient having a fasting glycaemia < 1.26 g/l; patient who can be followed during the study period; patient able to understand the protocol and complete the questionnaire; patient initiating spa therapy for "metabolic disease" (alone or with a second indication); patient accepting to participate in the study.

## CATHY - THYROID CANCER STUDY IN CALVADOS, MARNE AND THE ARDENNES

Last update : 09/05/2017

Head :

Guénel Pascal, CESP - Inserm UMR S 1018 Epidémiologie environnementale des cancers de l'adulte

Responsible organization :

Institut National de la Santé et de la Recherche Médicale

### Type of database

Study databases

## Study databases (details)

Case control study

### Main objective

To identify hormonal, reproductive environmental, and lifestyle risk factors of thyroid cancers.

### Inclusion criteria

- Men or women - adult

## FLVS II - FLEURBAIX LAVENTIE VILLE SANTÉ STUDY II

Last update : 09/05/2017

Head :

Charles Marie-Aline, INSERM U1018 CENTRE DE RECHERCHE EN ÉPIDÉMIOLOGIE ET SANTÉ DES POPULATIONS

Responsible organization :

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

To investigate the determining factors for changes in adiposity in adolescents and their parents in the general population

## Inclusion criteria

School-age children in primary schools that agreed to participate in the first survey in 1992 (FLVS I). Families living in Fleurbaix and Laventie.

## CASC - BEHAVIORS AND ATTITUDES OF CYCLISTS CONCERNING ROAD SAFETY AND USE OF HELMET

Last update : 08/01/2020

Head :

LAGARDE Emmanuel, U897 PRÉVENTION ET PRISE EN CHARGE DES TRAUMATISMES ( Equipe de l'ISPED - Institut de Santé Publique, d'Epidémiologie et de Développement)

Responsible organization :

INSERM - Institut National de la Santé et de la Recherche Médicale

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Understand the relation between the use of the helmet and risk behaviors in urban cyclists. Enlighten the decision concerning road safety politics.

## Inclusion criteria

People borrowing a bicycle in a municipal structure for a minimum period of 4 months

## - BAROMÈTRE SANTÉ MÉDECINS GÉNÉRALISTES

Last update : 01/01/2019

Head :

Gautier Arnaud

Responsible organization :  
Institut National de Prévention et d'Education pour la Santé (INPES)

### **Type of database**

Study databases

### **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

### **Main objective**

Know the opinions, behaviors and practices of general practitioners in terms of prevention; various themes are addressed here, in particular opinions with regards to vaccination, cancer screening and viral hepatitis, treatment for addiction problems, etc.

It also aims to identify preventive practices (primary, secondary), detect the difficulties encountered? especially in order to guide, strengthen preventive actions with health professionals.

### **Inclusion criteria**

General practitioner exercising in metropolitan France through liberal activity and included in the CEGEDIM file.

## **PREVAS - ETUDE DE PREVALENCE DE L'ASTHME PERSISTANT SÉVÈRE MAL CONTRÔLÉ**

Last update : 09/05/2017

Head :  
Anne Ponthieux, Direction Relations Economiques et Institutionnelles

Responsible organization :  
Novartis Pharma S.A.S.

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Assessment of prevalence of uncontrolled severe persistent asthma in a representative population of asthmatic patients followed by pneumologist or allergy specialist

### **Inclusion criteria**

severe persistent asthmatic patients viewed in routine consultation, with controlled or uncontrolled asthma in the previous year.

## **DN4 MIXTE - EVALUATION STUDY OF THE DN4 QUESTIONNAIRE**

## IN MIXED PAIN

Last update : 01/01/2019

Head :  
Millet Véronique, Pfizer

Responsible organization :  
Pfizer

### Type of database

Study databases

### Study databases (details)

Longitudinal study (except cohorts)

### Main objective

Main:

Compare the average scores of the DN4 questionnaire observed in 4 sub-groups of patients monitored for lumbar radiculalgia.

Secondary objectives:

- For each item on the DN4, compare the proportion of the positive items between each sub-group.
- Compare the proportions of the subjects that have a DN4 score > 4/10 between each sub-group.
- Compare the proportions of the subjects that have been considered as having neuropathic component by a 1st investigator between each sub-group.
- Estimate the sensitivity, specificity, the positive and negative predictive value of the DN4 scale in relation to the opinion of the 1st investigator within each sub-group:
  - o By taking a DN4 threshold at 4/10
  - o Using the ROC curve
- Compare the averages of the scores of the modified Schöber index and the ""hand-to-floor distance"" according to the presence or not of a neuropathic component in the 4 sub-groups.

### Inclusion criteria

Patients ? 18 years,

- First consultation in an investigating center,
- Patients having a lumbar pain with or without radiculalgia (patients presenting another pain of moderate to severe were excluded),
- Pain present for at least 3 months,
- Moderate to severe pain (intensity ? 4 on the 11-point Lickert scale),
- Patients who have signed the written consent.

## - DECENNIAL HEALTH SURVEY (SURVEY ON HEALTH AND MEDICAL CARE)

Last update : 09/05/2017

Head :  
de Saint Pol Thibaut, Bureau "Etat de santé de la population"

Responsible organization :

INSEE - Institut national de la statistique et des études économiques

DREES - Direction de la recherche, des études, de l'évaluation et des statistiques

## **Type of database**

Study databases

## **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

## **Main objective**

The first health survey was conducted in 1960-1961. The following surveys took place in 1970-1971, 1980-1981 and 1991-1992, and the last in 2002-2003. The objectives of the decennial health survey are: - to describe population health: reported morbidity and perceived health; - to provide quantitative data on health care usage and prevention (by volume and value); - to analyse health, health care usage and prevention in relation to sociodemographic characteristics (age, gender, profession ...) of individuals and their households.

## **Inclusion criteria**

French people from 11 years of age (for 2002-2003)

# **- TRANSVERSAL STUDY ON FRENCH NATIONAL PREVALENCE OF DENTAL DECAY IN CHILDREN (2006)**

Last update : 09/05/2017

Head :

Desfontaine Jacques

Hescot Patrick

Responsible organization :

Union Française pour la Santé Bucco-Dentaire (UFSBD)

## **Type of database**

Study databases

## **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

## **Main objective**

Perform reliable and representative studies in France of the dental health of school children from 6 to 12 years old and at regular intervals (every 5 years). Through these studies, it must be possible to assess the health, pathology, care administered and treatment needs of children as well as the trends in the population's dental health.

## **Inclusion criteria**

children born between 1/07/1999 and 31/12/1999 for 6 year-old children

children born between 1/01/1994 and 30/06/1994 for 12 year-old children

## **CECILE - BREAST CANCER: EPIDEMIOLOGICAL STUDY ON ENVIRONMENT IN CÔTE D'OR AND ILLE-ET-VILAINE**

Last update : 01/01/2018

Head :

Guénel Pascal, CESP - UMR S 1018 Epidémiologie environnementale des cancers de l'adulte

Responsible organization :

Institut National de la Santé et de la Recherche Médicale

### **Type of database**

Study databases

### **Study databases (details)**

Case control study

### **Main objective**

To highlight environmental risk factors for breast cancer and investigate the role of lifestyle or exposure to harmful conditions. To identify genetic susceptibility factors and study gene?environment interaction.

### **Inclusion criteria**

- women aged 75 years old and under

## **ESTHER - ESTROGEN THROMBOEMBOLISM RISK**

Last update : 09/05/2017

Head :

Scarabin Pierre-Yves

Responsible organization :

Institut National de la Santé et de la Recherche Médicale

### **Type of database**

Study databases

### **Study databases (details)**

Case control study

### **Main objective**

To evaluate the risk of venous thromboembolism in connection with hormone therapy for menopause while considering the administration route for oestrogen and types of progestin.

### **Inclusion criteria**

- Women - between 45 and 65 years of age - menopausal - with no cardiovascular history or breast cancer

## **ORBIT - COMORBIDITIES ASSOCIATED WITH PERIPHERAL NEUROPATHIC PAIN (PNP)**

Last update : 01/01/2018

Head :

Millet Véronique, Pfizer

Responsible organization :

Pfizer

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Primary:

- Evaluate the frequency of anxiety and mood disturbances and the degree of sleeping disorders in patients monitored for PNP

Secondary:

- Describe the links between the components of the PNP and the existence of anxiety, mood disturbances and the degree of sleeping disorders,

- Describe the relations between the components of PNP, anxiety, sleeping disorders, mood disturbances on the one hand and the quality of life on the other hand,

- Describe the relation between the components of the PNP, the existence of anxiety and mood disturbances, the degree of sleep disorders on the one hand and the current drug care of patients on the other hand.

### **Inclusion criteria**

Patients > 18 years,

- Peripheral neuropathic pain (in relation with a lesion of the peripheral nervous system).

- Intensity of the pain > 3 on the NVS,

- Radiculalgia without lumbar pain or with lumbar pain of which the intensity of the pain is < 3 on the NVS (intensity of the lumbar pain from 0 to 2),

- Through general neurology: patient consulting for his PNP or for the etiology of this PNP,

- Through home consultation: patient consulting for a PNP for the 1st or 2nd consultation.

## **CADISP - ÉTUDE CAS-TÉMOINS SUR LES FACTEURS DE RISQUE DES ACCIDENTS ISCHÉMIQUES CÉRÉBRAUX ET LA DISSECTION D'UNE ARTÈRE CERVICALE CHEZ LES JEUNES ADULTES**

Last update : 01/01/2019

Head :

Debette Stéphanie, Inserm U740, Université Paris 7

Responsible organization :  
INSERM - Institut National de Santé et Recherche Médicale  
AP-HP

## **Type of database**

Bases de données issues d'enquêtes

## **Study databases (details)**

Etudes cas-témoins

## **Main objective**

Effectuer des recherches sur la dissection d'une artère cervicale et les accidents ischémiques cérébraux chez les jeunes adultes, et en particulier sur les facteurs de risque, le traitement et les prédicateurs de résultats

## **Inclusion criteria**

Patients hospitalisés dans un des centre neurologiques participants

- Patient victime d'une dissection d'une artère cervicale
- Patients victimes d'un accident ischémique cérébral sans dissection d'une artère cervicale
- Patients de la population générale sans antécédents de maladie vasculaire

# **PREVENIR 5 - PERSISTENCE OF COMBINATION OF EVIDENCE-BASED MEDICAL THERAPIES IN PATIENTS WITH RECENT HISTORY OF ACS IN PRIMARY CARE SETTING**

Last update : 09/05/2017

Head :

Amelineau Élisabeth, Bristol-Myers Squibb  
Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :

Bristol-Myers Squibb France (BMS)

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

Describe the persistence of secondary prevention therapeutics, between 3 and 30 months after hospitalization for a first acute coronary syndrome (ACS), in patients monitored by general practitioners

## **Inclusion criteria**

Patient  $\geq 18$  years, outpatient seen in a general practice consultation, hospitalized in 2004 or 2005 for a 1st ACS



# SU.VI.MAX 2 - STUDY OF THE RELATIONS BETWEEN GLOBAL FOOD BEHAVIORS AND/OR SPECIFIC NUTRITIONAL FACTORS AND AGEING ON A LARGE POPULATION OF MATURE ADULTS

Last update : 01/01/2019

Head :

Hercberg Serge, Unité de Recherche en Epidémiologie Nutritionnelle (UREN), U557 Inserm / U1125 Inra / Cnam / Université Paris 13 (Sorbonne Paris-Cité)

Galan Pilar, Unité de Recherche en Epidémiologie Nutritionnelle (UREN), U557 Inserm / U1125 Inra / Cnam / Université Paris 13 (Sorbonne Paris-Cité)

Kesse Emmanuelle, Unité de Recherche en Epidémiologie Nutritionnelle (UREN), U557 Inserm / U1125 Inra / Cnam / Université Paris 13 (Sorbonne Paris-Cité)

Responsible organization :

INSTITUT NATIONAL DE LA SANTE ET DE LA RECHERCHE MEDICALE - INSERM

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

General objective: study the relations between global eating behaviors and/or some specific nutritional factors (provisions and status of anti-oxydants, folates, Omega 3 PUFA, energetic and protein provision, etc.) evaluated from data collected between 1994 and 1996 on a large population of mature adults, and:

- The global quality of ageing among this population, evaluated 12 and 15 years later (2006 and 2009) through a global indicator, realized from different components determining successful aging (cognitive state, mood, nutritional status, bone status, balance disorders, sensory disorders, perceived health, integration, social, physical activity, pathologies existence).

- Each specific different specific functional component related to the quality of aging: cognitive functions and depression, risk of sarcopenias or osteoporosis, insulin resistance, life quality, chronic diseases (cancers, cataract, macular degeneration). Secondary objective: evaluate the role of oxidative stress, inflammatory status, insulin resistance, some genetic factors implied in the relation between nutritional behaviors, and/or some more specific nutritional factors, and aging quality

## Inclusion criteria

Voluntary subjects, having participated to the SUI.VI.MAX cohort, and aged between 55 and 70.

# - ASBESTOS EXPOSURE AND DIGESTIVE CANCERS

Last update : 09/05/2017

Head :

Launoy Guy, Cancers et Populations

Responsible organization :

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

To determine whether occupational asbestos exposure increases the risk of digestive cancers. To define the relationship between exposure to asbestos and cancer of the small intestine and oesophagus in humans.

## Inclusion criteria

- Men or women - exposed to asbestos in their place of work

## PREMODA - MODE OF DELIVERY AND TERM BREECH PRESENTATION IN THE PREMODA COHORT.

Last update : 09/05/2017

Head :

Goffinet François, CIC901 CIC - GH NECKER-COCHIN-PARIS CENTRE DESCARTES Équipe/activité : Recherche clinique U953 Recherche épidémiologique en santé périnatale et santé des femmes et des enfants Équipe/activité : Épidémiologie clinique et organisation des soins

Responsible organization :

Assistance Publique - Hôpitaux de Paris Hôpital Cochin - AP-HP  
Institut national de la santé et de la recherche médicale - INSERM

## Type of database

Study databases

## Study databases (details)

Case control study

## Main objective

The purpose of this work was to report labor and birth management practices for term breech presentation in France and Belgium in 2001-2002 and to describe indications for cesarean sections (before labor, emergency situations) in breech presentations.

## Inclusion criteria

woman - adult - term of pregnancy greater than or equal to 37 weeks of amenorrhea - single pregnancy

## REN-ART NEPHROTEST - IDENTIFICATION OF ARTERY STIFFENING AS A RISK FACTOR ASSOCIATED WITH

# DETERIORATION IN RENAL FUNCTION FOR INDIVIDUALS WITH CHRONIC KIDNEY DISEASE

Last update : 04/01/2019

Head :

BOUTOUYRIE Pierre, U970 PARCC (PARIS - CENTRE DE RECHERCHE CARDIOVASCULAIRE)

Responsible organization :

ASSISTANCE PUBLIQUE - HÔPITAUX DE PARIS

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

General objective: to show that aortic stiffness predicts the rate of renal function decline in patients with moderate to severe CKD. Secondary objectives: - To identify the relationship between arterial parameters, renal function and metabolic parameters; - to examine the evolution of arterial parameters throughout kidney disease progression; - to examine the evolution of arterial parameters other than aortic stiffness and renal function deterioration; - to correlate vascular properties with the occurrence of cardiovascular and renal events with a composite endpoint (end-stage KD...).

## Inclusion criteria

Glomerular filtration rate less than 60 ml/min

# SALT-I - STUDY OF ACUTE LIVER TRANSPLANT

Last update : 10/27/2017

Head :

Gülmez Sinem Ezgi, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Moore Nicholas, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Responsible organization :

INSERM

## Type of database

Study databases

## Study databases (details)

Case control study

## Main objective

Acute liver failure, Nimesulide, non-steroidal anti-inflammatory drugs (NSAIDs), liver transplantation, drug intoxication, retrospective, case-population, Europe, Pharmacoepidemiology, Department of

### **Inclusion criteria**

Patient who was at least 18 years old at the time of registration on transplant list; Patient with acute liver failure placed on liver transplantation list and exposed to NSAIDs during the 30 days before the 1st signs or symptoms of liver disease, whether or not transplantation was performed; Patient resident of the participating country; Excluding elective liver transplants for chronic diseases such as cirrhosis, chronic hepatitis or cancer.

## **POLA - LONGITUDINAL STUDY ON AGE-RELATED EYE DISEASES**

Last update : 09/05/2017

Head :

Delcourt Cécile, Centre de Recherche Inserm U897 "Epidémiologie et Biostatistiques"

Responsible organization :

INSERM

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

To investigate risk factors of age-related eye diseases (Age-Related Macular Degeneration (AMD), cataracts)

### **Inclusion criteria**

Elderly people over 60 years of age living in Sète (Hérault)

## **- ÉTUDE TRANSVERSALE SUR LA SATISFACTION DES PATIENTS FACE AUX SERVICES DE CONSULTATION AMBULATOIRE DES HÔPITAUX**

Last update : 09/05/2017

Head :

Falissard Bruno, Inserm U1018 - Centre de Recherche en Épidémiologie et Santé des Populations Équipe: Genre, Santé sexuelle et Reproductive  
Gasquet Isabelle, UPRES JE 2360

Responsible organization :

INSERM - Institut National de la Santé et de la Recherche Médicale

### **Type of database**

Bases de données issues d'enquêtes

## Study databases (details)

Etudes transversales non répétées (hors enquêtes cas-témoins)

### Main objective

Sélectionner des éléments, identifier les dimensions, la fiabilité, la mesure interne et la validité concurrente des items du questionnaire.

### Inclusion criteria

Patients venant en consultation ambulatoire dans 3 hôpitaux d'enseignement publics de la région parisienne (Paul Brousse, Bichat et George Pompidou hôpitaux européens), dans les services suivants:

- 7 départements médicaux (médecine interne, rhumatologie, 2 cardiologie, la dermatologie, les maladies infectieuses et de l'oncologie)

- 3 départements de chirurgie ambulatoire (urologie, orthopédie, gynécologie chirurgicale).

Les patients hospitalisés avant ou immédiatement après la consultation ont été exclus.

## PROXAIR - ETUDE DE PROXIMOLOGIE DANS L'ASTHME PERSISTANT SÉVÈRE

Last update : 09/05/2017

Head :

Ponthieux Anne, Direction Relations Économiques et Institutionnelles

Responsible organization :

Novartis Pharma S.A.S.

### Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

### Main objective

Assess physical, psychic and socio-economic impact on patients and spouse of severe persistent asthma treated with high-dose inhaler steroids and long-acting  $\beta_2$  agonists (LABA), according to the asthma control level

### Inclusion criteria

Patient inclusion Criteria :

- Ambulatory patients, able to cooperate, of either sex, at least 18 years of age.

- Patients with severe persistent asthma receiving for at least three months a continuous and stable treatment of high-dose inhaler steroids (? 1 000  $\mu\text{g}/\text{d}$  of beclometasone dipropionate excluding micronized forms in metered-dose inhalers, ? 800  $\mu\text{g}/\text{d}$  of beclometasone dipropionate in micronized form in metered-dose inhalers or ? 800  $\mu\text{g}/\text{j}$  of budesonide or ? 500  $\mu\text{g}/\text{d}$  of fluticasone propionate) and of inhaled long-acting  $\beta_2$  agonists, administered:

either in the form of two specialties using one or two of the following inhalers: Aerolizer®, standard metered-dose inhaler, Autohaler®, Diskus®, Turbuhaler®,

or in the form of a fixed association using one of the following inhalers: standard metered-dose inhaler, Diskus®, Turbuhaler®.

- Patients with FEV measurement in the previous month.
- Patients who brought their inhaled steroid treatment and inhaled long-acting  $\beta$ 2-agonist at the time of consultation.
- Patients in couple whether or not married
- Patients and relatives agree to participate

Patient non-inclusion Criteria :

- Patients with a non-asthmatic OCPD.
- Patients who had inhaled steroids or LABA treatment change in the previous three months (add-on or change of drug, posology change).
- Patients and relatives refusing to participate to the study
- Parents/ those close unable to complete a self-questionnaire.
- Patients who do not live as a couple

## **ESCAL - SURVEY ON HEALTH AND EATING BEHAVIOURS IN MARTINIQUE**

Last update : 09/05/2017

Head :

Quénel Philippe, Directeur du LERES - IRSET U1085 Inserm

Responsible organization :

Cellule interrégionale d'épidémiologie (Cire) Antilles-Guyane  
Observatoire de la Santé de la Martinique

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

The initial objectives of the health and eating habits survey were to identify: - The frequency of major diseases or their risk factors that affect the population of Martinique; - The nutritional status of the population and food intake in relation to potential risk factors; - Population habits in terms of food choices and frequency of consumption; - the population's habitual access to food.

### **Inclusion criteria**

- 3 years and over - living in Martinique for 12 months preceding the survey and at the time of the survey

## **HEPATYS - COHORT STUDY ON MANAGEMENT AND OUTCOME OF PATIENTS WITH CHRONIC HEPATITIS C TREATED WITH PEG-IFN-ALPHA2A IN COMBINAISON WITH RIBAVIRIN IN HOSPITAL-BASED OR PRIVATE OFFICE-BASED**

Last update : 09/05/2017

Head :  
Pinta Alexandrina, ROCHE

Responsible organization :  
ROCHE

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

Describe the management and outcome of patients with chronic Hepatitis C treated with PEG-IFN-alpha2a in combination with ribavirin in hospital-based or private office-based

### **Inclusion criteria**

Patients suffering from chronic hepatitis C for whom a treatment via Pagasys was decided and then initiated, followed in a hospital and general environment

Patients who have been informed of the study orally and in writing

## **EMU - CROSS-SECTIONAL STUDY ON MIGRAINE PATIENTS IN EMERGENCY SERVICE DEPARTMENTS: DIAGNOSIS AND TREATMENT**

Last update : 01/01/2018

Head :  
Grun-Overdyking Aurélie, Pfizer

Responsible organization :  
Pfizer

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Primary:

- To estimate the frequency of patients admitted with headaches and to estimate the frequency of patients diagnosed with migraine.
- To characterise patients diagnosed with migraines attending emergency services based on demographic data and migraine history (age, severity).
- To describe the treatment of patients diagnosed with a migraine during emergency service consultation.

Secondary:

- To describe the treatment of patients diagnosed with a migraine according to the type of migraine.
- To describe the frequency of patients diagnosed with a migraine and their treatment by emergency structure type.
- To describe treatment follow-up.

### **Inclusion criteria**

- Patients aged 18 or older.
- Patients deemed capable of answering the questions.
- Patients agreeing to participate in the survey.
- Patients admitted to emergency department

## **MENUI - COHORT OF CHILDREN WITH ISOLATED NOCTURNAL ENURESIS : A STUDY OF THE SAFETY OF MINIRINMELT IN ACTUAL PRESCRIPTION**

Last update : 01/01/2020

Head :

Niez Philippe, Laboratoire FERRING S.A.S

Responsible organization :

Laboratoire FERRING SAS

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

Show that the two oral forms (tablet and lyophilisate) of desmopressin have a similar safety profile and in particular in terms of frequency of the symptoms of alarms of an intoxication via water, in actual prescription situations, i.e. at general practitioners.

### **Inclusion criteria**

Criteria for inclusion:

- patient having an isolated nocturnal enuresis, defined by the number of wet nights per week,
- patient aged 6 to 18 years,
- patient in which the family doctor has decided to prescribe a treatment via desmopressin, or Minirin® tablets or Minirinmelt® lyophilisate.

Criteria for non-inclusion:

- patient who has already received prior treatment via desmopressin regardless of its form,
- patient who has a treatment in progress via desmopressin,
- patient participating in a therapeutic study.

## **- PREVALENCE OF GASTROESOPHAGEAL REFLUX DISEASE (GERD) DURING CHILDHOOD IN FRANCE**



Last update : 01/01/2018

Head :

Piedbois Pascal

Thomas-Delecourt Florence, AstraZeneca

Responsible organization :

ASTRAZENECA

## **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To assess prevalence of gastroesophageal reflux during childhood and teenage years in France

### **Inclusion criteria**

Children / teenagers from 0 to 17 years of age

## **- MANAGING FEVER IN CHILDREN: A NATIONAL SURVEY OF PARENTS' KNOWLEDGE AND PRACTICES IN FRANCE**

Last update : 06/01/2018

Head :

Chalumeau Martin, U953 Recherche épidémiologique en santé périnatale et santé des femmes et des enfants  
Équipe/activité : Epidémiologie clinique et organisation des soins

Responsible organization :

Institut National de la Santé et de la Recherche Médicale - Inserm

Assistance Publique - Hôpitaux de Paris Saint-Vincent-de-Paul AP-HP

## **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Study parents' knowledge and practices and their determinants in managing fever symptoms in children in France as compared with current recommendations.

### **Inclusion criteria**

man or woman - children between 1 month and 12 years old - having fever since less than 48h

# ADHOC - OBSERVATIONAL STUDY ON THE VARIOUS ARRANGEMENTS FOR CARING FOR PATIENTS WITH TYPE-2 DIABETES TREATED BY ASSOCIATING ORAL DIABETES MEDICATION AND INSULIN

Last update : 01/01/2018

Head :  
VIGNAL Franck, Sanofi Aventis

Responsible organization :  
Sanofi-aventis France

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Describe the various arrangements for caring for patients with type-2 diabetes treated by associating oral diabetes medication and insulin as well as their repercussion on HbA1c

## Inclusion criteria

man or woman, over the age of 18 years, having type-2 diabetes treated by the association of oral diabetes medication and insulin for at least 6 months, accepting to participate in the study

# PREFACE - STUDY ON THE PREVALENCE OF FACIAL LIPOATROPHY IN PATIENTS INFECTED BY HIV VIRUS RECEIVING ANTIRETROVIRAL DRUGS - FRAN08-005

Last update : 07/01/2019

Head :  
Cohen-Codar Isabelle

Responsible organization :  
ABBOTT France

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Describe the prevalence of the facial lipoatrophy in patients infected by HIV virus and receiving antiretroviral drugs. Describe the prevalence of lipoatrophies and lipodystrophic mixed syndromes.

Evaluate the score of quality of life, using the patients multi-dimensional auto-questionnaire and specific ABCD "Assessment of Body Change and Distress".

### **Inclusion criteria**

Patient infected by HIV virus, receiving antiretroviral drugs for at least 1 month and coming to consultation at his usual follow-up center, during the period of the study, obtaining of the written consent for the collect and exploitation of his personal data.

## **FLUREC - RECURRING INFLUENZA VIRUS INFECTIONS AND IMPLICATIONS IN TERMS OF EPIDEMIC RECURRENCES AND PANDEMIC RISK.**

Last update : 01/01/2020

Head :

Carrat Fabrice, UMR-S 707 / Epidémiologie des maladies infectieuses et modélisation  
De Lamballerie Xavier, UMR-D 190, Emergence des pathologies virales

Responsible organization :

APHP

### **Type of database**

Study databases

### **Study databases (details)**

Case control study

### **Main objective**

To explore cross-protective immunity involving different A/H1N1 viruses (seasonal and pandemic) and the seasonal A/H3N2 virus through data gathered from this study. This study also aims to identify the epidemiological, immunological and virological factors that determine the risk of recurring influenza infections as well as characterise the population's herd immunity and its evolution.

### **Inclusion criteria**

The "influenza" group comprises of patients enrolled after general medical consultation for influenza-like illness during seasonal epidemic. The "non-influenza" group comprises of patients enrolled after general medical consultation during seasonal epidemic for a reason other than influenza, yet that was acute in nature e.g. infectious diarrhoea, minor trauma.

## **CEPHEUS - ASSESSMENT OF LIPID-LOWERING TREATMENT IN FRANCE**

Last update : 09/05/2017

Head :

Ferrières Jean, Département d'épidémiologie INSERM UMR1027

Responsible organization :

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Determine the proportion of patients on lipid-lowering drugs who reach the LDL-C goals recommended in guidelines.

## Inclusion criteria

man or woman - adult - treated with lipid-lowering drugs for at least three months, with no dose adjustment for a minimum of six weeks

# ABSINTHE - OBSERVATIONAL STUDY OF THE THERAPEUTIC STRATEGY IN THE MANAGEMENT OF ACUTE SINUSITIS IN PRIMARY CARE

Last update : 09/05/2017

Head :

Moore Nicholas, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Responsible organization :

INSERM

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

The study objectives were to describe the management of acute sinusitis in real-life conditions of prescription and to evaluate the effectiveness of the initial therapeutic strategy.

## Inclusion criteria

Patient consulting for acute sinusitis diagnosed according to criteria of the practitioner; Patient without a previous episode of acute sinusitis during the two months preceding diagnosis; Patient aged 18 years and over; Patient agreeing to participate in the study; Patient not included in a clinical trial (Huriet-Sérusclat); Patient who may be followed for two months; Patient showing no severe active disease (life-threatening in the next three months).

# ESCALE - CASE-CONTROL STUDY ON CHILDHOOD CANCERS

Last update : 09/05/2017

Head :

CLAVEL Jacqueline, UMR-S 1153 Equipe 7 / Épidémiologie des cancers de l'Enfant

Responsible organization :

Institut National de la Santé et de la Recherche Médicale

## **Type of database**

Study databases

## **Study databases (details)**

Case control study

## **Main objective**

To investigate the influence of the following on childhood cancer risk: pregnancy outcome, child's characteristics at birth, child's environment in utero or during childhood, parents' profession and lifestyle, successive domicile communities, family history of cancer, birth defects, or autoimmune diseases. To identify genetic predispositions capable of modulating the function of these factors, including polymorphisms in genes involved in the transport and metabolism of xenobiotics, DNA repair, immune response, or in embryonic development and cell differentiation. To investigate the interactions between these polymorphisms and environmental factors.

## **Inclusion criteria**

- Men or women - under 15 years old

# **SIRS - HEALTH, INEQUALITIES AND SOCIAL RUPTURES**

Last update : 01/01/2019

Head :

Chauvin Pierre, UMRS 707, equipe ds3 - UPMC

Responsible organization :

INSERM

## **Type of database**

Study databases

## **Study databases (details)**

Longitudinal study (except cohorts)

## **Main objective**

General objective: to investigate the impact of living conditions and social integration on health and health care usage. Secondary objective: to investigate territorial factors (social, urban and available services) in relation to health and health care usage.

## Inclusion criteria

Voluntary French-speaking adults

# SURVSEIN - STUDY ON LONG-TERM IATROGENIC EFFECTS FOR WOMEN TREATED FOR BREAST CANCER BETWEEN 1954 AND 1985 AT THE INSTITUT GUSTAVE ROUSSY

Last update : 09/05/2017

Head :

De Vathaire Florent, U1018 CESP équipe 3

Rubino Carole, U1018 CESP équipe 3

Responsible organization :

Institut National de Santé et Recherche Médicale

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

SurvSein is a prospective and retrospective cohort study of women treated for breast cancer between 1954 and 1983 at the Institut Gustave Roussy in order to measure the iatrogenic effects of long-term treatment.

## Inclusion criteria

Subjects of female gender Women aged 18 or over Women treated for breast cancer

# DEFI - DETERMINATION OF EPIDEMIOLOGY OF FIBROMYALGIA

Last update : 01/01/2019

Head :

Millet Véronique, Pfizer

Responsible organization :

Pfizer

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

Main:

- Evaluate the prevalence of fibromyalgia in the general population, in five "French regions", by identifying:

? in a first step, by telephone, potential or known fibromyalgia patients,

? then in a second step, from among these patients, those that have fibromyalgia validated by a rheumatologist.

Secondary:

- Describe the characteristics of the patients with a diagnosis of fibromyalgia validated by a rheumatologist.

- Describe the characteristics of patients for whom this diagnosis has not been validated by the rheumatologist.

- Compare the characteristics of these two groups of patients to one another and each of them with the characteristics of the general population of the study.

### **Inclusion criteria**

- Patients aged 18 years or older.

- Patients deemed apt to answer the questions.

- Randomly selected patients who accept to participate in the survey (in the event of a refusal, the selected person was not replaced within the home or in the study)."

## **- MANAGEMENT OF HIGH CARDIOVASCULAR RISK PATIENTS TREATED WITH STATINS IN FRANCE**

Last update : 01/01/2019

Head :

Piedbois Pascal

Thomas-Delecourt Florence, AstraZeneca

Responsible organization :

ASTRAZENECA

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To describe utilisation of newly prescribed statins in French primary care

### **Inclusion criteria**

Any patient for whom a treatment with statin was initiated for at least 3 months and less than 6 months

## **E-TUD - TRANSVERSAL STUDY ON MEDICAL MANAGEMENT OF GLAUCOMA /OCULAR HYPERTENSION**

Last update : 01/01/2019

Head :  
TROY Sylvie, Pfizer

Responsible organization :  
pfizer

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Primary:

The main objective of this survey is to describe the diagnostic and therapeutic care for glaucoma and/or ocular hypertensions (defined by an intraocular pressure  $\geq 21$  mm Hg) examined on the day of the survey.

Secondary:

The secondary objectives are:

- The description of the satisfaction and of the observance perceived by the doctors according to the treatments,
- The diagnostic techniques used: visual fields, ocular imaging and gonioscopy,
- The risk factors for patients in routine practice,
- The exercise mode for ophthalmologists in caring for glaucoma and/or ocular hypertensions.
- Putting the data into relation with the data from the survey conducted in 2003.

## Inclusion criteria

All of the patients having glaucoma or an OHT coming to a consultation in ophthalmology on the given day were included in the survey, if they met the following conditions:

- at least 18 years of age,
- diagnosis of glaucoma, or OHT already established at the time of the consultation.
- were informed about the survey and gave oral agreement for the use of the data concerning them.

# ECRHS - EUROPEAN COMMUNITY RESPIRATORY HEALTH SURVEY

Last update : 09/05/2017

Head :  
Leynaert Benedicte, U 700  
Pin , PNEUMOLOGIE INFANTILE DÉPARTEMENT DE PÉDIATRIE  
Neukirch , U 700

Responsible organization :  
Institut National de la Santé et de la Recherche Médicale

## Type of database

Study databases

## Study databases (details)



Cohort study

## **Main objective**

General objective: To investigate the changes in lung function as well as the onset and development of allergic and non-allergic asthma and chronic obstructive bronchopneumopathy (COBP) in adults as they age. Secondary objectives: - To study the onset and development of respiratory allergies and COBP in men and women from the general population, and to research associations with lifestyle and environmental factors, taking into account interactions between these factors and other personal and biological determinants. - To research associations with sensitisation to allergens (atopy), respiratory allergies, accelerated lung function decline and the development of COBP. - To investigate the natural history of allergic and non-allergic asthma (incidence, remission, severity) according to age and gender. - To evaluate the effects of obesity and importance of weight loss and physical activity on asthma severity and decline in lung function, as well as investigate the role of sex hormones and hormones linked to obesity such as leptin.

## **Inclusion criteria**

Men and women from the general population between 20 and 44 years old in 1990-92

# **ASSESS - ASSESSMENT OF SYSTEMIC INVOLVEMENT AND PROGRESSION FOR PATIENTS WITH PRIMARY SJOGREN'S SYNDROME**

Last update : 01/01/2019

Head :

Gottenberg Jacques-Eric, U1109 Immuno-rhumatologie moléculaire

Responsible organization :

CHU Strasbourg

## **Type of database**

Study databases

## **Study databases (details)**

Longitudinal study (except cohorts)

## **Main objective**

To prospectively determine for the first time the prevalence and risk factors for systemic complications and lymphoma in primary Sjögren's syndrome.

## **Inclusion criteria**

- Men or women - adults - patients in tertiary centres for autoimmune diseases

# **ADEOS - ADHERENCE EVALUATION OF OSTEOPOROSIS TREATMENT**

Last update : 01/01/2018

Head :

Leclerc-Zwirn Christel, Laboratoire GSK

Responsible organization :

Laboratoire GSK

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

Finalize and validate a self-administered instrument developed to evaluate adhesion to the treatment of postmenopausal osteoporosis

## **Inclusion criteria**

Woman;

Over the age of 50 years;

Patient to whom has been prescribed one of the following osteoporosis treatments in the six months prior to the inclusion starting date:

bisphosphonates, selective estrogen receptor modulators (SERM), or strontium ranelate;

Patient fluent in French and who has the cognitive and functional ability required to complete the self-questionnaires included in the study on their own;

Informed patient who has given their oral consent to participate in this study

# **B009 - FOLLOW-UP STUDY OF PATIENTS TREATED WITH DROTRECOGIN ALFA (ACTIVATED) (XIGRIS®) IN FRANCE**

Last update : 01/01/2019

Head :

Laboratoire , Eli Lilly France

Responsible organization :

Eli Lilly

## **Type of database**

Study databases

## **Study databases (details)**

Longitudinal study (except cohorts)

## **Main objective**

Primary objective: evaluate 1-month mortality (at 28 days and 31 days) of patients treated with drotrecogin alpha (activated) in intensive care for severe sepsis and describe causes of death.

Secondary objectives: characteristics of patients and conditions of use.

## **Inclusion criteria**

All adult patients treated with drotrecogin alpha (activated) in France in the course of routine care in intensive care services.

## **- HEALTH BAROMETER**

Last update : 01/01/2020

Head :  
BECK François

Responsible organization :  
INPES

### **Type of database**

Study databases

### **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

### **Main objective**

Follow the evolution of some basic indicators about behavior and attitudes towards health on general population. Follow the evolution by age group, gender, sub-populations? according to the large size of the sample. Feed the different INPES information programs concerning behaviors, opinion and attitudes towards health as well as determinants (associates factors) of these opinions.

Study, thanks to the large size of the sample, sub-populations specificities (young people, old people, students, unemployed people) and unusual behaviors.

Study the links between representations and behaviors concerning health

(the difference between barometers and other health surveys is that they are surveys about prevention practices and lifestyles concerning health factors. There is no attempt to measure exactly health conditions. A relevant part of the questionnaire concerns sensitive topics such as addictions, mental health, sexual behaviors, etc.)

Being complementary to other big health surveys (avoid repetition, research coherence), in a collaborative effort.

Participate, thanks to the large size of the base, to the production of regional data, in collaboration with the ORS.

In the dimension study/observation, it will also be a point of support for the evaluation of public politics: indicators of the follow of Public Health laws, assessment of government plans (addictions plan, mental health plan, ageing well?)

### **Inclusion criteria**

Question people belonging to households:

- Equipped with a fixed telephone line (white, orange or red list)
- Equipped with just a mobile phone
- Full unbundled access

# CESAME - CANCER AND INCREASED RISK ASSOCIATED WITH INFLAMMATORY BOWEL DISEASE IN FRANCE

Last update : 12/01/2017

Head :

Beaugerie Laurent, UMRS-S 707

Responsible organization :

INSERM - Institut National de Santé et Recherche Médicale

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

General objective: To test the hypothesis that the use of immunosuppressants (IS) for inflammatory bowel disease (IBD) leads to an increased risk of lymphoma in a cohort population. Secondary objective: to assess - relative risk associated with IS treatment for all cancers, to assess the standardised incidence rate for all other cancers observed during IBD; -The increased risk of colorectal cancer associated with extensive and former colonic inflammation for ulcerative colitis (UC) and Crohn's disease: - Increased risk of ileal adenocarcinoma associated with chronic ileal inflammation in Crohn's disease; - Evaluation of mortality of patients with IBD diseases and risk factors; - Protective role of salicylates, pregnancy and malformation syndromes.

## Inclusion criteria

Subjects with IBD (Crohn's disease, ulcerative colitis, and indeterminate colitis) defined according to the usual diagnostic criteria, who attended consultation at least once between May 2005 and May 2006.

# TROPHEE - BENIGN PROSTATIC HYPERPLASIA: MANAGEMENT AND MEDICAL NEEDS NOT COVERED IN GENERAL PRACTICE IN FRANCE

Last update : 01/01/2020

Head :

Leclerc-Zwirn Christel, Laboratoire GSK

Responsible organization :

LABORATOIRE GSK

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

To describe the general practitioner long term management plan for patients with symptoms of BPH in France

## Inclusion criteria

- ? Patients over 50 years old
- ? Patients with a diagnosis of BPH for at least 1 year
- ? Patients currently treated with at least one of the treatments for symptoms of BPH

## EPIMAN-II-MILLE - LONGITUDINAL STUDY ON HOSPITALISED PATIENTS WITH ACUTE MANIA PERTAINING TO PRIMARY BIPOLAR DISORDER

Last update : 09/05/2017

Head :

Hantouche Elie, Centre de l'Humeur, département de psychiatrie adulte, hôpital Pitié-Salpêtrière  
Azorin Jean-Michel

Responsible organization :

Centre des Troubles Anxieux et de l'Humeur - CTAH

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

To demonstrate the influence of temperament on the expression of mania; particularly the link between mixed mania and depressive temperament.

## Inclusion criteria

- male and female
- adult
- hospitalised and presenting with acute mania pertaining to primary bipolar disorder

## O-PREDICT - OBSERVATOIRE DE LA PRISE EN CHARGE ET DE L'ÉVALUATION DU RISQUE D'INSUFFISANCE CARDIAQUE CHEZ LES PATIENTS HYPERTENDUS

Last update : 09/05/2017

Head :

Hagège Albert, Inserm U 633Pôle cardiovasculaire, hôpital européen Georges-Pompidou

Responsible organization :

Assistance Publique - Hôpitaux de Paris

## Type of database

Bases de données issues d'enquêtes

## Study databases (details)

Etudes transversales non répétées (hors enquêtes cas-témoins)

## Main objective

Évaluer les composantes du risque d'insuffisance cardiaque chez des patients hypertendus vus en médecine générale.

Comparer les estimations par le médecin du risque d'insuffisance cardiaque et du risque cardiovasculaire global par rapport au risque calculé par des échelles validées.

Évaluer la concordance entre ces deux estimations.

## Inclusion criteria

- homme ou femme
- adulte
- patient hypertendu sans insuffisance cardiaque vu en consultation

# I-NORM - MANAGEMENT BY FRENCH GENERAL PRACTITIONERS OF UNCONTROLLED HYPERTENSIVE PATIENTS TREATED BY DUAL THERAPY

Last update : 01/01/2018

Head :

Herrmann Marie-Annick, Bristol-Myers Squibb

Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :

Bristol-Myers Squibb France (BMS)

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Describe medical strategies used by general practitioners for uncontrolled hypertension despite dual-therapy: the I-NORM survey

## Inclusion criteria

Register: Patient  $\geq 18$  years, seen in routine consultation, monitored for HBP, known for more than 3 months, treated via fixed or free dual therapy for his HBP, dual therapy initiated by the participating doctor or the corresponding specialist.

Study: pt from the register, uncontrolled HBP (SBP  $\geq 140$  mmhg and /or DBP  $\geq 90$  mmhg all comer pt or

SBP  $\geq$ 130 mmhg and /or DBP $\geq$ 80mmhg diabetic pt or with kidney failure), not participating into a clinical trial

## **MANIF-2000 FOLLOWED BY MANIF-2 - COHORT OF HIV-SEROPOSITIVE INDIVIDUALS INFECTED BY USE OF INTRAVENOUS DRUGS: PROSPECTIVE STUDY IN MARSEILLE, AVIGNON, NICE AND ILE-DE-FRANCE**

Last update : 09/05/2017

Head :

Carrieri Maria-Patrizia, ORS PACA - INSERM U912

Obadia Yolande, ORS PACA

Responsible organization :

OBSERVATOIRE REGIONAL DE LA SANTE PACA

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

General objective: MANIF 2000 is a cohort of HIV-1 seropositive patients that have been infected through intravenous drug use. Patients should have an initial CD4 count of  $> 300/\text{mm}^3$ . The initial aim of this study was to compare HIV progression according to whether patients were still active addicts, substituting or sober. This goal was revised in light of the changes in HIV history with the release of highly active combination therapy and the treatment of addicts with buprenorphine substitution. MANIF 2000 has become an observational cohort study focused on risk-related sexual behaviours associated with drug addiction, as well as adherence to therapy and interaction between substitution and treatment. Secondary objectives: - To determine the impact of depression and HIV treatment adherence on clinical progression; - to determine the impact of retention in substitution treatment on virological success.

### **Inclusion criteria**

HIV-1 or HIV-2 seropositive patients that have been infected through intravenous drug use with an initial CD4 count greater than  $300/\text{mm}^3$  and at clinical stage A or B.

## **YOUNG-THYR - RISK FACTORS OF DIFFERENTIATED THYROID CANCER IN YOUNG PATIENTS IN EASTERN FRANCE: A CASE-CONTROL STUDY**

Last update : 01/01/2020

Head :

De Vathaire Florent, Equipe 3 Épidémiologie des cancers : radiocarcinogénèse et effets iatrogènes des traitements UMR 1018 Center for research in epidemiology and population health (CESP)

Responsible organization :

## Type of database

Study databases

## Study databases (details)

Case control study

## Main objective

To highlight genetic and environmental factors that contribute to the onset of differentiated thyroid cancer in children and young adults. To study the potential consequences of the Chernobyl accident.

## Inclusion criteria

- Men or women - residing in Eastern France

# **SPRIK - GENERAL PRACTITIONER/PAEDIATRICIAN-BASED SURVEILLANCE TO ESTIMATE THE DISEASE BURDEN OF ROTAVIRUS GASTROENTERITIS IN CHILDREN <5 YEARS OF AGE IN FRANCE**

Last update : 01/01/2018

Head :

Leclerc-Zwirn Christel, Laboratoire GSK

Responsible organization :

Laboratoire GSK

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Estimate the incidence of RVGE leading to a consultation of a general practitioner or pediatrician in a monitored population of children under the age of 5 years.

## Inclusion criteria

? Any child under the age of 5 years from the monitored population who spontaneously consults an investigating center for a acute gastroenteritis

# **LAPS - FEMALE PATIENTS WITH BREAST CANCER TREATED WITH LAPATINIB IN THE CONTEXT OF AN EXPANDED ACCESS PROGRAM (ATU): DESCRIPTION OF THE CARE TRAJECTORY AND CLINICAL COURSE**



Last update : 01/01/2020

Head :

Leclerc-Zwirn Christel, Laboratoire GSK

Responsible organization :

Laboratoire GSK

## Type of database

Study databases

### Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

### Main objective

Describe the care pathways of patients with breast cancer who have received and/or are receiving treatment with lapatinib under Temporary Use Authorization (TUA)

### Inclusion criteria

Patient who has received lapatinib for the treatment of breast cancer within the framework of a TUA between January 1, 2007 and 3 months prior to the beginning of the study

## **EVASCAN - ETUDE LONGITUDINALE SUR DES PATIENTS PRÉSENTANT UNE CORONAROPATHIE SUSPECTÉE OU CONNUE, STABLE OU STABILISÉE ET NÉCESSITANT UNE CORONAROGRAPHIE : ÉVALUATION MÉDICO-ÉCONOMIQUE DU SCANNER MULTICOUBE**

Last update : 01/01/2019

Head :

Durand-Zaleski Isabelle, Directrice de l'Unité de recherche clinique en économie de la santé d'Ile-De-France Responsable du service de Santé Publique (Groupe hospitalier Albert Chenevier- Henri Mondor)

Darlington Meryl, Économiste de la santé de l'Unité de recherche clinique en économie de la santé d'Ile-De-France

Responsible organization :

Assistance Publique - Hôpitaux de Paris

## Type of database

Bases de données issues d'enquêtes

### Study databases (details)

Etudes longitudinales (hors cohortes)

### Main objective

Valider la valeur diagnostique du scanner coronaire multicoupe dans une population de patients stables ressentant des douleurs thoraciques évoquant une coronaropathie et chez lesquels une coronarographie

est nécessaire.

### **Inclusion criteria**

- homme ou femme
- adulte
- patient présentant une coronaropathie suspectée ou connue, stable ou stabilisée et nécessitant une coronarographie avec la possibilité de la différer d'au moins 48 heures.

## **IDALGIA - STUDY OF PREDICTIVE FACTORS OF THE INDUCED PAIN LINKED TO GESTURES IN RHEUMATOLOGY**

Last update : 09/05/2017

Head :

Payen-Champenois Catherine, Bristol-Myers Squibb

Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :

Bristol Myers Squibb (BMS)

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Evaluate the frequency of the various types of gestures and the intensity of the associated pain

Evaluate the various therapeutic preventive and curative approaches for pain linked to gestures in rheumatology

### **Inclusion criteria**

For the gesture observatory: adult patient consulting in rheumatology for which the rheumatologist deems that it is necessary to immediately perform a diagnostic or therapeutic gesture

For the Study portion: patient eligible in the Observatory for which the rheumatologist has scheduled or has deemed immediately necessary, during this consultation, a one-off gesture (puncture and/or unilateral infiltration or an infiltration of the lumbar spine).

## **CASTOR - STUDY FOR THE COMPARISON OF THE RATE OF HOSPITALIZATION FOR RSV BRONCHIOLITIS BETWEEN PRETERM INFANTS BORN AT 32 WEEKS OF GESTATIONAL AGE OR LESS WITHOUT BRONCHOPULMONARY DYSPLASIA AND FULL-TERM INFANTS**

Last update : 01/01/2019

Head :

Miloradovich Tatiana, Abbott France

Responsible organization :  
Abbott France

## **Type of database**

Study databases

## **Study databases (details)**

Longitudinal study (except cohorts)

## **Main objective**

Compare the rate of hospitalization for bronchiolitis to RSV between premature children born  $\leq 32$  weeks of gestational age without bronchopulmonary dysplasia (BPD) and full-term children, during the 2008-2009 RSV season

## **Inclusion criteria**

- All aged less than 6 months at the beginning of the RSV epidemic season
- Very premature: born at 32 weeks of gestational age or less, without Bronchopulmonary Dysplasia (BPD)
- Full-Term: born between 39 and 41 weeks of gestational age

# **- ASSESSMENT OF CARDIOVASCULAR RISK IN PRIMARY CARE IN FRANCE**

Last update : 01/01/2020

Head :  
Piedbois Pascal  
Thomas-Delecourt Florence, AstraZeneca

Responsible organization :  
ASTRAZENECA

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

Evaluate the correlation between the cardiovascular risk estimated by the physician and the theoretical cardiovascular risk

## **Inclusion criteria**

Adults 50 years and older

# **- ETUDE LONGITUDINALE SUR DES PATIENTS SOUFFRANT DE DEPRESSION SEVERE : ETUDE DE L'IMPULSIVITE**

Last update : 09/05/2017

Head :

Falissard Bruno, Inserm U1018 - Centre de Recherche en Épidémiologie et Santé des Populations Équipe:  
Genre, Santé sexuelle et Reproductive  
Corruble Emmanuelle, INSERM U669 CHU DE BICETRE - SERVICE DE PSYCHIATRIE

Responsible organization :

INSERM - Institut National de la Santé et de la Recherche Médicale

## **Type of database**

Bases de données issues d'enquêtes

## **Study databases (details)**

Etudes transversales répétées (hors enquêtes cas-témoins)

## **Main objective**

Etudier la relation entre dépression sévère et impulsivité et le rôle de cette impulsivité dans les tentatives de suicide.

## **Inclusion criteria**

Patients nouvellement hospitalisés pour motif de dépression majeure (DSM-IV)

Critères d'exclusion: troubles bipolaires, troubles schizophréniques, abus d'alcool ou de drogue, troubles neurologiques.

# **ARISTOTE - USE OF FONDAPARINUX IN CURRENT CLINICAL PRACTICE FOR THROMBOPROPHYLAXIS FOLLOWING MAJOR ORTHOPEDIC SURGERY IN FRANCE**

Last update : 01/01/2019

Head :

Leclerc-Zwirn Christel, Laboratoire GSK

Responsible organization :

Laboratoire GSK

## **Type of database**

Study databases

## **Study databases (details)**

Longitudinal study (except cohorts)

## **Main objective**

? Describe the actual conditions of use of ARIXTRA® 2.5 mg in routine practice after an orthopedic surgical intervention.

? Observe the frequency of occurrence of VTEs during the 6 weeks following the initiation of the treatment

via ARIXTRA® 2.5 mg.

? Observe the frequency of occurrence of major bleeding during the 6 weeks following the initiation of the treatment via ARIXTRA® 2.5 mg.

### **Inclusion criteria**

? Patients of at least 18 years of age.

? Patients hospitalized in orthopedic surgery and for whom a treatment via ARIXTRA® 2.5 mg is initiated.

## **EMA - ÉTUDE LONGITUDINALE SUR LA RELATION ENTRE CRAVING ET TEMPS DE RÉACTION DANS LA DÉSINTOXICATION À L'ALCOOL. : UTILISATION DE L'EVALUATION ECOLOGIQUE MOMENTANÉE**

Last update : 09/05/2017

Head :

Falissard Bruno, Inserm U1018 - Centre de Recherche en Épidémiologie et Santé des Populations Équipe: Genre, Santé sexuelle et Reproductive  
Lukasiewicz Michael

Responsible organization :

INSERM - Institut National de la Santé et de la Recherche Médicale

### **Type of database**

Bases de données issues d'enquêtes

### **Study databases (details)**

Etudes longitudinales (hors cohortes)

### **Main objective**

Utiliser l'EMA (Evaluation Ecologique Momentanée) pour tester la relation entre le temps de réaction et le craving pendant la désintoxication alcoolique.

Décrire l'intensité, la fréquence et les facteurs prédictifs de craving, avec un enregistrement simultané d'autres états affectifs.

### **Inclusion criteria**

Dépendance à l'alcool (évaluation DSM-IV), hospitalisation pour désintoxication à l'alcool, âge compris entre 18 et 65 ans.

Les critères d'exclusion: autres abus de substances (à l'exception du tabac), diagnostic psychiatrique ou handicaps physiques incompatibles avec l'utilisation / compréhension de l'EMA, et l'absence de consentement verbal.

## **MDG 2007 - CHILDCARE AND SERVICES FOR YOUNG CHILDREN - METROPOLITAN FRANCE - 2007**

Last update : 11/01/2017

Head :

Legendre Emilie

Responsible organization :

La Direction de la recherche, des études, de l'évaluation et des statistiques (DREES)

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

To identify the different care solutions adopted by parents and to understand how they reconcile their professional and family life.

## **Inclusion criteria**

Households in metropolitan France that have at least one child under seven and a half years old (born 1st May 2000 and after).

# **CARDIORENAL - LONGITUDINAL OBSERVATIONAL STUDY ON CARDIOVASCULAR AND RENAL RISK IN PATIENTS WITH HYPERTENSION AND AT HIGH CV RISK**

Last update : 09/05/2017

Head :

Herrmann Marie-Annick, Bristol-Myers Squibb

Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :

Bristol-Myers Squibb France (BMS)

Sanofi Aventis

## **Type of database**

Study databases

## **Study databases (details)**

Cohort study

## **Main objective**

Evaluate, over a cohort of patients with a high cardiovascular risk according to the ANAES definition and monitored by a general practitioner, the incidence of the major cardiovascular event (MI, CVA and cardiovascular death) according to the various sub-classes of risks defined by ANAES in the "high risk" level

## **Inclusion criteria**

M/F  $\geq$ 18 years, examined in ambulatory consultation by general practitioners and cardiologists, having a treated essential HBP, or such that SBP/DBP  $>$  140/90 mmHg in the absence of treatment, monitored in a

normal manner by the investigating doctor

## **EVA - VASCULAR AGEING, DECLINE IN BRAIN AND COGNITIVE FUNCTIONS**

Last update : 09/05/2017

Head :

Berr Claudine, U 888

Responsible organization :

Institut National de la Santé et de la Recherche Médicale

### **Type of database**

Study databases

### **Study databases (details)**

Cohort study

### **Main objective**

General objective: to investigate vascular ageing, decline in brain and cognitive function and associated factors with a longitudinal follow-up study. Secondary objective: to determine the role of biological factors linked with oxidative stress and brain ageing.

### **Inclusion criteria**

Subjects from both genders born between 1922 and 1932 (59 to 71 years old at enrolment), included in electoral registers in Nantes, can speak French and living in Nantes.

## **BBMM IN POITOU-CHARENTES - BIEN BOUGER-MIEUX MANAGER (GET ACTIVE-EAT RIGHT) IN POITOU-CHARENTES**

Last update : 06/01/2020

Head :

Giraud Julien, Observatoire Régional de la Santé de Poitou-Charentes  
Roux , MÉDECIN CONSEILLER DU RECTEUR DE POITIERS

Responsible organization :

Rectorat de Poitiers

### **Type of database**

Study databases

### **Study databases (details)**

Cohort study

### **Main objective**

General objective: to halt the increase in the prevalence of excess weight and obesity in secondary

school children, who are monitored from 6th class (6e) to 3rd year (3e), in the Poitou-Charentes region - to assess physical and sports activities and the time 6th class (6e) and 3rd year (3e) students (n + 4) spend doing these activities; - to evaluate meal structure and food consumption over 24 hrs in accordance with the French National Nutrition and Health Program (PNNS).

### **Inclusion criteria**

Secondary school students from 32 institutions volunteering to participate in the programme (2 classes per institution). Secondary school students from 32 other randomly selected establishments (2 classes per institution)

## **ENGELA - ENVIRONMENT AND ADULT LYMPHOMA GENETICS**

Last update : 09/05/2017

Head :  
Clavel Jacqueline

Responsible organization :  
Institut National de la Santé et de la Recherche Médicale

### **Type of database**

Study databases

### **Study databases (details)**

Case control study

### **Main objective**

To analyse the relationship between lymphoid malignancies and family history of cancer.

### **Inclusion criteria**

- Men or women - aged between 18 and 75 years old

## **PASS-ARTHROSE - PATIENT ACCEPTABLE SYMPTOMATIC STATE**

Last update : 01/01/2020

Head :  
VIGNAL Franck, Sanofi Aventis

Responsible organization :  
Sanofi-aventis France

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)



## **Main objective**

Determine, in patients suffering from arthrosis of the lower limbs, the percentage of patients that find their pain "acceptable" at rest and when movement, after one week of treatment, according to the PASS concept (Patient Acceptable Symptomatic State).

## **Inclusion criteria**

Patients must:

- ? be ambulatory,
- ? be aged 50 years or older,
- ? consult for a defined gonarthrosis and/or coxarthrosis according to the clinical or radiological criteria of the American College of Rheumatology,
- ? be symptomatic with pain greater or equal to 3 on a numerical pain scale from 0 to 10, at rest or during movement, in the 24 hours preceding entry in the study,
- ? requiring the taking of an antalgic treatment for a duration of at least one week,
- ? be able and accept to complete a self-evaluation questionnaire,
- ? accepting to participate in the survey.

## **DRIM - FRENCH GENERAL PRACTITIONERS' WILLINGNESS TO PARTICIPATE IN PRIMARY CARE RESEARCH**

Last update : 09/01/2019

Head :

Letrilliart Laurent, Département de médecine générale

Responsible organization :

Université de Lyon 1

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

To assess French GPs' willingness to participate in primary care research as well as their expectations

## **Inclusion criteria**

General practitioners from the Rhône-Alpes region in France

## **- STUDY ON MENTAL HEALTH INDICATORS FOR HEALTH CARE PLANNING**

Last update : 09/05/2017

Head :

Kovess-Masfety Viviane, INSERM U705 et U669

Responsible organization :

Université Paris Descartes EHESP Hotel Dieu, Parvis Notre Dame 75004 Paris

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

- To use a specific epidemiological survey to generate simple indicators in order to - enable better allocation of resources in accordance with needs. Specific objectives: - To highlight different mental health issues and their risk factors. - To gather data in order to monitor these mental health issues. - To investigate health care access for mental health issues (medication/psychotherapy)

## **Inclusion criteria**

All individuals over 18 years of age from 4 participating regions (Ile de France, Upper Normandy, Lorraine, Rhône Alpes), in a household with a land-line phone or a mobile phone if there is no land-line phone (mono-equipped mobile phone).

# **- BAROMÈTRE CANCER**

Last update : 01/01/2020

Head :

Beck François

Gautier Arnaud

Responsible organization :

Institut national de prévention et d'éducation pour la santé (INPES)

Institut national du Cancer (INCa)

## **Type of database**

Study databases

## **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

## **Main objective**

Evaluate the behaviors, screening practices, representations on the care, attitudes and knowledge of individuals residing in France in terms of cancer.

## **Inclusion criteria**

Querying people in households:

- equipped with a landline (white, orange or red list) with geographical number (01 to 05)

- equipped solely with a mobile telephone

- equipped with a mobile telephone and a fully unbundled landline with a number starting with 08 or 09

---

# IMPACT - DETERMINANTS OF THE NON-CONTROL OF BLOOD PRESSURE IN OVERWEIGHTED TREATED HYPERTENSIVE PATIENTS

Last update : 01/01/2019

Head :  
Vignal Franck, Sanofi Aventis

Responsible organization :  
Sanofi-aventis France

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Find the determinants of the non-control of blood pressure in treated hypertensive patients who are overweight or obese compared to treated hypertensive patients of normal weight

## Inclusion criteria

Man or woman, over the age of 18 years, Having essential high blood pressure (HBP) treated for at least 1 year, Having a BMI < 25 kg/m<sup>2</sup> for patients of normal weight, Having a BMI ≥ 25 kg/m<sup>2</sup> for overweight or obese patients

# ARIANE - ARIXTRA AND PLATELET MONITORING (STUDY OF THE ACTUAL USAGE OF ARIXTRA® 2, 5 MG IN GENERAL PRACTICE

Last update : 01/01/2019

Head :  
Leclerc-Zwirn Christel, Laboratoire GSK

Responsible organization :  
Laboratoire GSK

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Evaluate and compare in general practice the level of prescription for platelet monitoring of patients

receiving Arixtra® 2.5 mg (fondaparinux sodium) or an LMWH (low-molecular-weight heparin) in venous thromboprophylaxis.

### **Inclusion criteria**

? Patient of at least 18 years of age

? Bedridden patient or with reduced mobility

Patient for whom a treatment via ARIXTRA® 2.5 mg or an LMWH is initiated within the framework of venous thromboprophylaxis

## **EDITH - OBSERVATIONAL STUDY TO EVALUATE EDUCATIONAL MODALITY OF TYPE 2 DIABETIC PATIENTS AT THE TIME OF INSULIN INITIATION BY GPS**

Last update : 01/01/2018

Head :

Vignal Franck, Sanofi Aventis

Responsible organization :

Sanofi-aventis France

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

describe the effects on the fasting glucose at 1 month for main methods of education of type-2 diabetes patients during the initiation of insulin therapy in general practice

### **Inclusion criteria**

man or woman, over the age of 18 years, having a type-2 diabetes responding to the criteria of HAS (French National Authority for Health), treated with oral diabetes medication at the maximum dose (or maximum tolerated dose where applicable), having needed the adding of a basal insulin initiated since at least 1 month and no more than 6 months (+ 1 week) before inclusion, reviewed 1 month after the initiation of insulin, during a routine visit in the framework of following his or her diabetes, accepting to participate in the study after information given by the observing doctor.

## **EULEV - EFFECTIVENESS AND USE OF LEVETIRACETAM IN REAL LIFE**

Last update : 10/26/2017

Head :

Fourrier-Reglat Annie, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen  
Moore Nicholas, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Responsible organization :

## Type of database

Study databases

### Study databases (details)

Longitudinal study (except cohorts)

### Main objective

The objectives of this study were to describe patients initiating treatment with levetiracetam, the prescribing patterns, and to evaluate the effectiveness of levetiracetam in a real-life situation in terms of treatment retention over one year.

### Inclusion criteria

Patients who initiated treatment with levetiracetam between 01/01/2005 and 31/08/2005 or between 01/01/2006 and 31/08/2006 (irrespective of whether or not treatment was continued), and having never previously been treated with levetiracetam; patient agreeing to participate; patient can be followed over the next 12 months; Patient not participating in a clinical trial (Huriet-Sérusclat).

## **EXTRA - OBSERVATOIRE SUR LA FRÉQUENCE DES MANIFESTATIONS EXTRA-ARTICULAIRES ET PLUS PARTICULIÈREMENT DE L'UVÉITE CHEZ LES PATIENTS PRÉSENTANT UNE SPONDYLARTHROPATHIE ET SUIVIS EN CONSULTATION DE RHUMATOLOGIE**

Last update : 01/01/2020

Head :

Thibout Emmanuel

Responsible organization :

Abbott

## Type of database

Study databases

### Study databases (details)

Longitudinal study (except cohorts)

### Main objective

Evaluation of the frequency of extra-articular manifestations and more specifically of uveitis in patients with a spondylarthropathy

### Inclusion criteria

Patients suffering from a spondylarthropathy

## **- THE IMPACT OF MALE GONADAL DOSE OF IONISING**

# RADIATION AND PROBABILITY OF PREGNANCY

Last update : 09/05/2017

Head :

Slama Remy, U823 Centre de recherche institut Albert Bonniot Équipe: Épidémiologie environnementale appliquée à la reproduction et à la santé respiratoire

Responsible organization :

Institut National de la Santé et de la Recherche Médicale

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

To determine the effects of ionising radiation on male reproductive function. To assess the risks of ionising radiation on humans concerning the probability of pregnancy.

## Inclusion criteria

- Women - between 18 and 60 years old - recognised pregnancy between 1985 and 2000

# TACTIC - THE CLINICAL PRESENTATIONS OF HIGH BLOOD PRESSURE IN 2005 AND ITS CARE IN GENERAL PRACTICE

Last update : 01/01/2018

Head :

Millet Véronique, Pfizer

Responsible organization :

Pfizer

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

"Evaluate the distribution of the three clinical presentations of high blood pressure (family, metabolic or vascular hypertension) at general practitioners in 2005 in subjects treated or not treated for hypertension - Evaluate the care of hypertensive subjects according to the various clinical presentations of high blood pressure (family, metabolic or vascular hypertension) diagnosed by general practitioners."

## Inclusion criteria

"Male or female aged 18 years or older,

- Hypertensive subject consulting, regardless of the reason. The investigator should include 4 patients as follows:

- 2 hypertensive subjects treated for their hypertension 60 years or older
- 1 hypertensive subject treated for their hypertension under the age of 60 years
- 1 hypertensive subject not treated for their hypertension for at least 6 months.
- Subject accepting to participate in the survey."

## **- EVALUATION OF IMMUNOCHEMICAL FAECAL OCCULT BLOOD TESTS FOR COLORECTAL CANCER SCREENING**

Last update : 09/05/2017

Head :

Launoy Guy, Cancers & Populations

Responsible organization :

Institut National de Santé et Recherche Médicale

CHU Caen

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

To demonstrate the use of immunochemical faecal occult blood tests for colorectal cancer screening.

### **Inclusion criteria**

- Men or women - aged between 50 and 74 years old

## **CONNECT - OBSERVATIONAL STUDY OF ADHERENCE TO EUROPEAN CLINICAL PRACTICE GUIDELINES FOR THE MANAGEMENT OF ACUTE CORONARY SYNDROME IN REVASCULARIZED VERSUS NON REVASCULARIZED PATIENTS**

Last update : 09/05/2017

Head :

Herrmann Marie-Annick, Bristol-Myers Squibb

Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :

Bristol-Myers Squibb France (BMS)

### **Type of database**

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

### Main objective

Management of acute coronary syndromes and adequation to international guidelines at the acute phase and at hospital discharge, according to the realization or not of a percutaneous coronary revascularization

### Inclusion criteria

Patient  $\geq$  18 years, hospitalized in cardiology for ACS with or without ST-segment elevation, selected the day of hospital discharge

## EFG SENIOR - INFLUENZA BURDEN ASSESSMENT IN ADULTS AGED OF 65 YEARS AND MORE VISITING A GENERAL PRACTITIONER FOR ACUTE RESPIRATORY ILLNESS IN FRANCE

Last update : 01/01/2020

Head :

Leclerc-Zwirn Christel, Laboratoire GSK

Responsible organization :

LABORATOIRE GSK

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

### Main objective

Describe and compare the burden of acute respiratory infections (IRA) linked to influenza virus, in terms of morbidity and medical consumption, according to the vaccinal status, in those 65 years and older spontaneously consulting in general practice

### Inclusion criteria

? Patient aged 65 years or older

? Patient having an acute respiratory infection defined as a clinical presentation combining the abrupt appearance of respiratory signs (coughing, rhinitis, coryza) in the context of acute infection (fever, asthenia, headache, myalgia, etc.), in less than 48h.

? For patients 80 years and older, the clinical presentations can associate other general signs (mental confusion, dehydration, anorexia, digestive disorders, general malaise, body aches, headache) and respiratory signs (from rhinitis to pneumopathy)

## TRAJAN - ETUDE TRANSVERSALE CHEZ DES PATIENTES ATTEINTES DE CANCER DU SEIN : ÉVALUATION DU PARCOURS DE SOINS



Last update : 01/01/2020

Head :

Defossez Gautier, Unité d'épidémiologie, biostatistique et registre général des cancers de Poitou-Charentes

Responsible organization :

CHU de Poitiers

## **Type of database**

Bases de données issues d'enquêtes

## **Study databases (details)**

Etudes transversales non répétées (hors enquêtes cas-témoins)

## **Main objective**

Identifier les déterminants associés aux délais d'accès aux traitements des patientes atteintes de cancer du sein en appliquant un algorithme informatique sur les données collectées en routine par le registre des cancers de la région Poitou-Charentes.

## **Inclusion criteria**

- femme
- âgée de 85 ans et plus
- résidant en Poitou-Charentes
- diagnostiquée entre 2008 et 2010 pour un cancer du sein infiltrant non métastatique
- prise en charge thérapeutique partielle ou complète en région Poitou-Charentes
- absence de traitement concomitant pour un autre cancer

# **ATHMOS - VALIDATION OF THE ACT QUESTIONNAIRE IN GENERAL PRACTICE**

Last update : 01/01/2018

Head :

Pribil Céline, Laboratoire GSK

Responsible organization :

Laboratoire GSK

## **Type of database**

Study databases

## **Study databases (details)**

Longitudinal study (except cohorts)

## **Main objective**

Describe in actual practice the level of control of asthma using the ACT questionnaire (Asthma Control Test)

## **Inclusion criteria**

Male or female aged 18 years or older

Patient having asthma diagnosed at least 12 months ago

Patient informed of the objectives of the study and accepting the collection and the analysis of the data concerning him.

## **CEREPHY - BRAIN TUMOURS AND PLANT PROTECTION PRODUCTS**

Last update : 09/05/2017

Head :

Salamon Roger, INSERM U 897

Baldi Isabelle

Responsible organization :

ISPED- Institut de santé publique d'épidémiologie et de développement

## **Type of database**

Study databases

## **Study databases (details)**

Case control study

## **Main objective**

To investigate the relationship between exposure to pesticides and central nervous system disorders (brain and spinal cord)

## **Inclusion criteria**

Adults aged 16 years or more living in Gironde who have malignant or benign primitive brain tumours that were diagnosed between 1st May 1999 and 30th April 2001. Recurrence, secondary tumours, pituitary adenomas, asymptomatic tumours, incidentally discovered and vascular malformations were excluded from the study.

## **EULEVP - EFFECTIVENESS AND USE OF LEVETIRACETAM OF PEDIATRICS IN REAL LIFE**

Last update : 10/26/2017

Head :

Fourrier-Reglat Annie, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Moore Nicholas, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Responsible organization :

INSERM

## **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

#### **Main objective**

The objectives of this study were to describe the children initiating treatment with levetiracetam, the prescribing patterns, and to evaluate the effectiveness of levetiracetam in a real situation in terms of treatment retention for one year.

#### **Inclusion criteria**

Patient aged under 16 years; Obtainment of consent from the guardian of the child; patient having initiated treatment with levetiracetam between 01/10/2006 and 31/03/2007 (whether or not the treatment was continued), and having never been previously treated with levetiracetam; patient who may be followed over the following 12 months; Patient not participating in a clinical trial (Huriet-Sérusclat).

## **POSTEPI - MANAGEMENT OF OSTEOPOROSIS AND ASSOCIATED QUALITY OF LIFE IN POST MENOPAUSAL WOMEN**

Last update : 01/01/2018

Head :

Leclerc-Zwirn Christel, Laboratoire GSK

Responsible organization :

LABORATOIRE GSK

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

#### **Main objective**

In the framework of the new screening and reimbursement conditions for osteoporosis medications, describe the profile of postmenopausal women with osteoporosis and treated for less than 6 months

#### **Inclusion criteria**

Postmenopausal women diagnosed with osteoporosis less than 6 months ago via low BMD (Femoral total T score and/or spinal average  $<-2.5$ ) or on a fracture not linked to a major trauma or to a cancer and treated for less than 6 months

## **CALIPSO - PROGNOSTIC VALUE OF PERIPHERAL ARTERIAL DISEASE ON 1-YEAR ISCHEMIC OUTCOMES IN OUPATIENTS WITH A RECENT HISTORY OF ACS**

Last update : 01/01/2019

Head :

Herrmann Marie-Annick, Bristol-Myers Squibb

Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :

Bristol-Myers Squibb France (BMS)

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

Determine the prognostic value of PAD (symptomatic or asymptomatic) on 1-year ischemic outcomes (death, myocardial ischemic recurrence, stroke, revascularization, rehospitalization for ischemic events) in patients with a recent ACS.

## Inclusion criteria

Register part : M/F patient,  $\geq 18$  years, outpatient seen by private cardiologists, with a recent history of ACS with or without ST-segment elevation in the last 12 months.

Study Part : patients from the register (2 first consecutive pts in the register without PAD) + (2 first consecutive patients in the register with symptomatic PAD or asymptomatic PAD and  $ABI < 0.9$ )

# IMPACT-VECTOR - EVALUATION AND PREDICTION OF THE IMPACT OF VECTOR-BORNE DISEASES (MALARIA PLASMODIUM FALCIPARUM AND ARBOVIRUSES) AMONG FRENCH MILITARY PERSONNEL DURING SHORT-DURATION MISSIONS (4 MONTHS) IN INTER-TROPICAL AFRICA

Last update : 11/01/2017

Head :

Rogier Christophe, Unité de recherche en biologie et épidémiologie parasitaires (équipe membre de l'UMR 6236, unité de recherche en maladies infectieuses et tropicales émergentes - URMITE) Ministère de la Défense.

Responsible organization :

Université de la Méditerranée (Aix-Marseille II)

Centre National de la recherche scientifique (CNRS)

Institut de recherche pour le développement (IRD)

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

General objective: To identify human and environmental determinants of French military personnel exposure to plasmodium falciparum malaria during short-duration missions (4 months) in inter-tropical Africa. Secondary objective: - To identify human and environmental determinants of French military personnel exposure to arboviruses during short-duration missions (4 months) in inter-tropical Africa - To identify human and environmental determinants of French military personnel exposure to malaria vector bites (s.s. Anopheles gambiae) and arboviruses (Aedes aegypti) during short-duration missions (4 months) in inter-tropical Africa..

## Inclusion criteria

Belonging to a military company taking part in short-duration missions (4 months) in inter-tropical Africa. Agreeing to participate in the study (questionnaires and blood tests).

## FLVS I - FLEURBAIX LAVENTIE VILLE SANTÉ STUDY I

Last update : 09/05/2017

Head :

Charles Marie-Aline, INSERM U1018 CENTRE DE RECHERCHE EN ÉPIDÉMIOLOGIE ET SANTÉ DES POPULATIONS  
Borys Jean-Michel, PRIVE

Responsible organization :

Protéines

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

To answer the question: does nutritional education in school change family eating behaviours?

## Inclusion criteria

Families with school-age children in primary school in Fleurbaix and Laventie and case-control cities

## PRELUDE - STUDY OF FACTORS AFFECTING THE LIVES OF PEOPLE TREATED BY ANTI-HIV DRUGS IN LONG TERM - FRAN09-014

Last update : 07/01/2020

Head :

Cohen-Codar Isabelle

Responsible organization :

ABBOTT France

Abbott

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Describe the factors affecting the life of individuals treated in the long term by anti-HIV drugs.

Describe the factors affecting the life of individuals treated in the long term by anti-HIV drugs according to their age.

Evaluate the correlation between the age, duration of seropositivity, the duration under treatment and the disorders reported.

Compare the patient's view and that of his doctor concerning the factors described in the main objective.

## Inclusion criteria

\* HIV-infected patients, treated with antiretroviral agents for at least 5 years and examined in consultation in their normal care center for monitoring their infection

\* Obtaining written consent for the collection and use of personal data

# ACTUEL - ÉTUDE TRANSVERSALE SUR DES PATIENTS ADULTES DÉPRESSIFS UNIPOLAIRES : DÉPRESSION RÉCURRENTÉ ET ÉVÉNEMENTS DE VIE : DE LA CLINIQUE À DE NOUVEAUX MODÈLES EXPLICATIFS ?

Last update : 09/05/2017

Head :

Falissard Bruno , Inserm U1018 - Centre de Recherche en Épidémiologie et Santé des Populations Équipe:

Genre, Santé sexuelle et Reproductive

CORRUBLE Emmanuelle, U 669

Responsible organization :

INSERM - Institut National de la Santé et de la Recherche Médicale

## Type of database

Bases de données issues d'enquêtes

## Study databases (details)

Etudes transversales non répétées (hors enquêtes cas-témoins)

## Main objective

Etudier la relation entre le nombre d'épisodes dépressifs antérieurs et le poids des événements de vie dans le déclenchement de l'épisode dépressif actuel

## Inclusion criteria

Patients adultes dépressifs unipolaires

## **EPIPAP - ORAL DISEASE AMONG PREGNANT WOMEN AND PRETERM BIRTH**

Last update : 09/05/2017

Head :

Nabet Cathy, U1027 EPIDEMIOLOGIE ET ANALYSES EN SANTE PUBLIQUE : RISQUES, MALADIES CHRONIQUES ET HANDICAPS

Responsible organization :

Institut de la santé et de la recherche médicale - Inserm

### **Type of database**

Study databases

### **Study databases (details)**

Case control study

### **Main objective**

To study the relationship between maternal periodontitis and preterm birth, according to different causes of preterm birth

### **Inclusion criteria**

Adult, with at least 6 teeth, having given birth to an alive-born child between 22 and 36 completed weeks of gestation (case) or at 37 weeks or more (control), consenting

## **OMA - EVALUATION OF THE MANAGEMENT OF ACUTE OTITIS MEDIA IN CHILDREN AGED ? 6 YEARS IN FRANCE**

Last update : 01/01/2019

Head :

Leclerc-Zwirn Christel, Laboratoire GSK

Responsible organization :

Laboratoire GSK

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

Describe in general practice the care for AOM in children aged 0 to 6 years in France.

### **Inclusion criteria**

- ? Child aged 0 to 6 years
- ? Child with an episode of AOM diagnosed by a doctor
- ? Parent(s)/tutor(s) who accept the inclusion of the child in the study

## **GN-PROGRESS - STUDY OF GENETIC AND ENVIRONMENTAL FACTORS OF GLOMERULAR NEPHROPATHY PROGRESSION**

Last update : 09/05/2017

Head :

Stengel Bénédicte, INSERM U1018 CENTRE DE RECHERCHE EN ÉPIDÉMIOLOGIE ET SANTÉ DES POPULATIONS  
Rossert , U780

Responsible organization :

AP-HP

### **Type of database**

Study databases

### **Study databases (details)**

Cohort study

### **Main objective**

To investigate genetic and environmental factors of glomerular nephropathy progression.

### **Inclusion criteria**

IgA-type glomerulonephritis, focal and segmental hyalinosis or extramembranous glomerulopathy, having had a biopsy between 1994 and 2001

## **MELODIE - ELABORATION OF A PREDICTIVE CLINICAL SCORE TO EVALUATE THE ADHERENCE OF OBESE, TYPE 2 DIABETIC PATIENT TO THE PHYSICIAN'S RECOMMENDATIONS IN TERMS OF MEDICATIONS, EXERCISE AND DIET**

Last update : 01/01/2018

Head :

Vignal Franck, Sanofi Aventis

Responsible organization :

Sanofi-aventis France

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)



## Main objective

Using the cardiovascular risk factors of unbalanced obese type-2 diabetes patients, develop a clinical predictive score of the adherence of patients to the modifications in their therapeutic care and/or in their life hygiene that are prescribed for them

## Inclusion criteria

Man or woman over the age of 18 years; Type-2 diabetes treated via oral diabetes medication monotherapy; HbA1c  $\geq$  6.5%; BMI  $\geq$  30 kg/m<sup>2</sup>; Abdominal obesity defined according to the NCEP/ATPIII 2001 criteria: waist  $>$  88 cm for women and  $>$  102 cm for men.

## ALMA+ - ALUMINIUM IN WATER CONVEYANCE AND ALZHEIMERS DISEASE

Last update : 09/05/2017

Head :

Rondeau Virginie, Unité Inserm : U897

Responsible organization :

ISPED- Institut de santé publique d'épidémiologie et de développement

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

Study the relation between minerals in drinking water and especially aluminium, and the development of Alzheimers disease in 75 or more years old subjects. Verify if high concentrations of aluminium (more than 50mg/l of water) can improve the risk of dementia.

## Inclusion criteria

75 or more years old subjects, exposed to high concentrations of aluminium in drinking water (Al $>$ 50mcg/l), residents in 14 towns of Dordogne, different than the ones included in the PAQUID cohort. 75 or more years old subjects participating to the PAQUID cohort, residents in one of the non-exposed towns (Al $<$ 50mcg/l) in Dordogne. Exclusion criteria: subjects recently arrived in town. Inclusion period: May 1999 - January 2001.

## - PREVALENCE AND TIME TRENDS OF DISABILITIES IN SCHOOL-AGE CHILDREN

Last update : 09/05/2017

Head :

Grandjean Hélène, CJF 94-04

Responsible organization :

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

To determine time trends and prevalence of disabilities in school-age children.

## Inclusion criteria

- Men or women - born between 1976 and 1985 - whose parents lived in one of the départements where subject was included.

# **PRESERV - OBSERVATOIRE SUR LES STRATÉGIES DE PRISE EN CHARGE DE LA POLYARTHRITE RHUMATOÏDE (PR) DE MOINS DE 12 MOINS SELON LA PRÉSENCE OU NON D'ANTICORPS ANTI-CCP ET/OU DE ACTEUR RHUMATOÏDE.**

Last update : 01/01/2020

Head :

Thibout Emmanuel

Responsible organization :

Abbott France

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

Describe in routine practice the possible influence of the presence of the rate of Ac anti-CCP and/or that of the rheumatoid factor on the therapeutic choices made by rheumatologists in patients with a diagnosis of rheumatoid polyarthritis of less than 12 months

## Inclusion criteria

Patients suffering from rheumatoid polyarthritis and an anti-CCP test

# **ZEN - AN OBSERVATIONAL STUDY OF OLANZAPINE COATED AND ORODISPERSIBLE TABLETS EFFECTIVENESS IN SCHIZOPHRENIC AND BIPOLAR OUTPATIENTS**

Last update : 01/01/2019

Head :  
Médecin pharmacoépidémiologiste , Eli Lilly France

Responsible organization :  
Eli Lilly

## **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

The primary objective of this study is to assess the proportion of schizophrenic patients and the proportion of bipolar patients who discontinued olanzapine within 12 months of treatment in outpatient, ambulatory or community settings.

### **Inclusion criteria**

Patients aged 18 years and older,  
- suffering from bipolar disorders or schizophrenic disorders according to DSM-IV criteria,  
- initiating or having initiated a treatment via olanzapine in tablet or orodispersible form.

## **ATLANTIS - LONG TERM OBSERVATIONAL STUDY ON THE THERAPEUTIC MANAGEMENT OF KNEE OSTEOARTHRITIS BY GENERAL PRACTITIONERS**

Last update : 09/05/2017

Head :  
Payen-Champenois Catherine, Bristol-Myers Squibb  
Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :  
Bristol-Myers Squibb (BMS)

## **Type of database**

Study databases

### **Study databases (details)**

Cohort study

### **Main objective**

Evaluate, in general practice, the change in the first-line therapeutic strategy in patients suffering from knee osteoarthritis, over one year, specially:

- Evaluate the percentage of patients treated with each therapeutic modality at the end of the one year observation period.
- Describe the modifications implemented in the therapeutic management and identify the reasons

justifying these modifications.

### **Inclusion criteria**

1. Males or females 50 years or older suffering from symptomatic osteoarthritis of the knee
2. Not treated over the last 6 months for this pathology (non-pharmaceutical treatments and self-medication by analgesics or NSAIDs at low doses will be allowed).
3. For whom the physician will have decided to initiate a first-line treatment in line with the recommendations of EULAR 2000-2003 (paracetamol - systematically taken at a dose of 3 to 4g/d - as pharmacological monotherapy in combination with non-pharmaceutical treatments).

## **- HEALTH AND ENVIRONMENT BAROMETER**

Last update : 09/05/2017

Head :  
Menard Colette

Responsible organization :  
Institut National de Prévention et d'Education pour la Santé (INPES)

### **Type of database**

Study databases

### **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

### **Main objective**

Evaluate the representations, attitudes and behaviors of the population residing in France faced with environmental risks

### **Inclusion criteria**

Querying people in households:  
-equipped with a landline telephone (white, orange or red list)

## **PSOLEMEE - LOCALIZED PSORIASIS: A PHARMACO-ECONOMIC EVALUATION**

Last update : 09/05/2017

Head :  
Moore Nicholas, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Responsible organization :  
Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

### **Type of database**

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

### Main objective

The main objective of this study was to evaluate the cost of psoriasis care for patients corresponding to the target population of BETESIL®, i.e. those suffering from chronic plaque psoriasis, localized in areas difficult to treat (knees, elbows, and anterior surface of the tibia, not exceeding 5% of body surface area).

### Inclusion criteria

Patient aged 18 years or above; Patient with chronic plaque psoriasis; Patient with plaques located in places difficult to treat (knees, elbows, and anterior tibia, not exceeding 5% of the body surface area); Patient able to understand or complete the study questionnaires; patient who can be followed for 6 months; patient agreeing to participate in the study.

## CORONS - RISK OBSERVATORY FOR COHORT OF PERSONS REGISTERED WITH THE INDEPENDENT NATIONAL SOCIAL SECURITY FUND FOR MINES IN NORTHERN FRANCE

Last update : 03/01/2019

Head :

Flahault Antoine, UMR S 707

Responsible organization :

Institut National de la Santé et de la Recherche Médicale

### Type of database

Study databases

## Study databases (details)

Cohort study

### Main objective

General objective: to observe the risks for a cohort of persons registered with the Independent National Social Security Fund for Mines in Northern France.

### Inclusion criteria

All individuals registered with the Independent National Social Security Fund for Mines that have agreed to the usage of their medical data.

## C.A.R.O.L.E: ACTIVE NETWORK COORDINATION OF LONGITUDINAL EPILEPSY OBSERVATORY - LONGITUDINAL STUDY OF PATIENTS WITH EPILEPSY: MEDICAL AND SOCIAL OUTCOME STUDY

Last update : 09/05/2017

Head :

Jallon Pierre, Unité d'épileptologie et d'EEG

Responsible organization :  
Fondation française pour la recherche sur l'épilepsie

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

To assess the medical and social outcome of subjects with spontaneous epileptic seizures and to determine drug resistance predictors.

## Inclusion criteria

- men and women
- having experienced first epileptic seizure between 01/05/1995 and 30/06/1996

# I-SYPRES - A 6 MONTH FOLLOW-UP OF STUDY OF HOSPITALISED PATIENTS WITH CARDIAC FAILURE WITH PRESERVED SYSTOLIC FUNCTION (PSF-HF) IN CARDIOLOGY, INTERNAL MEDECINE ET GERIATRY

Last update : 01/01/2019

Head :  
Herrmann Marie-Annick, Bristol-Myers Squibb  
Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :  
Bristol-Myers Squibb (BMS)  
Sanofi Aventis

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

Determine the demographic characteristics and of the disease of patients hospitalized in the cardiology department for cardiac failure with intact systolic function

## Inclusion criteria

Register Portion: M/F  $\geq 18$  years, hospitalized for confirmed acute cardiac failure on the clinical symptoms and signs,

Study Portion: Patient in the register, diagnosed for HF-ISF with an EF >45% evaluated within 8 days after the date of hospitalization

## **CORINES - IDENTIFICATION OF ADRENAL INSUFFICIENCY AND / OR CUSHING'S SYNDROME IN PATIENTS TREATED WITH INHALED CORTICOSTEROIDS**

Last update : 09/05/2017

Head :

Molimard Mathieu, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen  
Moore Nicholas, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Responsible organization :

Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

### **Type of database**

Morbidity registers

### **Main objective**

The objectives were to identify cases of adrenal insufficiency and / or Cushing's syndrome in patients taking inhaled corticosteroids during the years 2000 to 2005 and describe the circumstances of occurrence of these events in order to propose recommendations to prevent them.

### **Inclusion criteria**

Children or adults treated with inhaled corticosteroids; Having presented symptoms suggestive of adrenal insufficiency and / or Cushing's syndrome during the years 2000 to 2005; First symptoms having appeared during treatment or within 3 months after discontinuation; Patients without adrenal insufficiency and / or Cushing's syndrome known before initiation of treatment with inhaled corticosteroids.

## **EDONIS - EPIDEMIOLOGICAL STUDY ON POST-SURGICAL NEUROPATHIC PAIN**

Last update : 06/01/2020

Head :

Duale Christian, CPC-CIC

Responsible organization :

CHU de Clermont-Ferrand

### **Type of database**

Study databases

### **Study databases (details)**

Cohort study

### **Main objective**

General objective: The EDONIS projects (PHRC national 2005) piloted by the Clermont-Ferrand Clinical Investigation Centre is a national multicentric epidemiological study whose primary objective is to know the prevalence of persistent pain at 3 and 6 months after surgery. This is the first prospective study of this size on chronic post-surgical pain which is a public health problem accounting for almost 15% of new consultations in specialist pain management institutions. Secondary objective: EDONIS also aims to identify pre-and peri-operative pain risk factors and to characterise its mechanisms with a focus on neuropathic factors. Eight types of conventional surgery were selected because (a) they are already known to induce persistent pain (thoracotomy, mastectomy, inguinal hernia repair), (b) they are frequently practised in France and chronic pain is an attribute (sternotomy, caesarean section, cholecystectomy, saphenectomy) or (c) surgery has a suspected risk of nerve injury (knee arthroscopy). There is also an associated ancillary genetic study.

### **Inclusion criteria**

Scheduled surgery identified as neuropathy-inducing

## **ESCAPE - ESTIMATED COST OF THE LOWER LIMB OSTEOARTHRITIS (KNEE ARTHRITIS AND COXARTHROSIS) IN THE GENERAL POPULATION AND IN THOSE TREATED BY SPA THERAPY**

Last update : 10/27/2017

Head :

Moore Nicholas, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Responsible organization :

INSERM

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

The main objective of this study was to develop a reference statistical model predictive of the average cost per patient of the lower limb osteoarthritis in the general population (ESCAPEg) and in those treated by spa therapy (ESCAPEc), and to identify the main determinants.

### **Inclusion criteria**

Patient with tibiofemoral osteoarthritis and / or patellofemoral uni and / or bilateral, and / or hip osteoarthritis and uni / or bilateral; aged more than 20 years; Patient affiliated personally to the Caisse Nationale de l'Assurance Maladie des Travailleurs Salariés (CNAM-TS) for at least six months; Patient giving written informed consent for participation in the study and to extract the patient data from the ERASME database of the CNAM-TS.

## **- COHORT OF SWIMMERS: HEALTH EFFECTS OF EXPOSURE TO WATER DISINFECTION BY-PRODUCTS**



Last update : 09/05/2017

Head :

Nadif Rachel, Épidémiologie, santé, travail

Responsible organization :

Institut National de la Santé et de la Recherche Médical

## **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

To investigate the health effects on exposure to water disinfection by-products from swimming pool water on experienced swimmers.

### **Inclusion criteria**

- Male;
- Between 15 and 28 years old;
- Having daily swimming training sessions.

## **SPARCLE - LONGITUDINAL STUDY ON CHILDREN WITH CEREBRAL PALSY LIVING IN EUROPE**

Last update : 01/01/2020

Head :

Colver Allan, The Institute of Health and Society

Responsible organization :

European Commission Research

Wellcome Trust

## **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

The study aims to identify which environmental factors, if improved, will yield the greatest benefits for children with disabilities and their families. This knowledge will inform EU policy in the health, educational and social sectors and generate protocols to optimise outcomes

### **Inclusion criteria**

Children with cerebral palsy (local registers) aged 8-12 years at the beginning of the study

## **B030 - OBSERVATIONAL STUDY: CONDITIONS OF USE OF DULOXETINE IN FRANCE**

Last update : 01/01/2020

Head :  
Laboratoire , Eli Lilly France

Responsible organization :  
Eli Lilly

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Primary objective: evaluate the conditions for use of duloxetine in routine practice;  
Secondary objectives: characteristics of the patients treated by duloxetine, profiles of duloxetine prescribers, characteristics of treatment.

### **Inclusion criteria**

Patient receiving duloxetine in a pharmacy regardless of the indication, whether treatment initiation or renewal

## **ALCOROUTE - ENQUÊTE SUR LES CONTREVENANTS À L'ALCOOLÉMIÉ ROUTIÈRE**

Last update : 09/05/2017

Head :  
Craplet Michel

Responsible organization :  
Association Nationale de Prévention en Alcoologie et Addictologie (ANPAA)

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

The purpose of the Alcoroute survey was to analyze what is currently going on in the ANPAA care structures for drinking drivers, i.e.: what is their typology? What is their future in the care structures? What

consumption modifications are observed in terms of treatment

### **Inclusion criteria**

The subjects studied were those consulting for "driving while intoxicated" in participating Alchology ambulatory cure centers (CCAA) and who were consulting for the first time either spontaneously or sent by justice.

## **ANRS CO2 SEROCO/HEMOCO - MULTICENTRIC COHORT OF HIV POSITIVE PATIENTS**

Last update : 01/01/2019

Head :

Meyer Laurence, U822

Responsible organization :

Agence Nationale de Recherches sur le Sida et les hépatites virales (ANRS)

### **Type of database**

Study databases

### **Study databases (details)**

Cohort study

### **Main objective**

General objective: to investigate spontaneous HIV disease progression and the treatment response of haemophilic (HEMOCO) or non-haemophilic (SEROCO) subjects. The prognostic role of markers, in particular CD4, HIV-RNA and HIV-DNA.

### **Inclusion criteria**

SEROCO: Seropositive HIV-1 patients (ELISA test confirmed by Western blot) with known date of contamination or whose HIV status is less than a year old (period between first positive sample date and the date the patient is enrolled in survey) and who do not have AIDS at baseline; attending physicians from centres participating in the survey; who are 18 years old or over and have given their informed consent.

HEMOCO: HIV-seropositive haemophiliacs (ELISA test confirmed by Western blot); attending physicians from centres participating in the survey, who are 3 years old and over and have given their informed consent. Consent is signed by the parties with parental authority of dependent minors or by the legal guardians of minors under tutelage.

## **EMBLEM - EUROPEAN MANIA IN BIPOLAR LONGITUDINAL EVALUATION OF MEDICATION**

Last update : 09/05/2017

Head :

Bellivier Franck, U1144 Variabilité de réponse aux psychotropes Équipe/activité : Biomarqueurs de la rechute et de la réponse aux traitements dans les addictions et les troubles de l'humeur

Responsible organization :  
Institut national de la santé et de la recherche médicale - Inserm

## **Type of database**

Study databases

## **Study databases (details)**

Longitudinal study (except cohorts)

## **Main objective**

To assess and monitor subjects presenting with a manic or mixed episode of bipolar disorder.

## **Inclusion criteria**

- male and female
- legal age
- presenting with bipolar disorder

# **QUALIMORPH - EMBRYO QUALITY EVALUATION IN INTRA CYTOPLASMIC SPERM INJECTION (ICSI) CYCLES WITH FERTIMORPH SYSTEM**

Last update : 01/01/2020

Head :  
Niez Philippe  
Arbo Elisangela

Responsible organization :  
Ferring SAS

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

Evaluate the concordance between the embryo quality appreciated by the biologist according to his normal technique and according to the Fertimorph system

## **Inclusion criteria**

Embryos coming from couples having the following characteristics:

- Criteria linked to the woman: woman with normal response, all infertility etiologies, aged up to 36 years, maximum of one prior assisted reproduction technique (ART) cycle, current ART cycle with ICSI with ejaculated or frozen sperm, normal ovarian reserve (evaluated via hormone assays and where applicable by an antral follicle count), with a normal uterus and without hydrosalpinx.
- Criteria linked to the spouse: diagnosed male infertility, where applicable normal male karyotype,

-Criteria linked to the studied attempt: at least 5 mature (MII) oocytes retrieved, maximum of two embryos transferred.

## TRACY - FRENCH COHORT OF URANIUM CYCLE WORKERS

Last update : 01/01/2020

Head :

Samson Eric, Laboratoire d'épidémiologie des rayonnements ionisants (LEPID)

Responsible organization :

IRSN (Institut de radioprotection et de sûreté nucléaire)

### Type of database

Study databases

### Study databases (details)

Cohort study

### Main objective

To study mortality risks for workers exposed to uranium.

### Inclusion criteria

- Men or women - statutory - employed for at least 6 months

## ANRS CO8 APROCO-COPILOTE - COHORT OF HIV-INFECTED PATIENTS WHO BEGAN TREATMENT WITH PROTEASE INHIBITOR IN 1997-1999. FOLLOW-UP OVER 4 YEARS ON THE LONG-TERM EFFECTS LINKED WITH ADHERENCE AND TOLERANCE (COPILOT).

Last update : 01/01/2019

Head :

Leport Catherine, LABORATOIRE DE RECHERCHE EN PATHOLOGIE INFECTIEUSE

Responsible organization :

Agence Nationale de Recherches sur le Sida et les hépatites virales (ANRS)

### Type of database

Study databases

### Study databases (details)

Cohort study

### Main objective

General objective: To investigate determinants for long-term effects of adherence and tolerance to highly active antiretroviral drugs, in relation to patients infected with HIV-1 who were treated for the first time

under a protease inhibitor in 1997-99. Secondary objectives: - to determine the part that host genetic polymorphisms play in the progression of the infection and onset of serious adverse events - to investigate viral load and antiretroviral resistance in plasma cell compartments in relation to antiretroviral concentrations over time.

### **Inclusion criteria**

Patients included in the APROCO cohort (adults infected with HIV-1 at first prescription of protease inhibitor in 1997-99), monitored for four years in APROCO after initial enrolment, monitored by voluntary services in order to participate in extended follow-up, having given their written consent.

## **DECLIC - DESCRIPTION OF THE PROFILE OF HYPERTENSIVE PATIENTS TREATED WITH CALCIUM CHANNEL BLOKERS**

Last update : 01/01/2018

Head :

Millet véronique, Pfizer

Responsible organization :

Pfizer

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Describe the profile and the care for hypertensive patients treated with calcium channel blockers and their distribution according to the level of the cardio-vascular risk as defined by HAS (French National Authority for Health) 2005.

- Describe the distribution of the risk factors according to sub-groups of age.
- Describe the profile of hypertensive patients on calcium channel blockers according to blood pressure control.
- Describe the therapeutic strategy when the high blood pressure is not controlled.

### **Inclusion criteria**

Patient over the age of 18 years.

- Patient seen through routine liberal medical practice, regardless of the reason for consultation.
- Hypertensive patient whether or not balanced, treated with calcium channel blockers in monotherapy or in association.
- Patient informed of the objectives of the survey.

## **SMP - ÉTUDE TRANSVERSALE SUR LA SANTÉ MENTALE DES PERSONNES DÉTENUES : ENQUÊTE "LONGUES PEINES"**

Last update : 09/05/2017

Head :

Falissard Bruno, Inserm U1018 - Centre de Recherche en Épidémiologie et Santé des Populations Équipe:  
Genre, Santé sexuelle et Reproductive

Responsable organization :  
Direction générale de la Santé (DGS)  
Direction de l'Administration Pénitentiaire (DAP)

### **Type of database**

Bases de données issues d'enquêtes

### **Study databases (details)**

Etudes transversales non répétées (hors enquêtes cas-témoins)

### **Main objective**

Initial : étudier qualitativement l'évolution de détenus présentant des troubles psychiatriques avérés, et condamnés à de longues peines.

Modifié du fait du changement de méthodologie de l'étude : étudier, qualitativement, les conditions de détention, le vécu et la situation psychiatrique de personnes incarcérées pour de longues peines et ayant déjà vécu plus de 10 années de détention.

### **Inclusion criteria**

60 détenus condamnés à de longues peines et incarcérés sans discontinuité depuis au minimum 10 ans, dans 3 établissements sélectionnés

## **- STUDENT HEALTH SURVEY IN 2005**

Last update : 09/05/2017

Head :  
Fontaine Danièle  
Dion Nicolas

Responsable organization :  
Union nationale des Sociétés Etudiantes Mutualistes régionales (USEM)  
Fédération nationale des observatoires de santé (FNORS)

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

To study student health problems with a focus on mental health issues; to identify the relationship between socio-demographic characteristics and student perceptions of health and behaviour; student perception and use of health care system; to identify prevention requests in order to refine and develop regional student health-insurance plans and programmes.

## Inclusion criteria

Students associated with six regional student health-insurance companies: Mutuelle Étudiants de Provence (MEP), Mutuelle Générale des Étudiants de l'Est (MGEL), Société Mutualiste des Étudiants de Bretagne-Atlantique (SMEBA), Société Mutualiste des Étudiants du Nord-Ouest (SMENO) Société Mutualiste des Étudiants de la Région Parisienne (SMEREP), VITTAVI

## CADASIL - CADASIL PATIENT COHORT

Last update : 12/01/2017

Head :

Chabriat Hugues, CERVCO CENTRE DE RÉFÉRENCES POUR LES MALADIES RARES DES VAISSEaux DU CERVEAU ET DE L'OEIL, HOPITAL LARIBOISIÈRE

Responsible organization :

APHP

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

General objective: prospective study of patients with Notch 3 gene mutation thought to cause CADASIL: research into prognostic factors and useful markers for preventative treatment trial. Secondary objectives: - To determine clinical-radiological correlations - Modifying factors of clinical phenotype and MRI studies - Prognostic and progressive factors - Clinical phenotype variability - Natural history of disease

## Inclusion criteria

Adult patients with Notch 3 genetic mutation thought to cause CADASIL

## - ÉTUDE LONGITUDINALE SUR DES PATIENTS SOUFFRANT DE DÉPRESSION: CONCORDANCE ENTRE AUTO-ÉVALUATION ET ÉVALUATION DU CLINICIEN

Last update : 09/05/2017

Head :

Falissard Bruno, Inserm U1018 - Centre de Recherche en Épidémiologie et Santé des Populations Équipe: Genre, Santé sexuelle et Reproductive  
Corruble Emmanuelle, INSERM U669 CHU DE BICETRE - SERVICE DE PSYCHIATRIE

Responsible organization :

INSERM - Institut National de la Santé et de la Recherche Médicale

## Type of database

Bases de données issues d'enquêtes



## Study databases (details)

Etudes de cohortes

### Main objective

Evaluer les différences entre l'auto-évaluation et l'évaluation du clinicien dans la maladie dépressive

### Inclusion criteria

Soixante-quatre patients consécutivement nouvellement admis en hospitalisation dans le service psychiatrie de l'hôpital Paul Brousse (Villejuif) pour dépression (DSM-III-R) majeure nécessitant le début d'un traitement antidépresseur.

Critères d'exclusion: patients avec troubles bipolaires, troubles schizophréniques, abus d'alcool ou de drogues, troubles neurologiques.

## OCSIGEN - FOLLOW-UP COHORT OF ASTHMATIC PATIENT TREATED WITH INHALED CORTICOSTEROID

Last update : 01/01/2019

Head :

Pribil Céline, Laboratoire GSK

Responsible organization :

Laboratoire GSK

### Type of database

Study databases

### Study databases (details)

Longitudinal study (except cohorts)

### Main objective

Describe the conditions of use of the fluticasone and other inhaled corticosteroids in a pragmatic situation in general practice and in specialized pneumology, evaluate the severity of the asthma of patients cared for and measure the adequacy of the care with the national recommendations

### Inclusion criteria

GP section:

The eligible population is defined by all of the asthmatic patients over the age of 15 years who have consulted at least twice in the previous year and in whom an inhaled corticotherapy is in progress (at least one prescription in the last six months). Patients defined as such who come to consult spontaneously for asthma during the period of inclusion will also be included. Refusal to participate and the association of the asthma with an OCPD are criteria for non-inclusion.

Pneumologist section:

Asthmatic persons cared for in liberal pneumology present in the practice of the investigators of the study and meeting the inclusion criteria

# DÉPASS - DEPRESSION ? ANXIETY AND DISABILITY IN A COHORT OF PATIENTS IN GENERAL MEDICINE PRACTICE

Last update : 01/01/2020

Head :  
Crochard Anne

Responsible organization :  
Lundbeck SAS

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

To evaluate the change in self-reported personal, professional and social disability in patients with anxious or mood disorders three months after initiating antidepressant treatment.

## Inclusion criteria

Patient of at least 18 years of age

Patient having at least one of the 5 following diagnoses:

major depressive episode (i.e. characterized)

Generalized anxiety disorder

Social anxiety disorder

Panic disorder with or without agoraphobia

Obsessive compulsive disorders

Patient initiating a new treatment with an antidepressant

Patient being able to communicate and to evaluate his quality of life

Patient having received written information on the use of the medical data concerning himself within the framework of the study.

# FRAG'OS - ESTIMATION VIA THE WHO INDEX (FRAX®) OF THE RISK OF FRACTURE BY BONE FRAGILITY IN FRENCH WOMEN

Last update : 01/01/2018

Head :  
VIGNAL Franck, Sanofi Aventis

Responsible organization :  
Sanofi-aventis France

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Estimation via the WHO index (FRAX®) of the risk of fracture by bone fragility in French women aged 50 years and older, with osteoporosis

### **Inclusion criteria**

An antecedent of osteoporotic fracture, i.e. of a fracture occurring after the age of 45 years, without major trauma; A T-score at at least one of the two sites, lumbar or femoral,  $\geq -2.5$  SD  
; Treated or not

## **- BAROMETER ON LIVING CONDITIONS, HEALTH BEHAVIOUR AND MENTAL HEALTH OF STUDENTS IN PROVENCE-ALPES-CÔTE D'AZUR (PACA) REGION**

Last update : 09/05/2017

Head :

Verger Pierre , ORS PACA - UMR912-SESSTIM

Responsible organization :

Observatoire régional de santé Provence Alpes Côte d'Azur (ORS PACA), UMR 912 SESSTIM (Sciences Economiques et Sociales de la Santé et Traitement de l'Information Médicale)

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To investigate the prevalence of and factors that determine health behaviour in students. Specific objectives: to investigate the prevalence of and factors associated with psychological distress and certain psychiatric disorders (mood disorders, anxiety disorders and disorders related to use of psychoactive substances)

### **Inclusion criteria**

Enrolled in first year for the first time in one of six universities in the PACA region; Aged between 18 and 24 years old; Students enrolled in distance education and preparatory classes for grandes écoles were excluded

## **HIBOU - LONGITUDINAL STUDY ON THE IMPACT ON PARENTS OF THEIR INFANT'S BRONCHIOLITIS HOSPITALIZATION**

Last update : 01/01/2019

Head :

Lapillonne Alexandre

Responsible organization :  
Abbott France

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

Create and validate the Impact of Bronchiolitis Hospitalization Questionnaire (IBHQ) intended to specifically evaluate the impact on parents of their infants being hospitalized for bronchiolitis.  
Describe the impact on parents of their infants being hospitalized for bronchiolitis.

### **Inclusion criteria**

infants under 1 year old, hospitalized solely for bronchiolitis

## **HCC - HEPATOCELLULAR CARCINOMA**

Last update : 09/05/2017

Head :  
Stücker Isabelle

Responsible organization :  
Institut National de la Santé et de la Recherche Médicale

### **Type of database**

Study databases

### **Study databases (details)**

Case control study

### **Main objective**

To evaluate the role of UDP-glucuronosyltransferase enzymes in hepatocellular carcinoma.

### **Inclusion criteria**

- Men - adult - below 75 years of age - born in Europe

## **- SURVEY ON THE HEALTH OF CHILDREN ATTENDING PRIMARY SCHOOLS IN THE PROVENCE-ALPES-CÔTE D'AZUR (PACA) REGION**

Last update : 09/05/2017

Head :

Kovess-Masfety Viviane, INSERM U705 et U669

Responsible organization :  
Université Paris Descartes  
Inserm  
EHESP

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

1) To assess the prevalence of most the common mental health issues (with teachers and children) and physical issues (with parents) for primary school children. 2) to assess the principle social and psychological risk factors for mental health among young people (with parents). 3) To assess the use of health services (with parents).

## **Inclusion criteria**

Children attending private or public primary schools in Aix-Marseille and Nice.

# **D.E.S.I.R. - EPIDEMIOLOGICAL DATA ABOUT INSULIN RESISTANCE SYNDROME**

Last update : 01/01/2020

Head :

Balkau Beverley, U1018 Centre de Recherche en Epidémiologie et santé des Populations (CESP)

Responsible organization :

INSTITUT NATIONAL DE LA SANTE ET DE LA RECHERCHE MEDICALE - INSERM

## **Type of database**

Study databases

## **Study databases (details)**

Longitudinal study (except cohorts)

## **Main objective**

Describe natural history of insulin resistance syndrome and its consequences.  
Evaluate diabetes risk factors.

## **Inclusion criteria**

Men and women, aged between 30 and 65, consulting in the examination centers

# **MÉMOTS - ÉVALUATION DES FONCTIONS COGNITIVES DES**

# ENFANTS AYANT UN DÉFICIT EN HORMONE DE CROISSANCE ET DES ENFANTS NÉS PETITS POUR L'ÂGE GESTATIONNEL

Last update : 01/01/2020

Head :

Troy Sylvie, Pfizer

Responsible organization :

Pfizer

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Evaluate the level of verbal and non-verbal cognitive functions of children with a growth hormone deficiency (GHD) and children born small for the fertilization age before beginning a GH treatment.

- Evaluate the percentage of GHD children and the percentage of SGA children with language cognitive function disorders.
- Evaluate the percentage of GHD children and the percentage of SGA children with non-verbal cognitive function disorders.
- Compare the average level of the language and non-verbal cognitive functions of GHD and SGA children to a reference population of French children.
- Compare the level of verbal and non-verbal cognitive functions of GHD children and of SGA children.

## Inclusion criteria

Child between the ages of 4 to 9 years,

- Understanding French,
- For whom the GH treatment was approved by a health insurance fund,
- With an adequate supplementation of the other pituitary deficiencies in the case of a multiple deficiency.

# TARTARE - PRISON VIOLENCE PREVENTION

Last update : 10/01/2019

Head :

LAGARDE Emmanuel, U897 PRÉVENTION ET PRISE EN CHARGE DES TRAUMATISMES ( Equipe de l'ISPED - Institut de Santé Publique, d'Epidémiologie et de Développement)

Responsible organization :

INSERM - Institut National de la Santé et de la Recherche Médicale

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Identify individual and environmental factors associated to violence in prison  
Compile the results and propose new prevention strategies

### **Inclusion criteria**

All male prisoners of the prison of Bordeaux

## **EIPPAIN - LONG-TERM NEUROLOGICAL EFFECTS OF PAIN AND ANALGESIA AND SEDATION TREATMENT IN NEONATES**

Last update : 09/05/2017

Head :

Carbajal Ricardo, INSERM UNITE 149. RECHERCHES EPIDEMIOLOGIQUES EN SANTE PERINATALE ET SANTE DES FEMMES

Responsible organization :

ASSOCIATION URGENCES PEDIATRIQUES TROUSSEAU (HOPITAL ARMAND TROUSSEAU, PARIS)

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

To investigate the long-term neurological effects of pain as well as analgesia and sedation treatment administered to neonates

### **Inclusion criteria**

Children who were hospitalised in intensive care during neonatal period

## **INTENSIVE - INTENSIFICATION OF INSULIN THERAPY USED IN TYPE-2 DIABETES PATIENTS TREATED BY THE ASSOCIATION OF ORAL DIABETES MEDICATION AND BASAL INSULIN**

Last update : 01/01/2019

Head :

Vignal Franck, Sanofi Aventis

Responsible organization :

Sanofi-aventis France

### **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Describe the evolution of HbA1c at 6 months under different therapeutic strategies for the intensification of insulin therapy used in type-2 diabetes patients treated by the association of oral diabetes medication and basal insulin requiring the adding of rapid insulin

### **Inclusion criteria**

man or woman, over the age of 18 years, having type-2 diabetes meeting the criteria of the HAS: treated by the association of oral diabetes medication and basal insulin (semi slow or slow-acting) and requiring the adding of rapid insulin.

## **EGIR-RISC - RELATIONSHIP BETWEEN INSULIN SENSITIVITY AND CARDIOVASCULAR DISEASE**

Last update : 09/05/2017

Head :

Balkau Beverley

Ferrannini Ele

Responsible organization :

Centre de Recherche en Nutrition Humaine de Lyon (CRNHL)

### **Type of database**

Study databases

## **Study databases (details)**

Cohort study

### **Main objective**

Main objective: study of insulin resistance and atherosclerosis, determinants of insulin resistance, its evolution on 9 years and consequences, diabetes, cardiovascular diseases and premature death.

Secondary objectives:

1. The evaluation of insulin resistance and its association with cardiovascular diseases
2. Identification of genetic and environmental factors linked to insulin resistance
3. The proposition of a model to identify insulin resistance

### **Inclusion criteria**

- Aged between 30 and 60 years
- Good general health condition
- No symptoms of cardiovascular diseases
- blood pressure inferior to 170/90 mm Hg
- Total cholesterol inferior to 7,8 mmol/l
- Triglycerides levels inferior to 4,6 mmol/l



- Plasma glucose level (on an empty stomach and 2 hours after an oral glucose control) inferior to 7,0/11,1 mmol/l
- No treatment related to diabetes, hypertension or lipid disorders

## **PRESAGE - OBSERVATIONNAL STUDY ON POSTMENOPAUSAL WOMAN**

Last update : 01/01/2018

Head :

Hébuterne Xavier, Eli Lilly France

Responsible organization :

Eli Lilly

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Primary objective: estimate the percentage of osteoporotic patients among postmenopausal patients with a mineral bone density measurement for diagnosis purposes and who can benefit from therapeutic recommendations according to French recommendations.

Secondary objectives: factors associated with osteoporosis, characteristics and care for patients

### **Inclusion criteria**

Postmenopausal woman,

- Patient not treated for osteoporosis for at least one year (HRT, raloxifene, bisphosphonate, teriparatide, strontium ranelate),
- Patient coming for a bone densitometry for diagnosis purposes,
- Patient able to benefit from a therapeutic recommendation for the treatment of osteoporosis (osteopenic or osteoporotic patient with risk factors of fractures).

## **EPISURVEY HEPATITE B - A CROSS-SECTIONAL, MULTICENTRE, EPIDEMIOLOGICAL STUDY OF THE DRUG MANAGEMENT IN HEPATITIS B**

Last update : 09/05/2017

Head :

Bregman Bruno, Bristol-Myers Squibb

Schmidely Nathalie, Bristol-Myers Squibb

Responsible organization :

Bristol-Myers Squibb France (BMS)

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Describe the drug management of patients having chronic hepatitis B, in France, in 2006.

### **Inclusion criteria**

M/F  $\geq$ 18 years, infected by the hepatitis B virus for at least 6 months, seen in outpatient clinic visit at expert centers for caring of hepatitis (Hepatitis Referenced University Hospitals, General Hospital Centres, community-based hepato-gastroenterologists) between 1 Jan. and 31 March 2006.

## **EMPATHIE - ENFANTS - PARENTS - TROUBLE DÉFICIT DE L'ATTENTION AVEC HYPERACTIVITÉ : IMPACT ET EVALUATION**

Last update : 12/01/2018

Head :

Ponthieux Anne

Responsible organization :

Novartis Pharma S.A.S.

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Study of the impact of the Attention-Deficit Hyperactivity Disorder (ADHD) on the parents/tutors of afflicted school children.

The study also strives to evaluate the impact of the ADHD on the siblings of the children afflicted and likewise of the impact of the disease on the school environment, thanks to the evidence provided by parents/tutors.

### **Inclusion criteria**

Inclusion criteria for parents/tutors:

- Parents (father, mother or tutors) of a child aged 6 to 18 years suffering from ADHD
- Parents (father, mother, or tutors) accompanying their ADHD children for a routine consultation during the observation period,
- Parent (father, mother or tutor) of an child whether or not treated for his/her ADHD.

Inclusion criteria for siblings:

- Brother or sister without ADHD of a child corresponding to the aforementioned criteria,
- Brother or sister at least 9 years old and living with the ADHD child.

Non-inclusion criteria :

- ADHD child with important co-morbidity that could influence assessment of ADHD impact on psycho-socio-economic value,
- Patients sibling / parents with important co-morbidity that could influence the evaluation of their experience (i.e. at the level of the socio-psycho-economic impact),
- Parents unable to complete a self-questionnaire,
- Parents who participated to the feasibility study (interviews)

## **EUROPOP - EUROPEAN PROGRAMME ON THE RELATIONSHIP BETWEEN OCCUPATIONAL RISKS AND PREMATURE BIRTHS**

Last update : 09/05/2017

Head :

Ancel Pierre-Yves, U953 - Unité de recherche épidémiologique en santé périnatale et santé des femmes et des enfants

Responsible organization :

Institut national de la santé et de la recherche médicale - Inserm

### **Type of database**

Study databases

### **Study databases (details)**

Case control study

### **Main objective**

Consider the effect of particular work conditions on premature birth.

### **Inclusion criteria**

woman - adult - childbirth 22-36 weeks amenorrhea

## **FIBROSTIC - ETUDE TRANSVERSALE SUR DES PATIENTS PRÉSENTANT UNE INDICATION DE BIOPSIE POUR HÉPATITE B OU C : DIAGNOSTIC ET MÉDICO-ÉCONOMIE DES PROCÉDÉS NON INVASIFS DE LA FIBROSE HÉPATIQUE**

Last update : 01/01/2019

Head :

Durand-Zaleski Isabelle, Directrice de l'Unité de recherche clinique en économie de la santé d'Ile-De-France Responsable du service de Santé Publique (Groupe hospitalier Albert Chenevier- Henri Mondor)

Chevreur Karine, Directrice adjointe de l'Unité de recherche clinique en économie de la santé d'Ile-De-France Médecin de Santé Publique, Economiste de la santé

Responsible organization :

Assistance Publique - Hôpitaux de Paris

### **Type of database**

Bases de données issues d'enquêtes

### **Study databases (details)**

Etudes transversales non répétées (hors enquêtes cas-témoins)

### **Main objective**

Évaluer les performances du Fibroscan dans le diagnostic de la fibrose et de la cirrhose hépatiques comparativement à la ponction-biopsie hépatique.

### **Inclusion criteria**

- homme ou femme
- adulte
- présentant une indication de biopsie pour hépatite B ou C

## **IAD - SURVEY ON HOME CARE ASSISTANCE PROVIDERS FOR VULNERABLE PEOPLE (IAD)**

Last update : 07/01/2018

Head :

Marquier Rémy, -----

Responsible organization :

La Direction de la recherche, des études, de l'évaluation et des statistiques (DREES)

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To identify the socio-demographic profile and to explore career paths and training for provider To identify their role and responsibilities and the precise nature of their activities, in particular their employer situation (level of independence, isolation) and the context in which they provide care (service provider, employed by private individuals with or without company representation) To collect feedback from providers concerning their occupation, their working conditions and the difficulties they may encounter while working To provide elements that will help to define the actions undertaken in order to encourage home care for dependent elderly people and disabled people, to structure services that offer home care assistance, to promote the professional development of providers and the attractiveness of this activity sector.

### **Inclusion criteria**

Home care providers for vulnerable people (elderly people, disabled people or those requiring help in carrying out day-to-day activities)

## **EPURE - NATIONAL SURVEY ONE DAY ON SMOKING PREVALENCE IN GENERAL PRACTICE**

Last update : 01/01/2019

Head :

Millet Véronique, Pfizer

Responsible organization :

Pfizer

## **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Contribute, in the framework of an epidemiological cross-disciplinary approach, in determining the frequency of smoking whether or not diagnosed, whether or not treated, in patients consulting through general practice in France and accepting to participate in this survey.

- Describe the smoking behavior in the patient sub-groups according to the possible existence of associated pathologies.
- Study the professional practices concerning caring for smoking in general practice in France.

### **Inclusion criteria**

Age: 18 years or older

- Smoking patient consulting on the given day, with a general practitioner who has accepted to participate in the survey
- Patient who has received information concerning the survey and who has given his oral agreement for the use of anonymous data concerning him.

## **CHIMERE - MOTHER-TO-CHILD CHIKUNGUNYA VIRUS INFECTION: AN AMBISPECTIVE COHORT STUDY**

Last update : 10/01/2018

Head :

Gérardin Patrick, Centre d'Investigation Clinique (CIC 1410) - ex Centre d'Investigation Clinique - Epidémiologie Clinique (CIE2)

Fritel Xavier, Service de Gynécologie-Obstétrique et Médecine de la Reproduction, CHU de Poitiers, Université de Poitiers, Faculté de Médecine et Pharmacie. <http://medphar.univ-poitiers.fr/CESP> UMR Inserm U1018, Equipe 7 : Genre, Santé Sexuelle et Reproductive.

Responsible organization :

CHU de la Réunion

## **Type of database**

Study databases

### **Study databases (details)**

Cohort study

## Main objective

1- Primary objective: to assess the prevalence and the neurodevelopmental outcomes of maternal-fetal chikungunya infection ; 2-Secondary objectives: to decipher the mechanisms and risk factors of mother-to-child chikungunya virus transmission; to describe the chikungunya virus in the pregnant woman and in the neonate;

to propose a guideline for diagnosis and treatment of infections over pregnancy, childbirth and neonatal periods; to valid the virological diagnosis in the neonate using cord blood RT-PCRs; to study viral strains; to describe in children exposed in utero to maternal infection the kinetics of seroreversion of maternal transplacental IgG antibodies.

## Inclusion criteria

Maternal study: pregnant women with life insurance cover; Child study: children born from women enrolled in t the maternal cohort. Inclusion whatever the pregnancy story and the exposure the first two months, then only the exposed women reporting a chikungunya attack the next six following months.

# FUTURE - ETUDE TRANSVERSALE SUR DES PATIENTS SUIVIS PAR DES CARDIOLOGUES LIBÉRAUX POUR INSUFFISANCE CARDIAQUE SELON LA FRACTION D'ÉJECTION VENTRICULAIRE GAUCHE : TRAITEMENT MÉDICAL APRÈS HOSPITALISATION

Last update : 08/01/2018

Head :

Cohen Solal Alain, UMRS 942Université Paris 7 Denis-DiderotService de cardiologie, hôpital Lariboisière

Responsible organization :

Assistance Publique - Hôpitaux de Paris

## Type of database

Bases de données issues d'enquêtes

## Study databases (details)

Etudes transversales non répétées (hors enquêtes cas-témoins)

## Main objective

Déterminer la façon dont les cardiologues prennent en charge le traitement médical des patients en fonction de leur fraction d'éjection du ventricule gauche après la sortie de l'hôpital.

## Inclusion criteria

- homme ou femme
- adulte
- patient insuffisant cardiaque ambulatoire
- suivi par des cardiologues libéraux entre septembre 2007 et Août 2008
- hospitalisé au cours des 18 mois précédents, avec un diagnostic d'insuffisance cardiaque

# DOREMI - FACTORS INFLUENCING THE MOTIVATION OF THE

# UNBALANCED TYPE-2 DIABETIC OBESE PATIENT IN MONOTHERAPY BY ORAL DIABETES MEDICATION

Last update : 01/01/2018

Head :

VIGNAL Franck, Sanofi Aventis

Responsible organization :

Sanofi-aventis France

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Describe the factors influencing the motivation of the unbalanced type-2 diabetic obese patient in monotherapy by oral diabetes medication in adhering to the drug and hygiene and dietary care of its cardiometabolic risk factors.

## Inclusion criteria

Man or woman over the age of 18 years, Type-2 diabetes treated via oral diabetes medication monotherapy, Having a HbA1c rate  $\leq$  6.5%, A BMI  $\leq$  30 kg/m<sup>2</sup>, An abdominal obesity defined according to the NCEP/ATPIII criteria: Waist  $>$  88 cm for women and  $>$  102 cm for men,

Able to understand simple recommendations in terms of physical activity and complete a self-evaluation questionnaire

# JEUNES SEINE-SAINT-DENIS - YOUNG PEOPLE 13-25 YEARS OLD IN SEINE-SAINT-DENIS

Last update : 10/01/2019

Head :

Joubert Michel, CULTURE ET SOCIETE URBAINE (CSU)

Responsible organization :

CNRS - Centre national de la recherche scientifique

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

General objective: To understand the engagement process of risk behaviours (moderation to amplification)

and active factors in different stages of life for adolescents and young adults Secondary objective: To refine prevention policies regarding the possibility of reducing the most problematic engagements and associated risks.

### **Inclusion criteria**

Young people aged 13-25 years old in Seine-Saint-Denis

## **EPHA - OBSERVATIONAL STUDY ON THE DISTRIBUTION OF THE VARIOUS TYPES OF ATRIAL FIBRILLATION (AF)**

Last update : 01/01/2019

Head :

Vignal Franck, Sanofi Aventis

Responsible organization :

Sanofi-aventis France

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Describe the distribution of the various types of Atrial Fibrillation (AF) in patients with an antecedent or in AF followed by a cardiologist in metropolitan France.

### **Inclusion criteria**

? Patients seen in consultation by a cardiologist in general practice or at the hospital and followed by this cardiologist for at least 1 year

? Aged 18 years or older

? Having an antecedent or a documented AF (by ECG or Holter ECG) for at least one year, regardless of the rhythm on the day of inclusion

? Accepting to participate in the study and who have signed the consent agreement

## **- CANCER OF THE OESOPHAGUS ASSOCIATED WITH TOBACCO TYPE, ALCOHOL USE AND SPECIFIC DIET**

Last update : 09/05/2017

Head :

Launoy Guy, Cancers et populations

Responsible organization :

Institut National de Santé et Recherche Médicale

CHU Caen

### **Type of database**



Study databases

### **Study databases (details)**

Case control study

#### **Main objective**

To demonstrate that the risk of cancer of the oesophagus increases according to types of tobacco and alcohol use. To classify different types of tobacco and alcohol in accordance to risk of cancer of the oesophagus. To demonstrate certain dietary habits that increase the risk of cancer of the oesophagus.

#### **Inclusion criteria**

- Men - adult - under 85 years old - already presenting squamous carcinoma of the oesophagus

## **EMES - PRAGMATIC PHARMACOECONOMIC ASSESSMENT OF SINOVIAL® IN THE TREATMENT OF GONARTHROSIS**

Last update : 09/05/2017

Head :

Moore Nicholas, Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

Responsible organization :

Service de Pharmacologie, CIC-P 0005-INSERM U657- Université Bordeaux Segalen

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

#### **Main objective**

The objective was to compare the cost of two therapeutic strategies: Sinovial® and other care versus other care alone.

#### **Inclusion criteria**

Patient agreed to participate in this study and having signed the consent form; Patient having had a first injection of Sinovial® between 01/01/2003 and 31/03/2003; Patient having never previously received viscosupplementation treatment; Patient aged 20 years and over; Patient affiliated to the general health insurance system for at least 12 months at the time of first injection

## **ARCHIMED VILLE - PREVENTION OF VENOUS THROMBOEMBOLIC EVENTS WITH ARIXTRA 2.5MG FOR MEDICAL ILL PATIENTS IN GENERAL PRACTICE**

Last update : 01/01/2020

Head :

Leclerc-Zwirn Christel, Laboratoire GSK

Responsible organization :  
LABORATOIRE GSK

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

Evaluate the average duration of treatment via ARIXTRA® 2.5 mg with thromboprophylaxis in ambulatory general practice

### **Inclusion criteria**

Patient of at least 18 years of age,  
-bedridden or with reduced mobility,  
-for whom a treatment via ARIXTRA® 2.5 mg is initiated with thromboprophylaxis.

## **- 2008 HEALTH AND NUTRITION BAROMETER**

Last update : 01/01/2018

Head :

Escalon Hélène, Département Enquêtes et Analyses Statistiques

Beck François, Responsable du département Enquêtes et Analyses Statistiques

Responsible organization :

National Institute for prevention and health education

### **Type of database**

Study databases

### **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

### **Main objective**

Evaluate diet and physical activity behaviors, attitudes and knowledge of french people

### **Inclusion criteria**

Querying people in households:

-equipped with a landline telephone (white, orange or red list)  
-equipped solely with a mobile telephone  
- and fully unbundled

## **AMI - ETUDE LONGITUDINALE SUR DES PATIENTS ATTEINTS**

# D'INFARCTUS DU MYOCARDE: INTÉRÊT SUR LE LONG TERME DU POST-CONDITIONNEMENT

Last update : 09/05/2017

Head :

Ovize Michel, Laboratoire de Recherche en Cardiovasculaire, Métabolisme, diabétologie et Nutrition

Responsible organization :

Hospices Civils de Lyon

## Type of database

Bases de données issues d'enquêtes

## Study databases (details)

Etudes longitudinales (hors cohortes)

## Main objective

Déterminer l'action du post-conditionnement sur l'étendue de l'infarctus.

Mesurer l'amélioration de la fonction contractile du myocarde grâce au post-conditionnement.

## Inclusion criteria

- Homme ou femme
- Agé(e) de plus de 18 ans
- Présentant des douleurs thoraciques dans les 6 heures précédant un sus-décalage du segment ST supérieur à 1 mm sur 2 dérivations contiguës
- Patients nécessitant une intervention coronaire percutanée

# BINOME - OBSERVATIONAL STUDY OF THE SOCIODEMOGRAPHIC CHARACTERISTICS OF PATIENTS INFECTED WITH HIV B SUBTYPE VERSUS HIV NON-B SUBTYPE AND HAVING RECENTLY STARTED THEIR FIRST COURSE OF ANTIRETROVIRAL TREATMENT.

Last update : 07/01/2018

Head :

Cohen-Codar Isabelle

Responsible organization :

ABBOTT France

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Describe and compare the socio-demographic characteristics of patients infected with an HIV-1 of subtype B (group 1) compared to non-B (group 2) treated with a first anti-retroviral drug treatment.

Compare according to the HIV subtype the stage of the disease (clinical, viro-immunology, co-morbidities) at the time treatment is initiated, compare between the 2 groups (B and non-B) the conditions for taking the treatment and the perception by the patient of the therapy and of its effects.

Identification of any correlation between these latter parameters and the change in the disease between the initiation and the first line ARV and the inclusion in the cohort:

- change in the viral load (VL) according to the ARV treatment (PI, NNRTI) and of the HIV subtype (B or non-B), of the CD4,
- any occurrence of clinical events,
- any changes in treatments and reasons.

## Inclusion criteria

Age > 18 years, obtaining of written consent for the collection of this personal data, patient infected by the HIV-1 virus, patient coming to consult for the first or second routine follow-up visit (i.e. 1 to 4 months) after the initiation of the ARV treatment, genotype available when the treatment is initiated.

# AVANCE - OBSERVATIONAL STUDY OF THE REAL-WORLD USAGE, EFFICACY AND SAFETY OF ROSIGLITAZONE IN TYPE 2 DIABETICS

Last update : 01/01/2019

Head :

Leclerc-Zwirn Christel, Laboratoire GSK

Responsible organization :

Laboratoire GSK

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

To describe the profiles of patients being treated by general practitioners and specialists, compatibility of practice with the SPC, patient compliance, and the safety and efficacy of rosiglitazone in "real-world" conditions of use.

## Inclusion criteria

- Type 2 diabetic spontaneously consulting (i.e. independently of the study);
- Patient who is starting or has recently started (within a month) a course of treatment with Avandia® or Avandamet®;
- Patient who has given his/her consent to participate.

# - ETUDE TRANSVERSALE SUR LES JEUNES CALÉDONIENS ÂGÉS

# DE 16 À 25 ANS : SITUATION SOCIALE ET COMPORTEMENT DE SANTÉ

Last update : 09/05/2017

Head :

Hamelin Christine , Équipe Épidémiologie des Déterminants Professionnels et Sociaux de la Santé, INSERM CESP

Salomon Christine, Centre de recherches en épidémiologie et santé des populations

Responsible organization :

Inserm

## Type of database

Bases de données issues d'enquêtes

## Study databases (details)

Etudes transversales non répétées (hors enquêtes cas-témoins)

## Main objective

L'étude « Situation sociale et comportements de santé des jeunes en Nouvelle-Calédonie » vise à décrire les comportements de santé des jeunes, garçons et filles de 16 à 25 ans, en rapport avec leur situation familiale et sociale. Quatre grandes thématiques ont été explorées : les consommations de substances psychoactives, les mises en danger de soi et d'autrui sur la route, la santé sexuelle et reproductive et la santé mentale. Le présent rapport vise à décrire ces comportements de santé, à identifier les principales conduites à risque, à en évaluer l'importance et à examiner la manière dont ces conduites diffèrent ou non selon l'âge, le sexe la communauté et le lieu de résidence afin de cerner les groupes les plus exposés. L'analyse approfondie des données sur la santé mentale, est également proposée.

## Inclusion criteria

L'étude concerne les jeunes calédoniens âgés de 16 à 25 ans

# FOCUS / FRENCH OBSERVATIONAL COHORT OF USUAL SMOKERS - ETUDE TRANSVERSALE SUR LA PRÉVALENCE ET LE PROFIL DU TABAGISME EN FRANCE

Last update : 11/01/2018

Head :

Aubin Henri-Jean, U1178 Santé mentale et santé publique Equipe/activité : Addicologie

Responsible organization :

Assistance Publique - Hopitaux de Paris AP-HP

PFIZER

TNS Healthcare

CHR de Metz

Centre Régional de Lutte du Cancer Val d'Aurelle

## Type of database

Bases de données issues d'enquêtes

### **Study databases (details)**

Etudes transversales non répétées (hors enquêtes cas-témoins)

### **Main objective**

Mesurer la prévalence du tabagisme en France, identifier le profil type des fumeurs,

### **Inclusion criteria**

- hommes et femmes
- âgés de 15 ans et plus

## **ENDOCARDITIS - REPEATED CROSS-SECTIONAL STUDY FOR PATIENTS WITH INFECTIVE ENDOCARDITIS: SURGICAL IMPACT**

Last update : 01/01/2018

Head :

Alla Francois, CIC-EC INSERM CHU

Hoen Bruno, INFECTIOLOGIE CHU

Duval Xavier, CIC

Responsible organization :

CHU Nancy

### **Type of database**

Study databases

### **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

### **Main objective**

To assess the impact of surgical strategy on medium to long-term morbidity and mortality.

Secondary objective: descriptive and analytic.

### **Inclusion criteria**

Patients over 18 years of age treated in hospital for definite (Duke) infective endocarditis (IE) in 2008.

## **ISAAC - INTERNATIONAL STUDY OF ASTHMA AND ALLERGIES IN CHILDHOOD IN FRANCE**

Last update : 09/05/2017

Head :

Annesi-Maesano Isabella, UMRS 707 : Épidémiologie, systèmes d'information, modélisationÉpidémiologie des maladies allergiques et respiratoires EPAR

Responsible organization :

## Type of database

Study databases

### Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

### Main objective

To establish a global map of the prevalence of childhood allergic diseases and to determine the risk factors for these diseases by comparing contrasting population groups based on a particularly strong or weak prevalence.

The ISAAC study consists of three phases:

The objective of Phase I is to assess the prevalence of asthma, allergic rhinitis and eczema by using a simple questionnaire and possible video questionnaire (for asthma), in representative samples of children from two age groups: 13-14 years old (obligatory for all centres) and 6-7 years old (optional).

Phase II aims to identify the risk factors and treatment procedures for allergic diseases through a more detailed study of selected population groups with relatively high or low prevalence rates suggested during Phase I.

Phase III aims to assess the trends in allergy prevalence.

As such, the Phase I protocol will be repeated in areas that participated in this phase.

### Inclusion criteria

- primary school children
- between 8 and 13 years old

## SPIRALE - POST-OCCUPATIONAL SURVEILLANCE OF EXPOSED WORKERS

Last update : 05/11/2015

Head :

Carton Matthieu, INSERM U687 MIXTE : INSERM ET CNAMTS

Zins Marie, INSERM U687 MIXTE : INSERM, CNAMTS

Goldberg Marcel, INSERM U687 INSERM U687

Responsible organization :

INSERM - Institut National de Santé et Recherche Médicale

## Type of database

Study databases

### Study databases (details)

Cohort study

### Main objective

General objective: to identify retired workers that were exposed to carcinogens throughout their

professional life; to inform them of their right to post-occupational medical monitoring and to support them throughout their efforts.

Secondary objective: to conduct an epidemiological (cohort) study to describe the long-term health effects of the considered exposures, and to assess the benefits of post-occupational monitoring in terms of health and social awareness.

### **Inclusion criteria**

Occupational exposure to asbestos or wood dust;  
Insured under the general social security scheme.

## **- INTERVENTIONAL LONGITUDINAL STUDY ON THE EFFECTS OF LONG-CHAIN OMEGA-3 DOCOSAHEXAENOIC ACID ON PLATELET FUNCTION AND REDOX STATUS**

Last update : 09/05/2017

Head :

Vidal Hubert, U1060

Véricel Evelyne, U1060

Responsible organization :

INSERM - Institut National de la Santé et de la Recherche Médicale

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

To determine the effects of increased intake of docosahexaenoic acid on platelet function and oxidative stress in a group of healthy male volunteers.

### **Inclusion criteria**

Healthy male volunteers.

## **REAL.FR - LONGITUDINAL STUDY ON PATIENTS IN ALZHEIMER'S DISEASE NETWORK**

Last update : 09/05/2017

Head :

Vellas Bruno, Inserm U 1027

Responsible organization :

CHU de Toulouse

### **Type of database**



Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

#### **Main objective**

To ensure regular monitoring of the patient and those close to them in order to assess the cognitive and non-cognitive aspects of the disease.

To analyse treatment methods, as well as drug- and non-drug-related therapy for elderly people with AD across a national sample with diverse socio-cultural contexts. To identify potential treatment issues, such as repeated hospitalisation or emergency institutionalisation.

#### **Inclusion criteria**

- Male and female;
- Patient with Alzheimer's-type dementia.

## **ORSOSA - CROSS-SECTIONAL STUDY ON THE ORGANISATION OF HEALTHCARE AND THE HEALTH OF HEALTHCARE WORKERS**

Last update : 01/01/2018

Head :

Sobaszek Annie, Service de médecine du travail du personnel hospitalier, pôle de santé publique  
De Gaudemar Régis, Service de médecine et santé au travail

Responsible organization :  
CHU de Grenoble

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

#### **Main objective**

The study aims to construct and validate a methodological tool to assess psychological and organisational constraints (POC) in healthcare by adapting measurement tools, such as objective (observations) and subjective (self-administered questionnaires) tools.

The long-term objective is the transfer of skills and the developed methodological tool to smaller healthcare institutions that are also subject to the same regulatory risk assessment requirements.

#### **Inclusion criteria**

- Statutory and voluntary contracted staff;
- 50% of the subjects' work is as a nurse, nursing assistant or doctor.

Non-inclusion criteria

- Temporary staff and staff that are planning to leave the UHC in two years (retirement, moving, etc.).

# CAPITAL - CROSS-SECTIONAL STUDY ON PATIENTS WITH PROSTATE CANCER: HORMONAL THERAPY PRESCRIBED BY FRENCH UROLOGISTS IN EVERYDAY PRACTICE

Last update : 10/01/2017

Head :

Rouprêt Morgan, Services d'Urologie Hôpital Pitié-Salpêtrière, Groupe Hospitalo-Universitaire Est

Responsible organization :

Hôpital Pitié-Salpêtrière

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

To define professional practices for urologists and radiotherapists regarding the use of hormonal therapy in prostate cancer.

## Inclusion criteria

- male and female
- adult
- prostate cancer patient

# ELCAPA - LONGITUDINAL STUDY ON ELDERLY CANCER PATIENTS: IMPACT OF COMPREHENSIVE GERIATRIC ASSESSMENT

Last update : 01/01/2020

Head :

Paillaud Elena, Médecine gériatrique - Albert Chenevier

Responsible organization :

CHU Henri Mondor, AP-HP

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

The general aim of ELCAPA is to identify the impact of comprehensive geriatric assessment components

on changes in planned cancer treatment, and to investigate the frequency where cancer treatment plans were not fulfilled (with reasons for non-fulfilment), as well as associated risk factors.

### **Inclusion criteria**

Cancer patients aged 70 years and over that were referred for geriatric oncology consultation at Henri/Mondor/Paul Brousse and have undergone a comprehensive geriatric assessment (CGA).

## **OFICA - NATIONAL FRENCH OBSERVATORY OF ACUTE HEART FAILURE**

Last update : 03/01/2020

Head :

Logeart Damien, AP-HP, Groupe Hospitalier Lariboisiere Saint Louis, INSERM U942, Université Paris 7

Responsible organization :

Société Française de Cardiologie

Assistance Publique - Hôpitaux de Paris

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

To assess the characteristics of patients admitted for acute or decompensated heart failure, their treatment course and practices regarding their treatment until discharge, and to measure the impact of these data on prognosis at 3 months.

To obtain epidemiological data, evaluate practices in terms of diagnostic and therapeutic treatment and to provide answers to clinicians' questions.

### **Inclusion criteria**

- Male or female;
- Over 18 years old;
- In hospital on the day of enrolment, i.e., hospitalised on the same day or days before;
- Principal diagnosis or associated acute heart failure;
- Informed consent.

Exclusion criteria:

- Heart surgery;
- Scheduled hospitalisation;
- Diagnosis of acute heart failure reversed in favour of an alternative diagnosis before discharge from hospital;
- Patients that refused to participate.

## **PRELUD - FRONT-LINE SURVEY ON DRUG USERS**

Last update : 09/05/2017

Head :  
Cadet-Tairou Agnès

Responsible organization :  
OFDT - Observatoire Français des Drogues et des Toxicomanies

### **Type of database**

Study databases

### **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

### **Main objective**

This survey focuses on the knowledge and monitoring of psychoactive product users and their practices.

### **Inclusion criteria**

Users of facilities situated in clusters participating in the TREND system.

## **- CROSS-SECTIONAL STUDY ON THE PREVALENCE OF ALLERGIC RHINITIS IN ASTHMATIC PEOPLE**

Last update : 09/05/2017

Head :  
Magnan Antoine, UMR 915: l'Institut du thorax

Responsible organization :  
CHU Nantes  
Institut national de la santé et de la recherche médicale - Inserm

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To describe the frequency and severity of allergic rhinitis in French asthmatic patients.

### **Inclusion criteria**

- male and female
- adult
- asthmatic

## **EXPRESA - EXACERBATION PREDICTIVE FACTORS IN SEVERE**

# ASTHMA - COHORT ON PREDICTIVE EXACERBATION FACTORS IN SEVERE ASTHMATICS

Last update : 01/01/2020

Head :

Magnan Antoine, UMR 915: l'Institut du thorax

Responsible organization :

CHU Nantes

Institut national de la santé et de la recherche médicale - Inserm

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

To validate exacerbation prediction in certain lymphocyte populations by comparing markers already identified (NO, EEI), based on factors that do or do not cause infection.

## Inclusion criteria

- Male and female;
- Adult;
- Severe asthmatic.

# - LONGITUDINAL GENERAL POPULATION STUDY: PULMONARY NODULE DETECTION BY COMPUTED TOMOGRAPHY (CT) SCAN

Last update : 01/01/2019

Head :

Launoy Guy, Cancers et populations

Responsible organization :

Institut National de Santé et Recherche Médicale

CHU Caen

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

To prove that the presence of pulmonary nodules (by scan) increases the risk of developing lung cancer.  
To prove that there is no relationship between exposure to asbestos and the risk of developing lung

cancer.

### **Inclusion criteria**

- Male or female;
- Not-working or retired.

## **SAM - SURVEY ON DRUGS AND FATAL ROAD TRAFFIC ACCIDENTS**

Last update : 09/05/2017

Head :

Laumon Bernard, département TS2

Responsible organization :

INRETS- Institut National de Recherche sur les Transports et leur Sécurité

OFTD

LAB PSA Peugeot-Citroën/RENAULT

INSERM

CEESAR - Centre Européen d'Etudes de Sécurité et d'Analyse des Risques

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

The epidemiological analysis aims to assess the role played by drugs in fatal road accidents. This includes estimating the relative risk of fatal accidents that involve drugs and the prevalence of drugs among "circulating" (accident-free) drivers. Even though only drivers involved in a fatal accident are included, both of these parameters may be estimated, provided certain analysis conditions are met.

It also aims to characterise fatal accidents involving drivers under the influence and to compare these characteristics with those from other fatal accidents. The objective is to determine whether the first accident type bears specific traits that may be the basis for defining preventative initiatives that target drug users.

### **Inclusion criteria**

- Present at the accident with immediate life-threatening consequences;
- Consent.

## **- CROSS-SECTIONAL STUDY ON OPIATE ADDICTION TREATMENT IN GENERAL MEDICINE**

Last update : 01/01/2020

Head :

Lapeyre-Mestre Maryse

Responsible organization :  
Université de Toulouse

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To assess whether conducting urine-based toxicity screening tests in the general practitioner's surgery improves treatment for drug-substituted patients.

### **Inclusion criteria**

General practitioners exclusively practising in ambulatory care in the Midi-Pyrénées region.

## **EPIMART - CROSS-SECTIONAL STUDY ON EPILEPTIC PATIENTS LIVING IN MARTINIQUE**

Last update : 09/05/2017

Head :  
Jallon Pierre , Unité d'épileptologie et d'EEG

Responsible organization :  
CEA

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To determine the overall incidence of epileptic seizures within the island population and to identify clinical seizure type and factors.

### **Inclusion criteria**

- Male and female;
- Living in Martinique;
- Experienced an epileptic seizure between 01/05/1994 and 31/04/1995.

## **- CASE-CONTROL STUDY ON THE IMPACT OF BOTTOM-UP AND TOP-DOWN NEGATIVE EMOTION GENERATION ON MEMORY FORMATION AMONG HEALTHY AND TRAUMATISED ADOLESCENTS**

Last update : 01/01/2018

Head :

Guillery-Girard Bérengère, INSERM U1077

Responsible organization :

CHU Caen

Université de Caen

INSERM

## **Type of database**

Study databases

## **Study databases (details)**

Case control study

## **Main objective**

The aim of the study is to investigate the influence of bottom-up and top-down processes in the generation of emotions and their effect on attention and memory, and to assess the differential impact of attention deficits on memory through a case-control survey of traumatised adolescents with attentional bias.

## **Inclusion criteria**

- Individuals with a post-traumatic stress disorder with no stress for at least a year;
- Individuals with no other neurological, psychiatric or learning disorders;
- No intellectual disability;
- Not taking medication;
- Native level French;
- Signed consent form (parents and children).

# **- SCHOOL-RELATED INJURIES AS A FUNCTION OF GENDER**

Last update : 09/05/2017

Head :

Falissard Bruno, U 669

Chau Narkasen, U 669

Responsible organization :

INSERM - Institut National de la Santé et de la Recherche Médicale

## **Type of database**

Study databases

## **Study databases (details)**

Cohort study

## **Main objective**



To study the characteristics and risk factors of school-related injuries (accidents).

### **Inclusion criteria**

Pupils registered in institutions participating in the study.

## **AEPEI - REPEATED CROSS-SECTIONAL STUDY FROM THE ASSOCIATION FOR THE STUDY AND PREVENTION OF INFECTIVE ENDOCARDITIS**

Last update : 12/01/2018

Head :

Duval Xavier, CIC7 CIC BICHAT Recherche clinique : Cardio-vasculaire, pneumologie, diabétologie, maladies infectieusesU738 Modèles et méthodes de l'évaluation thérapeutique des maladies chroniquesÉquipe: Modélisation biostatistique et pharmacométrie

Responsible organization :

Assistance Publique des Hopitaux de Paris  
Institut National de la Santé et de la Recherche Médicale

### **Type of database**

Study databases

### **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

### **Main objective**

To describe trends in incidence, epidemiological, clinical, microbiological and prognostic factors, as well as surgical treatment procedures for infective endocarditis in France in 2008, in relation to 1991 and 1999.

### **Inclusion criteria**

- Male and female;
- Over 18 years old;
- Treated for infective endocarditis.

## **EPP1 - PROSPECTIVE PARISIAN SURVEY 1**

Last update : 09/05/2017

Head :

Ducimetière Pierre

Responsible organization :

Institut National de la Santé et de la Recherche Médicale

### **Type of database**

Study databases

## Study databases (details)

Cohort study

### Main objective

To determine the incidence and prevalence of ischaemic cardiomyopathy (IC) among the French population and to quantify the risk associated with various risk factors.

### Inclusion criteria

- Men;
- Born in France;
- Employed as policemen in the civil service of the City of Paris;
- Aged between 43 and 55 years old during the enrolment period;
- Free from cardiovascular disease.

## EPP2 - PROSPECTIVE PARISIAN SURVEY 2

Last update : 09/12/2017

Head :

Ducimetière Pierre, U780

Responsible organization :

Institut National de la Santé et de la Recherche Médicale

### Type of database

Study databases

## Study databases (details)

Cohort study

### Main objective

General objective: to investigate various causes of mortality, especially cardiovascular events.

Secondary objective: to investigate coronary morbidity over a period of 5 years.

### Inclusion criteria

Men born in France and employed as policemen in the civil service of the City of Paris or in the SNCF (French National Rail) and aged between 43 and 55 years old during the enrolment period.

## NUTRIMAY - FOOD, NUTRITION AND HEALTH IN MAYOTTE: 2006 NUTRIMAY STUDY

Last update : 09/05/2017

Head :

Deschamps Valerie, Unité de surveillance et d'épidémiologie nutritionnelle

Responsible organization :

INVS-Institut de Veille Sanitaire

## Type of database

Study databases

### Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

### Main objective

To describe food consumption, physical activity and nutrition status of the population living in Mayotte, with particular focus on children under 5 years old and women of childbearing age.

### Inclusion criteria

Individuals living in an ordinary household in Mayotte and capable of understanding the aims of the study.

## ENTRED 2007-2010 - NATIONAL REPRESENTATIVE SAMPLE OF PEOPLE WITH DIABETES 2007-2010

Last update : 07/01/2019

Head :

Fosse-Edorh Sandrine

Responsible organization :

INVS - Institut de Veille Sanitaire

## Type of database

Study databases

### Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

### Main objective

To describe the characteristics of people pharmacologically treated for diabetes, their health with regards to diabetes, the quality of care received, treatment plan and self-management education; their quality of life, experiences and requirements in terms of education and information, as well as the cost of diabetes.

### Inclusion criteria

Beneficiaries of the French general scheme of health insurance (CNAMTS) (excluding local mutualist sections) and the Independent Scheme for employees (RSI); residing in mainland France or overseas départements (DOM); and who have received at least three reimbursements for oral antidiabetic medications and/or insulin over the last twelve months.

## PREVAGAY - INCIDENCE OF HIV AMONG THE PARISIAN GAY COMMUNITY

Last update : 06/01/2019

Head :  
Velter Annie

Responsible organization :  
InVS

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

To determine the incidence of HIV in the Parisian gay community.

## **Inclusion criteria**

- male
- adult
- gay
- having had relations within 12 months preceding the study

# **SATURN-INF 2008-2009 SURVEY (ENQUÊTE SATURN-INF 2008-2009) - NATIONAL SURVEY ON LEAD POISONING PREVALENCE AND INFECTIOUS DISEASE SEROPREVALENCE IN CHILDREN BETWEEN 6 MONTHS AND 6 YEARS OLD (LEAD POISONING COMPONENT AND BIOBANK)**

Last update : 01/01/2019

Head :  
Invs-DSE

Responsible organization :  
Institut de Veille Sanitaire

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

- To estimate the national prevalence of lead poisoning in children between 6 months and 6 years old;
- To determine the blood lead level distribution of children between 6 months and 6 years old by administrative region;
- To update knowledge of factors that determine blood lead level;
- To validate indicators that characterise at-risk areas.

## **Inclusion criteria**

- Children between 6 months and 6 years old;
- Covered by social security health insurance;
- Inpatient or outpatient hospital stay during study period;
- Where a blood sample was prescribed as part of treatment or where a sampling system (catheter) is already in place;
- Residing in France at the time of enrolment.

# **ENNS - FRENCH NATIONAL NUTRITIONAL AND HEALTH STUDY**

Last update : 09/05/2017

Head :

Salanave Benoit, Equipe de surveillance et d'épidémiologie nutritionnelle (ESEN)

Responsible organization :

Agence nationale Santé publique France, ex Institut de veille sanitaire (InVS)

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

To describe food consumption, physical activity and nutrition of children and adults from a representative sample of the population living in France between 2006-2007.

## **Inclusion criteria**

Adults:

- Residing in metropolitan France during the study period (February 2006 to January 2007);
- Between 18 and 74 years old;
- Residing in an ordinary household with a landline or mobile phone;
- Living at least 5 days a week in the randomly selected household;
- Able to understand the primary implications of participating in the study by phone (hearing, understanding of French, no mental illness);
- Not suffering from a disease requiring enteral and parenteral artificial feeding;
- Agreed to participate in at least one part of the two-part study (food or clinical and biological assessment).

Children:

- Residing in metropolitan France during the study period (February 2006 to January 2007);
- Between 3 and 17 years old;
- Residing in an ordinary household with a landline or mobile phone;
- Living at least 5 days a week in a randomly selected household;
- Able to understand (child and child's guardian) the major implications of participating in the study by phone (hearing, understanding of French, no mental illness);
- Not suffering from a disease requiring enteral and parenteral artificial feeding;
- Agreed to participate in at least one part of the two-part study (food or clinical and biological

assessment).

## "AZF" SANTÉ COHORT - COHORT OF TOULOUSE CONURBATION WORKERS

Last update : 12/01/2019

Head :

Buisson Catherine, DÉPARTEMENT SANTÉ TRAVAIL (DST)

Diène Eloi, Département santé travail Institut de veille sanitaire 94415 Saint Maurice Cedex

Responsible organization :

Institut de veille sanitaire

### Type of database

Study databases

### Study databases (details)

### Main objective

General objectives:

- To investigate the medium-term health effects from the explosion at the AZF plant that occurred in September 2001 in the population of workers;
- To test access procedures for medical-administrative data (ERASMUS) for monitoring large prospective cohorts.

### Inclusion criteria

Working within the Toulouse conurbation and agreeing to participate in the cohort during the cross-sectional study, carried out between September 2002 and April 2003.

## MATERNELLE (PRESCHOOL) - SCHOOL CHILDREN COHORT: EARLY DETECTION OF SPECIFIC VERBAL AND WRITTEN LANGUAGE DEVELOPMENTAL DISORDERS IN CHILDREN.

Last update : 01/01/2019

Head :

Dellatolas Georges

Responsible organization :

COMMUNAUTE DE CREUSOT-MONTCEAU LES MINES, SAONE ET LOIRE

### Type of database

Study databases

### Study databases (details)

Cohort study

### Main objective

Early detection of specific language disorders and written language-learning difficulties in preschool.

### **Inclusion criteria**

All children attending preschools in the Creusot-Montceau-les-Mines community during academic years 2001 and 2002.

## **- LONGITUDINAL STUDY ON THE RELATIONSHIP BETWEEN PREMATURE BIRTH AND PERINATAL DEPRESSION**

Last update : 09/05/2017

Head :

Dayan Jacques, U1077 Neuropsychologie cognitive et neuroanatomie fonctionnelle de la mémoire humaine  
Équipe/activité :- Mémoire et amnésies

Responsible organization :

Institut national de la santé et de la recherche médicale - Inserm

### **Type of database**

Study databases

### **Study databases (details)**

Longitudinal study (except cohorts)

### **Main objective**

To determine the risk of premature birth based on the depressive state of the mother.

### **Inclusion criteria**

- female
- between 18 and 45 years old
- between the 20th and 28th week of pregnancy

## **DISCO - CASE-CONTROL STUDY ON THE IMPACT OF "THE BACK GUIDE (LE GUIDE DU DOS)" EDUCATIONAL PROGRAMME ON THE DEVELOPMENT OF COMMON LOWER BACK PAIN IN GENERAL MEDICINE**

Last update : 09/05/2017

Head :

Fernand Coriat, Direction Médicale Unité Thérapeutique Antalgie

Responsible organization :

Laboratoire Aventis

### **Type of database**

Study databases

## Study databases (details)

Case control study

### Main objective

To assess the impact of an educational programme on the chronic state of common lower back pain, through written information in "The Back Guide", as well as the usual verbal information provided to all lower back pain patients.

### Inclusion criteria

- male and female
- aged 18+
- with common lower back pain

## IREB SURVEYS - CROSS-SECTIONAL STUDIES ON ALCOHOL AND YOUNG PEOPLE - IREB

Last update : 09/05/2017

Head :

Com-Ruelle Laure

Choquet Marie, U 669

Responsible organization :

Institut de Recherches Scientifiques sur les Boissons (IREB)

Institut de Recherche et Documentation en Economie de la Santé (IRDES)

### Type of database

Study databases

## Study databases (details)

Repeated cross-sectional studies (except case control studies)

### Main objective

To provide a snapshot of the attitudes and behaviours of young French people attending or not attending school with regards to alcohol and associated risks. The use of repeated cross-sectional survey methodology also allows for a better understanding of consumption trends over time.

### Inclusion criteria

- between 13 to 20 years old for the 1996 and 2003 surveys and 13 to 24 years old for the 2007 survey.
- young people living in a household, outside an institution, whatever their circumstances (living with parents or not, active or not, etc.).
- living in metropolitan France excluding Corsica.

## HID - REPEATED CROSS-SECTIONAL STUDY: DISABILITY, INCAPACITY, DEPENDENCE

Last update : 09/05/2017



Head :

Boutron Isabelle, Département Epidémiologie, Biostatistique et Recherche CliniqueINSERM, U738

Responsible organization :

AP-HP - Assistance Publique - Hôpitaux de Paris Hôpital Bichat

Insee - Institut national de la statistique et des études économiques

## Type of database

Study databases

## Study databases (details)

Repeated cross-sectional studies (except case control studies)

## Main objective

To determine the impact of health problems on physical integrity, everyday life and social life for individuals.

## Inclusion criteria

- male and female
- surveyed on "everyday life and health"
- elderly or disabled individuals

# SIP - ENQUÊTE TRANSVERSALE SANTÉ ET ITINÉRAIRE PROFESSIONNEL

Last update : 01/01/2020

Head :

Barnay Thomas, Santé et itinéraire professionnel

Bouhia Rachid, Santé et itinéraire professionnel

Mette Corinne

Responsible organization :

DREES

DARES

## Type of database

Bases de données issues d'enquêtes

## Study databases (details)

Etudes transversales répétées (hors enquêtes cas-témoins)

## Main objective

L'enquête SIP vise d'une part à mieux connaître les déterminants de la santé liés aux conditions de travail et aux parcours professionnels : comment les conditions physiques et organisationnelles du travail, les expositions aux risques, les discontinuités et les ruptures du parcours professionnel contribuent-elles, dans la durée, à construire, à préserver ou à fragiliser leur état de santé ?

L'enquête devrait d'autre part permettre de repérer l'impact de l'état de santé sur le parcours professionnel des personnes, les aléas de carrière ou les discriminations éventuelles qu'elles ont pu rencontrer : en quoi des événements de santé, qu'ils soient déterminés ou non par l'environnement professionnel, induisent-ils des ruptures ou au contraire sont-ils compatibles avec une continuité de l'itinéraire professionnel, en fonction des ressources individuelles et collectives dont disposent les personnes ? Là aussi, il est important de prendre en compte d'éventuels décalages temporels.

Un objectif complémentaire a été fixé pour la vague 2010 : disposer, conformément aux recommandations intermédiaires du Collège d'expertise sur le suivi statistique des risques psychosociaux au travail, d'une observation des risques psychosociaux encourus par l'ensemble de la population active occupée au moment de l'enquête.

### **Inclusion criteria**

L'enquête a été effectuée auprès d'individus vivant en ménage ordinaire en France métropolitaine. Les individus étaient âgés de 20 à 74 ans en 2006 et ont été interrogés quelle que soit leur situation vis-à-vis du marché du travail (actifs occupés, chômeurs ou inactifs, retraités, etc.).

## **HSM-HSI - ETUDE TRANSVERSALE HANDICAP-SANTÉ (2008)**

Last update : 01/01/2019

Head :

Bouvier Gérard, Division Enquêtes et Etudes Démographiques

Robert-Bobée Isabelle, Division Enquêtes et Etudes Démographiques

Responsible organization :

Insee

DREES

### **Type of database**

Bases de données issues d'enquêtes

### **Study databases (details)**

Etudes transversales non répétées (hors enquêtes cas-témoins)

### **Main objective**

Les objectifs de l'enquête Handicap-Santé (volets 'Ménages' et 'Institutions') sont donc :

d'estimer le nombre des personnes handicapées ou dépendantes (en assurant autant que possible une comparabilité avec HID) ;

de relever la nature, la quantité et l'origine des aides existantes ainsi que les besoins non satisfaits ;

de mesurer les désavantages sociaux subis des personnes handicapées ;

d'évaluer les flux d'entrée en incapacité (à partir d'une interrogation rétrospective sur la date à laquelle les incapacités déclarées sont apparues) ;

de mesurer certains indicateurs annexés à la loi de santé publique ;

d'intégrer les modules européens élaborés par Eurostat (ECHIS).

### **Inclusion criteria**

L'échantillon est tiré dans la base des répondants à l'enquête de filtrage Vie Quotidienne et Santé (VQS),

effectuée en 2007 auprès d'environ 140 000 ménages. Tirage aléatoire  
Pour la partie "Institutions", 2200 structures volontaires

## **HID - ETUDE LONGITUDINALE HANDICAP-INCAPACITÉ- DÉPENDANCE**

Last update : 09/05/2017

Head :

Robert-Bobée Isabelle, Division Enquêtes et Etudes Démographiques

Responsible organization :

Insee

### **Type of database**

Bases de données issues d'enquêtes

### **Study databases (details)**

Etudes longitudinales (hors cohortes)

### **Main objective**

Trois objectifs : mesurer le nombre des personnes handicapées ou dépendantes ; évaluer les flux d'entrée et de sortie en incapacité ; relever la nature, la quantité, et les fournisseurs d'aides existantes, ainsi que les besoins non satisfaits

### **Inclusion criteria**

Tirage aléatoire (avec stratification) à partir de l'enquête de filtrage, dite "Vie Quotidienne et Santé" (VQS) qui a été adjointe au recensement de la population de mars 1999

Enfants internes des institutions relevant des annexes 24

Adultes handicapés accueillis en foyers d'hébergement, foyers de vie, MAS, FDT et structures expérimentales

Personnes hospitalisés dans un établissement psychiatrique

Personnes prises en charge dans des établissements en faveur des personnes âgées

## **SSIAD - ETUDE TRANSVERSALE RÉPÉTÉE DES SERVICES DE SOINS INFIRMIERS À DOMICILE**

Last update : 01/01/2020

Head :

Bertrand Dominique

Responsible organization :

DREES

### **Type of database**

Bases de données issues d'enquêtes

### **Study databases (details)**

Etudes transversales répétées (hors enquêtes cas-témoins)

### **Main objective**

Cette enquête permet de suivre l'activité des services de soins infirmiers à domicile, dispensés aux personnes âgées dépendantes ou aux personnes handicapées à leur domicile et dans les établissements non médicalisés. Le recueil d'informations apportera aux ministères en charge de la santé et des solidarités, des éléments nécessaires pour définir une politique de prise en charge de la dépendance des personnes âgées et handicapées à domicile.

Ces informations intéressent également les agences régionales de santé (ARS), notamment pour la mise en œuvre des schémas départementaux et des politiques locales en faveur des personnes âgées, ainsi que les autorités publiques à tous les échelons locaux.

### **Inclusion criteria**

Les services de soins infirmiers à domicile situés en France métropolitaine et dans les départements d'outre-mer (DOM)

## **DEVENIR - CROSS-SECTIONNAL STUDY ON PATIENTS' OUTCOME HOSPITALIZED FOR CARDIAC INSUFFICIENCY AND MONITORING IN PRIVATE CARDIOLOGY**

Last update : 08/01/2018

Head :

Cohen-Solal Alain

Responsible organization :

MENARINI France

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Describe modalities for the care of patients followed for cardiac insufficiency in private cardiology.

### **Inclusion criteria**

Men or women aged over 18 years having cardiac insufficiency hospitalized during the 18 months preceding inclusion consultation for cardiac insufficiency monitoring.

## **ICAPS 2 - ÉTUDE INTERVENTIONNELLE SUR DES ÉLÈVES DE 6ÈME : PROMOTION DE L'ACTIVITÉ PHYSIQUE**

Last update : 09/05/2017

Head :

Simon Chantal

Responsible organization :  
INPES

### **Type of database**

Bases de données issues d'enquêtes

### **Study databases (details)**

Etudes longitudinales (hors cohortes)

### **Main objective**

L'objectif est d'évaluer l'efficacité d'une action visant à lutter contre la sédentarité et à promouvoir l'activité physique sous toutes ses formes pour diminuer le risque cardiovasculaire et agir sur la corpulence

### **Inclusion criteria**

Inclusion de 1 050 élèves scolarisés en 6<sup>ème</sup> en 2002 (entre le 06/09/2002 et le 23/10/2002) dans huit collèges du Bas-Rhin

## **NORMANUT - ENQUÊTE TRANSVERSALE SUR LA PRÉVALENCE DES TROUBLES NUTRITIONNELS EN HAUTE NORMANDIE**

Last update : 01/01/2018

Head :  
Déchelotte Pierre  
Ladner Joel

Responsible organization :  
CHU Rouen

### **Type of database**

Bases de données issues d'enquêtes

### **Study databases (details)**

Etudes transversales répétées (hors enquêtes cas-témoins)

### **Main objective**

Normanut 1 : Le but de l'étude est d'évaluer, auprès de la population consultant en médecine générale, la prévalence des différents troubles nutritionnels, et d'autre part analyser leurs relations avec les comportements alimentaires et les déterminant sociaux.

Normanut 2: Evaluer la prévalence des troubles nutritionnels et son évolution chez des patients consultant en médecine générale en Haute-Normandie

### **Inclusion criteria**

## **PELAGIE - ASSESSMENT OF THE IMPACT OF EXPOSURE TO OCCUPATIONAL AND ENVIRONMENTAL CHEMICALS DURING PREGNANCY ON PREGNANCY OUTCOME AND CHILD DEVELOPMENT**

Last update : 09/05/2017

Head :

Chevrier Cécile, IRSET U1085 INSERM Université Rennes 1

Responsible organization :

Institut National de la Santé et de la Recherche Médicale

### **Type of database**

Study databases

### **Study databases (details)**

Cohort study

### **Main objective**

General objective: to assess the impact of early-life exposure to occupational and environmental chemicals on pregnancy outcome and child development, including exposure to solvents, pesticides, persistent pollutants and chlorinated derivatives in water.

Secondary objectives:

- to identify environmental sources and individual determinants (socio-demographic characteristics, lifestyle, genetic susceptibility) of exposure to pollutants measured by biomarkers
- to investigate the respective roles of prenatal exposure and exposure during childhood

### **Inclusion criteria**

Women in their first trimester of pregnancy (less than 19 weeks amenorrhoea) living in one of the "départements" participating in the study.

## **DAT'AIDS - ETUDE TRANSVERSALE SUR LES PATIENTS SÉROPOSITIFS SUIVIS DANS DES CHU DE FRANCE**

Last update : 09/05/2017

Head :

Pozot-Martin Isabelle

Cuzin Lize

Raffi François , Service de Maladies Infectieuses

Pugliese Pascal , Service d'Infectiologie

Responsible organization :

CHU

## Type of database

Bases de données issues d'enquêtes

## Study databases (details)

Etudes transversales non répétées (hors enquêtes cas-témoins)

## Main objective

Les objectifs de la cohorte sont :

- Fournir des caractéristiques épidémiologiques et démographiques
- Déterminer la proportion de patients traités/non traités
- Déterminer les profils viro-immunologique et thérapeutique
- Déterminer les caractéristiques des patients nouvellement diagnostiqués

## Inclusion criteria

Patients venus au moins une fois en consultation dans les CHU participants entre 01/06/04 e le 01/06/05

# - ETUDE TRANSVERSALE SUR DES PATIENTS DIABÉTIQUES : UTILISATION DU SCORE EPICES POUR DÉTERMINER LA RELATION CONTRÔLE GLYCÉMIQUE/COMPLICATIONS DU DIABÈTE ET LES CONDITIONS INDIVIDUELLES DE PRIVATION

Last update : 09/05/2017

Head :

Bihan Hélène, Centre de Recherche en Nutrition Humaine (CRNH)  
Cohen Régis, Internal Medicine and Endocrinology Departments

Responsible organization :  
CETAF

## Type of database

Bases de données issues d'enquêtes

## Study databases (details)

Etudes transversales non répétées (hors enquêtes cas-témoins)

## Main objective

Cette étude à pour objectif d'utiliser le score EPICES (évaluation de la précarité et les inégalités dans les centres d'examens de santé ), afin de déterminer la relation entre le contrôle glycémique, les complications du diabète, et les conditions individuelles de privation

## Inclusion criteria

Patients admis dans le services de Médecine Interne et d'Endocrinologie, de l'Hôpital Avicenne, Bobigny, France

# - SURVEY ON SUPPORT FOR ILLICIT DRUG USERS WHO USE HEALTH AND SOCIAL SYSTEM SERVICES

Last update : 09/05/2017

Head :  
DREES

Responsible organization :  
Direction de la Recherche, des Études, de Évaluation et des Statistiques (DREES)

## **Type of database**

Study databases

## **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

## **Main objective**

- To assess the number of drug users who turn to social and health institutions;
- To improve knowledge of socio-demographic characteristics, consumption;
- To help in the decision and development of treatment policies for the population of drug users seeking treatment;

## **Inclusion criteria**

Inclusion of treatment centres:

- Institutions specialising in the reception and treatment of drug users
- health facilities: General public hospitals or specialising in psychiatry, private psychiatric hospitals in the public sector and nursing homes for mental illness
- social institutions involved in the prevention, counselling or accommodation of drug users.

All patients present in these institutions are included in the survey.

# **SAMENTA - MENTAL HEALTH AND ADDICTION AMONG HOMELESS PEOPLE IN ILE-DE-FRANCE**

Last update : 04/01/2019

Head :  
Arnaud Amandine  
Chauvin Pierre, INSERM U707, equipe DS3

Responsible organization :  
Observatoire du Samu Social de Paris  
INSERM - Institut National de la Santé et de la Recherche Médicale

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)



## Main objective

To estimate the prevalence of primary psychiatric disorders and addictions in the homeless population in Ile-de-France. To study medical care sought in order to improve therapeutic treatment and adapt accommodation/housing conditions. To calculate the size of the concerned population in Ile-de-France (representative survey)

## Inclusion criteria

Homeless people in Ile-de-France over the age of 18, who speak French and can understand and answer questions

## ENEIS - NATIONAL SURVEY ON SERIOUS ADVERSE EVENTS IN HOSPITALS

Last update : 01/01/2019

Head :

Michel Philippe

Responsible organization :

Direction de la recherche, des études, de l'évaluation et des statistiques (DREES)

## Type of database

Study databases

## Study databases (details)

Repeated cross-sectional studies (except case control studies)

## Main objective

The survey's primary objective is to estimate the incidence of serious adverse events observed in hospitals and their avoidable character - for events resulting from admission in medicine and surgery units of health institutions or arising in such units during hospitalization.

The secondary objectives are:

- 1) estimate the severity and avoidable portion of such events;
- 2) describe the immediate care-related causes of such events arising.

## Inclusion criteria

All patient stays present during the survey in shortstay medicine and surgery units

## SCOPE - PATIENTS WITH SCHIZOPHRENIC OR SCHIZOAFFECTIVE DISORDERS

Last update : 09/05/2017

Head :

Azorin Jean-Michel, SHU Psychiatrie adultes

Responsible organization :

CHU St Marguerite Marseille

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

To observe the progress of patients being treated for an acute psychotic episode over a 2-year period.

## Inclusion criteria

- Men or women
- adult
- diagnosed with schizophrenia or affected by schizoaffective disorder

## - ETUDE TRANSVERSALE SUR DES SUJETS ATTEINTS D'UN CANCER OU DU VIH : POINT DE VUE SUR LES CONDITIONS DE VIE AU TRAVAIL

Last update : 01/01/2019

Head :

Lhuillier Dominique, Centre de recherche sur le travail et le développement (EA 4132)

Waser Anne-Marie , Laboratoire interdisciplinaire pour la sociologie économique (UMR 3320)

Responsible organization :

CNAM, Centre de recherche sur le travail et le développement (EA 4132)

## Type of database

Bases de données issues d'enquêtes

## Study databases (details)

Etudes transversales répétées (hors enquêtes cas-témoins)

## Main objective

Il s'agit de recherches qualitatives privilégiant le point de vue de personnes atteintes de cancer ou du VIH sur leur vie au travail, les difficultés rencontrées comme les stratégies déployées pour construire et entretenir leur santé tout en relançant ou maintenant leur vie professionnelle. Elles complètent et nuancent bon nombre de résultats d'enquêtes quantitatives portant sur l'impact d'une pathologie sur l'emploi ou la trajectoire professionnelle qui montrent invariablement l'effet négatif potentialisé selon la catégorie sociale d'appartenance ou le statut d'emploi.

Cette étude permet d'aborder les différences, mais aussi les similitudes dans le retour ou le maintien en emploi de personnes touchées par le VIH ou le cancer.

## Inclusion criteria

Personnes atteintes de VIH ou d'un Cancer et étant volontaire pour participer à l'étude

# CRYOVAS - ETUDE LONGITUDINALE SUR DES PATIENTS PRÉSENTANT DES VASCULARITES CRYOGLOBULINÉMIQUES NON ASSOCIÉES À UNE HÉPATITE C

Last update : 01/01/2020

Head :

Terrier Benjamin

Responsible organization :

AP-HP

## Type of database

Bases de données issues d'enquêtes

## Study databases (details)

Etudes longitudinales (hors cohortes)

## Main objective

Très peu de données sur les vascularites cryoglobulinémiques mixte d'origine non-infectieuse sont disponibles dans la littérature. Nous rapportons les caractéristiques cliniques et biologiques initiales des patients avec vascularites cryoglobulinémiques mixte inclus dans l'enquête CryoVas.

## Inclusion criteria

Patients avec vascularites cryoglobulinémiques non associée au VHC

Diagnostic de vascularites cryoglobulinémiques entre janvier 1995 et juillet 2010

Cryoglobuline détectable dans le sérum

Vascularite systémique définie par la présence de manifestations cliniques de vascularite en association au moins à un purpura et/ou une preuve histologique

# - ETUDE RESTROSPECTIVE SUR LES DÉCÈS D'ENFANTS DE MOINS D'1 AN DANS L'AGGLOMÉRATION DE LILLE : MORTALITÉ INFANTILE, DÉFAVEUR ET PROXIMITÉ AUX INDUSTRIES POLLUANTES

Last update : 01/01/2019

Head :

Padilla Cindy, IRSET- Institut de recherche santé environnement et travail

Responsible organization :

EHESP - Ecole des Hautes Etudes en Santé Publique

Université Sorbonne Paris Cité

IRSET- Institut de recherche santé environnement et travail

## Type of database

Bases de données issues d'enquêtes

## Study databases (details)

Etudes transversales non répétées (hors enquêtes cas-témoins)

### **Main objective**

Étudier la relation entre les variabilités géographiques des inégalités de mortalité infantile et la défaveur socioéconomique et environnementale à une échelle fine de l'agglomération de Lille.

### **Inclusion criteria**

- cas de décès d'enfants de moins d'1 an
- vivant dans l'agglomération de Lille

## **POLLIN'AIR - ENQUÊTE TRANSVERSALE SUR LA SÉVÉRITÉ DE LA RHINITE ALLERGIQUE SAISONNIÈRE EN FONCTION DE LA PRÉSENCE DE POLLEN ET DE LA POLLUTION DE L'AIR**

Last update : 09/05/2017

Head :

Annesi-Maesano Isabella, Epidemiology of Allergic and Respiratory Diseases

Desqueyroux helene

JANKOWSKI Roger

KLOSSEK Jean-Michel

THIBAUDON Michel

DEMOLY Pascal, General Allergology-Rhinitis-Asthma

DIDIER Alain, didier.a@chu-toulouse.fr

Responsible organization :

Inserm

### **Type of database**

Bases de données issues d'enquêtes

### **Study databases (details)**

Etudes transversales non répétées (hors enquêtes cas-témoins)

### **Main objective**

Le but de ce travail était de modéliser le risque de souffrir d'une rhinite allergique saisonnière sévère (SAR) en fonction à la fois de la présence de pollen et de la pollution de l'air dans un grand échantillon de patients.

### **Inclusion criteria**

Les patients consultant un médecin pour une rhinite allergique non traitée entre Mai et Août 2004

Le diagnostic est confirmé par la présence de symptômes nasaux (obstruction nasale, rhinorrhée, éternuements et prurit nasal), un symptôme oculaire (prurit oculaire) et la sensibilisation allergique à des allergènes majeurs

Consentement des participants

## **SMILE - ÉTUDE TRANSVERSALE SUR DES PATIENTS MIGRAINEUX CONSULTANT EN MÉDECINE DE VILLE**

Last update : 09/05/2017

Head :

Radat Françoise, 33076 Bordeaux Cedex

Geraud Gilles

Responsible organization :

Laboratoires Schwartz Pharma. TNS Healthcare

Société Française d'Etudes des Migraines et Céphalées

### **Type of database**

Bases de données issues d'enquêtes

### **Study databases (details)**

Etudes transversales non répétées (hors enquêtes cas-témoins)

### **Main objective**

Les objectifs de l'étude SMILE étaient d'évaluer l'anxiété, le stress, la dépression, l'impact fonctionnel et les comportements d'adaptation chez les patients migraineux consultant en médecine de ville en France

### **Inclusion criteria**

Chaque médecin participant devait inclure les quatre premiers patients consécutifs consultant pour une migraine, âgés de plus de 18 ans et actuellement non traité avec une substance connue pour être dotée de propriétés prophylactiques migraine (utilisées spécifiquement pour la prévention des migraines ou pour d'autres raisons)

## **PARITE - ÉTUDE TRANSVERSALE SUR LA PRISE EN CHARGE PAR LES CARDIOLOGUES DE L'HYPERTENSION ARTÉRIELLE EN FONCTION DU SEXE**

Last update : 01/01/2020

Head :

Mounier-Vehier Claire

SIMON MADJLESSI Tabassome , UPMC-Paris-06

Responsible organization :

Novartis Pharma S.A.S

CHRU de Lille

### **Type of database**

Bases de données issues d'enquêtes

### **Study databases (details)**

Etudes transversales non répétées (hors enquêtes cas-témoins)

## Main objective

L'objectif principal de cette étude française observationnelle, transversale, multicentrique, a été d'évaluer en fonction du sexe, la prise en charge par des cardiologues libéraux de l'hypertension artérielle (HTA) et du risque cardiovasculaire global.

## Inclusion criteria

Patients consultant chez un cardiologue participant pour une hypertension essentielle

## MIRABEL - ÉTUDE TRANSVERSALE SUR DES PATIENTS ALLERGIQUES À L'ARACHIDE EN FRANCE

Last update : 09/13/2017

Head :  
Crépet Amélie

Responsible organization :  
ANSES, INRA, Allergyvigilance Network (RAV)

## Type of database

Bases de données issues d'enquêtes

## Study databases (details)

Etudes transversales non répétées (hors enquêtes cas-témoins)

## Main objective

L'objectif est de proposer des outils méthodologiques et opérationnels pour quantifier le risque allergique, de tester des scénarios de gestion et de produire une analyse coûts-bénéfices

## Inclusion criteria

Patients volontaires consultant chez un des allergologues participant ayant une allergie à l'arachide

## - ÉTUDE TRANSVERSALE SUR DES PATIENTS ATTEINTS DE SARCOÏDOSE : INCIDENCE ET CARACTÉRISTIQUES EN GUADELOUPE

Last update : 01/01/2019

Head :  
Cadelis Gilbert  
Cordel Nadege, EA 4546

Responsible organization :  
CHU Guadeloupe

## Type of database

Bases de données issues d'enquêtes

## Study databases (details)

Etudes transversales non répétées (hors enquêtes cas-témoins)

### Main objective

Définir l'incidence et les caractéristiques de la sarcoïdose en Guadeloupe

### Inclusion criteria

Patients avec une sarcoïdose confirmée histologiquement consultant au CHU de de Pointe-à-Pitre

## DAIFI: OUTCOME FOLLOWING INITIATION OF IN VITRO FERTILISATION (DEVENIR APRÈS INITIATION D'UN TRAITEMENT PAR FÉCONDATION IN VITRO) - COHORT OF COUPLES STARTING IN VITRO FERTILISATION

Last update : 01/01/2020

Head :

de La Rochebrochard Elise, Ined ? Inserm ? University of Paris-Sud 11 (Ined UR14/Inserm CESP 1018)

Responsible organization :

Institut National de la Santé et de la Recherche Médicale (National Institute for Health and Medical Research)

Institut National d'Etudes Démographiques (National Institute for Demographic Studies)

### Type of database

Study databases

## Study databases (details)

Cohort study

### Main objective

General aim: To assess the likelihood of long-term parental project fulfilment among a population of couples beginning IVF treatment.

Secondary aim: To assess the cumulative IVF success rate, treatment discontinuation and spontaneous births after stopping IVF treatment.

### Inclusion criteria

? female

? adult

? started IVF treatment between 2000 and 2002 in one of the IVF centres participating in the study.

## ERISCAM - COHORT OF FRENCH FAMILIES WITH IDENTIFIED MMR GENE MUTATION: ASSESSMENT OF CANCER RISK

Last update : 01/01/2019

Head :

Lasset Christine, Prevention and genetics epidemiology  
Bonadona Valérie  
Bonaiti-Pellie Catherine , Genetics epidemiology and human population structures  
Olschwang Sylviane , Medical Genetics and Functional Genomics

Responsible organization :  
Centre Léon Bérard

## **Type of database**

Study databases

## **Study databases (details)**

Cohort study

## **Main objective**

The aim is to reliably and accurately determine the risks of different tumours in subjects with MMR gene mutations.

## **Inclusion criteria**

French families with an identified MMR gene mutation recruited throughout 40 cancer genetic consultation centres participating in the study; all members of the Cancer and Genetics Network.

# **- COHORT OF FEMALES RECEIVING SINGLE BLASTOCYST TRANSFER DURING IN VITRO FERTILIZATION**

Last update : 09/05/2017

Head :

Guerif Fabrice, UMR 0085 PRC Reproductive Physiology and Behaviour  
Royet Dominique, UMR 0085 PRC Reproductive Physiology and Behaviour

Responsible organization :  
INRA

## **Type of database**

Study databases

## **Study databases (details)**

Cohort study

## **Main objective**

This study aims to improve knowledge regarding in vitro fertilisation and to better understand the relationship between the number of implanted embryos/blastocyst transfers and birth.

## **Inclusion criteria**

Inclusion criteria: females under 36 years of age;  
First or second attempt at in vitro fertilisation;



## EXACO - COHORT OF PATIENTS WITH CHRONIC OBSTRUCTIVE PULMONARY DISEASE

Last update : 01/01/2020

Head :

Masure Frédéric

Cortot Alexis

Schuck Stephane

Responsible organization :

Société de Pneumologie de Langue Française

### Type of database

Study databases

### Study databases (details)

Cohort study

### Main objective

The main aims of the study are to qualitatively and quantitatively describe the exacerbations in a cohort of COPD patients at different levels of severity and to confirm the existence of a sub-group of patients defined as frequent exacerbators who experience a high number of exacerbations over time.

Where applicable, to determine the threshold value (number of exacerbations) that distinguishes frequent exacerbations by assuming that 3 exacerbations per year, regardless of severity, are sufficient to class the patient as a ?frequent? exacerbator.

Several secondary aims will also be pursued.

To identify factors associated with frequent exacerbators.

To identify criteria more readily associated with exacerbation severity.

To confirm the link between frequent exacerbators/accelerated decline in lung function.

A validation of the VSRQ scale will be performed with the following metrics: replication; clinical validity through comparison with the St. George questionnaire; VSRQ unidimensional structure; VSRQ internal consistency; sensitivity to change; minimal important distance.

### Inclusion criteria

Respiratory physicians (private practice or hospital: CHG, CHU) will enrol patients aged 40 and over with stage II or III COPD according to SPLF criteria (2003); smokers or ex-smokers (>15 pack-years); stable and with post-bronchodilator FEV1 lower than or equal to 80% of predicted value and FEV1/VC relationship <70%. Patients must accept and complete the self-monitoring log on a monthly basis and, at each exacerbation, can be contacted by phone every three months for 4 years. Participants shall sign a consent form.

The following may not be included: patients with active tuberculosis, cancer (or who received cancer treatment in the last 3 years), diffuse bronchiectasis, cystic fibrosis, asthma (in clinical history), or any other diagnosed lung diseases (sarcoidosis, pulmonary fibrosis, pneumoconiosis, etc.). Other non-inclusion criteria include exacerbation one month prior to enrolment; absence of a telephone and

participation in another clinical or epidemiological study.

## **EPI3 - COHORT OF PATIENTS WITH THREE COMMON DISEASES (MUSCULOSKELETAL PAIN; SLEEP, ANXIETY AND DEPRESSIVE DISORDERS AND UPPER RESPIRATORY INFECTIONS): STUDY ON THE IMPACT OF HOMOEOPATHIC TREATMENT**

Last update : 01/01/2020

Head :

Abenhaim Lucien

Begaud Bernard, U 657 Inserm

Responsible organization :

Société LA-SER

### **Type of database**

Study databases

### **Study databases (details)**

Cohort study

### **Main objective**

The aim of these three groups was to assess the difference between conventional and homoeopathic treatment.

### **Inclusion criteria**

Patients with:

? musculoskeletal pain (MSP);

? mixed-depressive disorder (MADD)

? upper respiratory infections (URI)

## **KBP-CPHG - COHORT OF PATIENTS WITH PRIMARY BRONCHIAL LUNG CANCER**

Last update : 01/01/2020

Head :

Grivaux Michel, Pulmonology

Responsible organization :

Collège des Pneumologues des Hôpitaux Généraux

### **Type of database**

Study databases

### **Study databases (details)**

Cohort study

## Main objective

The main aim is to study patient mortality at 5 years and risk factors. Secondary aims are:  
? to describe the characteristics in this patient population, their cancer and treatment methods;  
? to estimate the survival rate at 1, 4 and 5 years;  
? to compare data obtained with data from the KBP-2000-CPHG study (same study type, centre and patient recruitment methods, patient inclusion criteria and several common items in both questionnaires).

## Inclusion criteria

All patients with histological and cytological BPC diagnosed between 01 January and 31 December 2000 and 01 January and 31 December 2010. Patients monitored by the pulmonology unit in one of the general hospitals centres (GHC).

## FOCUS - COHORT OF PATIENTS WITH HYPERTROPHIC CARDIOMYOPATHY: PROSPECTIVE SCREENING FOR FABRY DISEASE

Last update : 07/01/2018

Head :

Hagège Albert, Inserm U633 Cardiology Department at Georges Pompidou European Hospital [Hôpital Européen Georges-Pompidou]

Responsible organization :

Paris Public Hospitals (AP-HP)

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

To estimate the prevalence of Fabry disease in a population of men and women with hypertrophic cardiomyopathy (HCM) consistent with primary cardiomyopathy.

## Inclusion criteria

? men and women

? between 18 and 79 years old

? with hypertrophic cardiomyopathy

## SOHO-FRANCE - COHORT OF SCHIZOPHRENIC PATIENTS RECEIVING ANTIPSYCHOTIC DRUGS

Last update : 09/05/2017

Head :

Gasquet Isabelle

Responsible organization :  
Assistance publique?hôpitaux de Paris

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

To provide information on patients monitored as outpatients, regardless of antipsychotic treatment received and to provide answers, especially regarding treatment adherence and remission rates, as well as relapse rates over a three-year monitoring period.

## Inclusion criteria

Schizophrenic adult patients monitored as outpatients and randomly recruited by psychiatrists.

## - COHORT OF PATIENTS WITH DEPRESSION: TOXIC EFFECTS ON BRAIN FUNCTION

Last update : 01/01/2018

Head :

Falissard Bruno , Inserm U1018 ? Centre for Research in Epidemiology and Population Health ? Team: Gender, Sexual and Reproductive Health

Corruble Emmanuelle, INSERM U669 BICETRE UNIVERSITY HOSPITAL ? PSYCHIATRIC DEPARTMENT

Responsible organization :

INSERM ? National Institute of Health and Medical Research

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

To assess the impact of depression on the hippocampus.

## Inclusion criteria

Outpatients from selected physicians who:

? meet the DSM-IV criteria for major depressive disorder based on clinical assessment criteria;

? were tested for delayed memory function during two visits several weeks apart.

## SMP: FIRST-TIME INCARCERATION SURVEY - COHORT OF DETAINED INDIVIDUALS: MENTAL HEALTH STUDY

Last update : 09/05/2017

Head :

Falissard Bruno, INSERM U669

Responsible organization :

Direction générale de la Santé (DGS)

Direction de l'Administration Pénitentiaire (DAP)

## Type of database

Study databases

### Study databases (details)

Cohort study

### Main objective

Detailed description of this specific prison population and comparison with prevalent prison population (sub-sample of cross-sectional maisons d'arrêt survey); outcome of disorders at 1 month and qualitative analysis at 9 months.

(Initial aim: to determine specific factors for mental health deterioration in the short term. Aim is not achievable given the small number of individuals still present at the institution 9 months after their arrival).

### Inclusion criteria

Individuals randomly drawn from available records at the Fresnes National Observation Centre and the diagnosis contained.

## - COHORT OF LOW-INCOME SUBJECTS RECEIVING FOOD VOUCHERS AND DIETARY ADVICE

Last update : 01/01/2018

Head :

Bihan Hélène, Human Nutrition Research Centre

Mejean Caroline, UMR 1125 UREN Nutritional Epidemiology. Jouy-en-Josas Research Centre

Galan Pilar, UMR 1125 UREN Nutritional Epidemiology. Jouy-en-Josas Research Centre

Peneau Sandrine, UMR 1125 UREN Nutritional Epidemiology. Jouy-en-Josas Research Centre

Responsible organization :

INRA

Inserm

## Type of database

Study databases

### Study databases (details)

Cohort study

### Main objective

To assess the effectiveness of using food vouchers to promote the consumption of fruit and vegetables among low-income populations.

### **Inclusion criteria**

People on low incomes (assessed by deprivation score within healthcare clinics) who are randomly selected.

18-60 years old.

## **HHORAGES - COHORT OF CHILDREN FROM WOMEN EXPOSED TO SYNTHETIC HORMONES DURING PREGNANCY**

Last update : 09/05/2017

Head :

Soyer-Gobillard Marie-Odile

Responsible organization :

HHORAGES-FRANCE

### **Type of database**

Study databases

### **Study databases (details)**

Cohort study

### **Main objective**

The aim of this cohort is to establish the causality between taking synthetic sex hormones during pregnancy and the onset of disorders in children from these pregnancies over a relative long term period.

Particular attention is paid to mental health disorders including: recurrent depression, anorexia, bulimia, manic depression, and schizophrenia that are related or not related to defects and/or malformations.

### **Inclusion criteria**

Volunteers born between 1946 and 1995.

## **EPIPOI - COHORT OF OVERWEIGHT CHILDREN OR CHILDREN AT RISK OF BEING OVERWEIGHT (HAUTE-GARONNE): SCREENING AND MANAGEMENT**

Last update : 09/05/2017

Head :

Jouret Béatrice, INSERM U558

Responsible organization :

Toulouse University Hospital

### **Type of database**

Study databases

## Study databases (details)

Cohort study

### Main objective

The general aims of the EPIPOI study are:

- ? To screen overweight children or children at the risk of being overweight at 3-4 years old and who are then followed up by their treating physicians with or without support from RéPOP (paediatric obesity prevention and management networks) in Midi-Pyrénées (RéPOP Toulouse);
- ? to carry out educational intervention programmes for children and to assess their effectiveness;
- ? to train private practitioners in screening and managing overweight and obese children;
- ? to determine the prevalence of excess weight and associated factors in children 3-4 years old.
- ? to characterize the life styles of children aged 3-4 years old.

### Inclusion criteria

Overweight children or children at risk of being overweight from voluntary preschools in Haute-Garonne.

## LOUIS MOURIER COHORT - RETROSPECTIVE COHORT OF NEWBORNS FROM MOTHERS USING MULTIPLE DRUGS

Last update : 01/01/2018

Head :

Simmat-Durand Laurence, CERMES 3 ? Centre for Research in Medicine, Science, Health, Mental Health and Society

Responsible organization :

Paris Descartes University

### Type of database

Study databases

## Study databases (details)

Cohort study

### Main objective

To analyse neonatal morbidity in terms of maternal psychoactive substance (PAS) use and within a sociodemographic context.

### Inclusion criteria

Newborns from mothers who were using at least 2 psychoactive substances at the start of pregnancy.

## INCANT - COHORT OF ADOLESCENTS TREATED FOR CANNABIS ABUSE AND ADDICTION

Last update : 12/01/2019

Head :

Falissard Bruno, INSERM U669

Phan Olivier, INSERM U669

Jouanne Celine

Responsible organization :

INSERM - National Institute of Health and Medical Research

Mission Interministérielle de Lutte contre les Drogues et les Toxicomanies (MILDT)

## **Type of database**

Study databases

## **Study databases (details)**

Cohort study

## **Main objective**

To assess the interest in a specific therapy; Multidimensional Family Therapy (MDFT) is given to adolescents who are misusing or addicted to cannabis.

To develop a therapeutic method in the European context, especially in France, that is effective in the field of adolescent addiction; to develop psychotherapy evaluation methods.

## **Inclusion criteria**

Presenting with a diagnosis for cannabis abuse and addiction and to having at least one parent (or legal guardian) agreeing to participate in the treatment and study. It should be noted that the misuse of psychoactive substances, behavioural and/or psychiatric disorders often associated with cannabis do not constitute grounds for exclusion.

# **PHASTE - COHORT OF PATIENTS IN GENERAL MEDICINE: HIGH BLOOD PRESSURE STUDY**

Last update : 09/05/2017

Head :

Mourad Jean-Jacques

Responsible organization :

Comité Français de Lutte contre l'HyperTension Artérielle

## **Type of database**

Study databases

## **Study databases (details)**

Cohort study

## **Main objective**

The aim is to extensively map hypertensive patients monitored by general practitioners, focusing on level of pulse pressure.



## **Inclusion criteria**

Subjects with uncontrolled hypertension (SBP >140 mmHg and/or DBP >90 mmHg), either treated or not treated.

## **- CASE-CONTROL STUDY ON THE RELATION BETWEEN SYSTEMIC SCLEROSIS AND OCCUPATIONAL EXPOSURE**

Last update : 01/01/2018

Head :

Marie Isabelle, INSERM U 905

Responsible organization :

Rouen University Hospital

## **Type of database**

Study databases

## **Study databases (details)**

Case control study

## **Main objective**

The aim of the study is to investigate the relation between systemic sclerosis and occupational exposure, and to define the risk of systemic sclerosis in male and female patients that is linked to occupational exposure.

## **Inclusion criteria**

Patients with systemic sclerosis (declared) consulting at one of 3 participating centres (Departments of Internal Medicine, Dermatology and Pneumology in Rouen, Elbeuf and Le Havre) between 2005 and 2008.

Controls: Subjects with a chronic illness or an acute illness who do not have: a connective tissue disorder; systemic vasculitis and other systemic autoimmune disorders; neoplasia and chronic interstitial lung disease.

## **KARU-PROSTATE - CASE-CONTROL STUDY ON THE IMPACT OF CHLORDECONE EXPOSURE AND RISK OF PROSTATE CANCER IN THE WEST INDIES**

Last update : 09/05/2017

Head :

Multigner Luc , U1085 Research Institute of Health, Environment and Work

Responsible organization :

Inserm - Institut national de la santé et de la recherche médicale

## **Type of database**

Study databases

## Study databases (details)

Case control study

### Main objective

To identify the environmental and genetic risk factors of prostate cancer and to assess the impact of chlordecone exposure in the onset of the disease.

### Inclusion criteria

? Male

? Adult

? Caribbean native (Guadeloupe, Martinique, Haiti, Dominica)

## - CASE-CONTROL STUDY ON PATIENTS WITH ADDICTIVE BEHAVIOUR

Last update : 09/05/2017

Head :

Falissard Bruno, Inserm U1018 ? Centre for Research in Epidemiology and Population Health ? Team: Gender, Sexual and Reproductive Health

Godart Nathalie

Responsible organization :

INSERM - National Institute of Health and Medical Research

### Type of database

Study databases

## Study databases (details)

Case control study

### Main objective

To verify whether anxiety disorders are more common in women with eating disorders.

### Inclusion criteria

Patients with eating disorders (multicentric random draw).

## ISICA - CASE-CONTROL STUDY ON THE RISK OF BREAST CANCER THROUGH INSULIN USE

Last update : 01/01/2019

Head :

Abenhaim Lucien

Grimaldi-Bensouda Lamiae, C121-SC of Vie-Pharma fourth group; U657 Pharmacoepidemiology and Evaluation of the Impact of Health Products on the Population?Pasteur Institute

Responsible organization :

Sanofi-Aventis  
La-Ser

## Type of database

Study databases

## Study databases (details)

Case control study

## Main objective

The aim is to assess the possible relation between using individual insulins (duration of exposure, administered dosage) and increased breast cancer risk.

## Inclusion criteria

For case subjects:

? Breast cancer diagnosed between 01 January 2008 and 30 June 2009

? Diabetes confirmed by the patient and anaesthesia records.

For controls:

? No breast cancer

? Diabetes confirmed by the patient and anaesthesia records

# EUROSC - THE EUROPEAN SCHIZOPHRENIA COHORT

Last update : 09/05/2017

Head :

Jean Michel Azorin

Bebbington Paul

Responsible organization :

Royal Free and University College Medical School.. Pour la partie Francaise : Hôpital Sainte Marguerite ,  
Marseille

## Type of database

Study databases

## Study databases (details)

Cohort study

## Main objective

The main aim of EuroSC is to report on the different types of treatment and care, as well as clinical outcome.

Secondary aims include: assessment of treatment requirements in relation to outcome, calculation of resource usage with regards to different care methods and identification of prognostic factors.

## Inclusion criteria

Patients between 18 and 64 years old at baseline;  
Established schizophrenia diagnosis;  
Participant consent

## **SMP: PREVALENCE STUDY - CROSS-SECTIONAL STUDY OF THE MENTAL HEALTH OF PRISONERS: PREVALENCE STUDY**

Last update : 10/20/2017

Head :

Falissard Bruno, Inserm U1018 - Centre de Recherche en Épidémiologie et Santé des Populations Équipe:  
Genre, Santé sexuelle et Reproductive  
Reynaud Michel

Responsible organization :

Direction générale de la Santé (DGS)  
Direction de l'Administration Pénitentiaire (DAP)

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To calculate the prevalence of psychiatric disorders in the prison environment

### **Inclusion criteria**

Prisoners (male and female) in institutions selected randomly in the protocol

## **- CROSS-SECTIONAL STUDY ON THE LABOUR FORCE IN LORRAINE: RELATIONSHIP BETWEEN LIFESTYLE/LIVING CONDITIONS, WORK AND WORKPLACE ACCIDENTS**

Last update : 10/20/2017

Head :

Falissard Bruno , Inserm U1018 - Centre de Recherche en Épidémiologie et Santé des Populations Équipe:  
Genre, Santé sexuelle et Reproductive  
Chau Nearkasen, U 669

Responsible organization :

INSERM - Institut National de la Santé et de la Recherche Médicale

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To study the relationship between work, lifestyle/life conditions and workplace accidents

### **Inclusion criteria**

Random selection from the Lorraine labour force

## **- CROSS-SECTIONAL STUDY OF THE BEHAVIOUR OF MOTHERS WITH REGARD TO THEIR BABIES' SLEEPING POSITION: EFFECTS OF THE LAST SUDDEN INFANT DEATH SYNDROME PREVENTION CAMPAIGN**

Last update : 10/20/2017

Head :

Falissard Bruno, INSERM U669

Vaivre-Douret Laurence, Inserm U483

Responsible organization :

INSERM - Institut National de Santé et Recherche Médicale

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To assess the sleeping practices of babies in two maternity wards and at the family home. To thereby evaluate the admissibility and application of the public health message regarding the prevention of sudden infant death syndrome.

### **Inclusion criteria**

Two general public samples were thus established for a one-month period in two maternity wards (Port-Royal and Créteil) and a maternal and infant protection (PMI) centre connected to the Port-Royal maternity ward.

## **ADOTECNO - CROSS-SECTIONAL STUDY OF THE HARMFUL USE OF ALCOHOL AND CANNABIS IN THE GENERAL POPULATION: VALIDATION OF A TRACKING QUESTIONNAIRE: THE CRAFFT-ADOSPA**

Last update : 10/20/2017

Head :

Falissard Bruno, Inserm U1018 - Centre de Recherche en Épidémiologie et Santé des Populations Équipe:  
Genre, Santé sexuelle et Reproductive  
Karila Laurent, INSERM U669

Responsible organization :  
INSERM - Institut National de la Santé et de la Recherche Médicale

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

To validate two levels of risk of the CRAFFT-ADOSPA for tracking the harmful use of alcohol, cannabis and other drugs.

## **Inclusion criteria**

High school and university students (Île-de-France)

# **- LONGITUDINAL STUDY OF PESTICIDE EXPOSURE IN OUTDOOR FARMING - CALVADOS**

Last update : 10/20/2017

Head :  
Lebailly Pierre, Groupe Régional d'Études sur le Cancer

Responsible organization :  
Université de Caen

## **Type of database**

Study databases

## **Study databases (details)**

Longitudinal study (except cohorts)

## **Main objective**

To demonstrate the contamination of farmers who are exposed to pesticides while treating their outdoor agriculture.

To determine the link between pesticide exposure for a farmer and the probability of developing cancer.

## **Inclusion criteria**

- male
- farmer
- using pesticides on crops

# EPIREUN - CROSS-SECTIONAL STUDY OF PATIENTS WITH NEWLY DIAGNOSED SEIZURES ON THE ISLAND OF RÉUNION

Last update : 10/20/2017

Head :

Preux Pierre-Marie, Neuroépidémiologie tropicale

Responsible organization :

University of Limoges

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

To evaluate the incidence of seizures in Réunion's population.

To determine the risk factors specific to Réunion among epileptic patients

## Inclusion criteria

- male or female
- resident of Réunion
- newly diagnosed as epileptic

# - CROSS-SECTIONAL STUDY OF THE INFLUENCE OF EXPOSURE TO GLYCOL ETHERS ON SEMEN QUALITY

Last update : 10/20/2017

Head :

Multigner Luc, Inserm U625

Responsible organization :

Institut National de la Santé et de la Recherche Médicale  
Université Rennes 1

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

To research the effect of past and current exposure to glycol ethers on semen quality.

## Inclusion criteria

- homme
- âgé entre 20 et 55 ans
- employé de la ville de Paris
- n'ayant pas de facteurs de risque d'infertilité

## **- CROSS-SECTIONAL STUDY IN THE GENERAL POPULATION [OF] METABOLIC SYNDROME AND IMPAIRED VENTILATORY FUNCTION**

Last update : 10/20/2017

Head :

Leone Nathalie, U700, Faculté de Médecine Xavier Bichat, Université Denis Diderot

Responsible organization :

Institut national de la Santé et de la Recherche Médicale

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To study the risks of deteriorating ventilatory function based on a metabolic syndrome.

### **Inclusion criteria**

- male or female
- age between 16 and 96 years
- living in Paris

## **- CASE-CONTROL SURVEY ON THE PARTICIPATION OF ADOLESCENTS IN ORGANISED SPORTS ACTIVITIES**

Last update : 10/20/2017

Head :

Sarrazin Philippe, "Laboratoire Sport et Environnement Social" (E.A. 3742) - UFR APS Université J. Fourier  
Boiché Julie, Laboratoire Epsilon EA Dynamique des Capacités Humaines et des Conduites de Santé

Responsible organization :

Université de la Réunion

### **Type of database**

Study databases

### **Study databases (details)**



Case control study

### **Main objective**

The objective of this survey is to study many determinants of discontinuation of an organised sports activity in French adolescents, in order to determine the distal and proximal factors of the discontinuation.

### **Inclusion criteria**

Subjects practising one of the following sports in a club: football, basketball, handball, rugby, skiing, judo, gymnastics, horse riding, cycling, climbing, tennis and table tennis. For athletes who have stopped:  
- stopped a year ago.

## **LYCOLL - CROSS-SECTIONAL STUDY OF YEAR-10 AND YEAR-11 STUDENTS: INFLUENCE OF A CHANGE OF ENVIRONMENTAL CONTEXT ON HIGH-RISK BEHAVIOURS**

Last update : 10/20/2017

Head :  
Binder Philippe

Responsible organization :  
INSERM - Institut National de la Santé et de la Recherche Médicale  
Observatoire Régional de la Santé Poitou-Charentes

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

To study the influence of change in environmental context on the consumption of psychoactive products and on other high-risk behaviours, such as suicide. To describe the conditions of social and family life, education, health condition and associated behaviours.

### **Inclusion criteria**

Students in year 10 and 11 assimilated in Charente-Maritime.

## **ALIMADOS - CROSS-SECTIONAL STUDY OF DIETARY BEHAVIOURS IN ADOLESCENTS**

Last update : 01/01/2019

Head :  
Pardo Véronique

Responsible organization :

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

To understand the dietary habits of adolescents considering cultural differences and population mixing at work today in France.

## **Inclusion criteria**

Be 12 to 19 years of age and live in the Alsace or Provence-Alpes-Côtes d'Azur region.

# **PPGP - PARIS PREVENTION IN GENERAL PRACTICE**

Last update : 10/20/2017

Head :

Rigal Laurent, Inserm U1018 - Centre de Recherche en Épidémiologie et Santé des Populations Team: Genre, Santé sexuelle et Reproductive

Responsible organization :

INSERM - Institut National de la Santé et de la Recherche Médicale

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

To analyse social inequalities in preventive practices in general medicine.

## **Inclusion criteria**

- hommes entre 35 et 64 ans pour la consommation de tabac et d'alcool
- femmes entre 50 et 69 ans pour le dépistage des cancers gynécologiques

# **- SURVEY ON HOMELESS PEOPLE IN PARIS (1994-1995)**

Last update : 10/20/2017

Head :

MARPSAT Maryse

Responsible organization :  
French National Institute for Demographic Studies (INED)

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

To study the conditions under which surveys of this type are conducted and extended on a larger scale, and to prepare recommendations for the purpose of achieving a better inclusion of homeless people in the general census of the population, as well as in surveys that are regularly carried out only among those with housing. A first examination of the processes leading to homelessness was also expected.

## **Inclusion criteria**

Persons 18 years of age or older, without a home or having housing but going to food and hot meal distribution centres and/or spending the night in shelters (emergency and long-term).

# **CONSTANT - CROSS-SECTIONAL STUDY OF THE GENERAL POPULATION IN GUADELOUPE: HEALTH, BEHAVIOUR AND SOCIAL ENVIRONMENT.**

Last update : 10/20/2017

Head :  
Atallah André, Cardiovascular rehabilitation unit

Responsible organization :  
CH de Basse-Terre

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

To provide data concerning the health, behaviour and social environment of the Guadeloupean population

## **Inclusion criteria**

Individuals selected randomly who agree to participate in the study

# **PHAPPG - LONGITUDINAL STUDY OF A PRECARIOUS GUADELOUPEAN POPULATION: ARTERIAL HYPERTENSION STUDY**

Last update : 10/20/2017

Head :

Atallah André, Cardiovascular rehabilitation unit

Responsible organization :

CH de Basse-Terre

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

The purpose of this study is to describe the prevalence and treatment of arterial hypertension in a population that is precarious but has good social coverage and good access to the health care system

## **Inclusion criteria**

The study's inclusion criteria for defining precarity were: receiving the minimum integration income (RMI) and/or unemployment and/or living exclusively on social and family services

# **COLLATION - CROSS-SECTIONAL STUDY OF CHILDREN IN NURSERY SCHOOL: PERCEPTION AND PRACTICE OF SNACKING IN AQUITAINE**

Last update : 01/01/2018

Head :

Thibault Hélène

Responsible organization :

ISPED

## **Type of database**

Study databases

## **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

## **Main objective**

The purpose is to make an inventory of the habitual practices of nursery school children in terms of snacks at school, including birthday treats, in Aquitaine. This inventory allows for the acquisition of information on the perceptions, practices, changes and expectations of the students, as well as guidance of the methods of intervention intended to improve these practices.

The renewal of this survey after intervention allows for the evaluation of its efficacy and the change in students' perceptions.

## Inclusion criteria

Nursery schools of the Bordeaux school district completing the questionnaire

## CE2 - CROSS-SECTIONAL STUDY OF YEAR-4 STUDENTS IN AQUITAINE SCHOOLS: DIETARY INTAKE AND CORPULENCE

Last update : 01/01/2019

Head :  
Thibault Hélène

Responsible organization :  
ISPED

### Type of database

Study databases

### Study databases (details)

Repeated cross-sectional studies (except case control studies)

### Main objective

The purpose of this study is:

- firstly, to find out the corpulence values and to therefore calculate the prevalence of overweight and obesity in year-4 children and, secondly, to gain knowledge of their dietary habits
- to complement the data initially collected with information on sedentary activity, physical activity and parents' socioprofessional categories.
- to gain knowledge of the evolution of these data between 2004-2005 and 2008-2009 and to evaluate the impact of the program's actions implemented since 2005.

### Inclusion criteria

Year-4 students in Aquitaine elementary schools

## - CROSS-SECTIONAL STUDY ON THE FUTURE OF INDIVIDUALS WHO HAVE STOPPED RECEIVING THE MINIMUM INTEGRATION INCOME (RMI)

Last update : 10/20/2017

Head :  
Accardo Jérôme, Division Conditions de Vie des Ménages  
Duée Michel, Household conditions

Responsible organization :  
Insee

### Type of database

Study databases

## Study databases (details)

Repeated cross-sectional studies (except case control studies)

### Main objective

The purpose of this methodological study is to specify the family configuration actually experienced by recipients of the minimum integration income (RMI).

### Inclusion criteria

All of the individuals receiving the minimum integration income (RMI) from the metropolitan France on 31 December 1996, except for old farmers enrolled in the Mutualité sociale agricole

## HSA - CROSS-SECTIONAL HANDICAP-HEALTH STUDY - INFORMAL CAREGIVERS

Last update : 01/01/2018

Head :

Soullier Noémie

Responsible organization :

DREES

Insee

### Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

### Main objective

The principal objective of the "Informal caregivers" survey is to characterise non-professional caregivers (family, friends, neighbours, etc.) for individuals who have stated that they have trouble performing certain daily activities due to a handicap, a health problem or their age in the 2008 Handicap-Health survey.

Conducted from April to September 2008, the survey includes 5,000 informal caregivers of handicapped individuals and dependent elderly individuals. This survey allows for the description and analysis of the living conditions and quality of life of individuals providing support to individuals they are close with (family, neighbours, friends, etc.) who stated that they have trouble in activities of daily life in the Handicap-Health survey in ordinary households in 2008 (handicapped or elderly individuals).

It addresses an interest identified in the recent public debate (conference on Family in 2006, decree dated January 2007 allowing the arrangement of the right to respite for family caregivers ), and addresses the initial questions from the Handicap-Health survey.

It allows for knowledge of the motivations and manifestations of informal support, description and analysis of living conditions and quality of life of caregivers and the manner in which non-professional aid connects or does not connect with aid provided by professionals. The survey also allows for the detailed measurement of the consequences of aid on the daily life of the caregivers, on the professional, family

and friendship plans.

The general objective of the survey can be broken down into five sub-objectives: to create a portrait of caregivers from the social network, to gain knowledge of the configuration of aid and the conditions under which it is provided, to see how the caregiver's activity is reconciled with that of other caregivers, to understand the consequences of aid and the resulting needs, and lastly to gain knowledge of the representations that the caregivers have regarding their economic consent to the aid provided.

### **Inclusion criteria**

The survey sample is made up of all of the caregivers named in the 2008 Handicap-Health survey, 16 years of age and older.

The "Informal caregivers" survey is a supplemental section of the Handicap-Health survey in ordinary households conducted by Insee in 2008: when a person indicates in the Handicap-Health survey that he or she is aided by one or more non-professional caregivers, a quick description of the caregiver and the nature of the aid that he or she provides is requested from the surveyed individual.

The contact information for the caregiver (or caregivers) is also collected (with a maximum of 3 cohabitating caregivers and 10 caregivers per surveyed individual): they constitute the sampling frame for the "Informal caregivers" survey.

The "Informal caregivers" study is limited to individuals 16 years of age or older. The individuals are questioned directly face to face by an Insee surveyor.

## **- CROSS-SECTIONAL STUDY OF WORKPLACE ACCIDENTS, OCCUPATIONAL ILLNESSES AND HANDICAPS - 2007**

Last update : 10/20/2017

Head :

Guggemos Fabien, Employment Division

Responsible organization :

Insee

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

The first European objective of the survey is to find out the number of workplace accidents and the number of days of work lost due to these workplace accidents. The survey will also allow for the determination of the number of illnesses caused or aggravated by the job, as well as the number of work days lost due to these illnesses.

The second European objective is to analyse the differences in results according to:

- the characteristics of the employer (economic activity, size of the business, etc.)

- the characteristics of the salaried worker's job (professional status, profession, full-time, part-time, duration of career, etc.)

This supplemental set of questions is intended specifically to enable comparisons with the results of the 2002 ad hoc module on the employment of handicapped individuals, in order to be able to evaluate the effects of the "Equal rights, opportunities, participation and citizenship of handicapped individuals act" adopted in 2004. The set of questions is divided into three parts. Part A address health problems within the last twelve months.

### **Inclusion criteria**

Individuals who are at least 15 years of age on 31 December 2007 and who are members of ordinary households.

## **- CROSS-SECTIONAL STUDY ON TRANSITIONS FROM ACTIVE LIFE TO RETIREMENT - 2006**

Last update : 10/20/2017

Head :

Guggemos Fabien, Employment Division

Responsible organization :

Insee

### **Type of database**

Study databases

### **Study databases (details)**

Repeated cross-sectional studies (except case control studies)

### **Main objective**

The first objective of this module is to compile an overview of the situation of the work-retirement transition. How is this transition made or how is it planned? Will the transition be made or will it be done directly from work to retirement? Have the individuals had or thought they would have a transition period (pre-retirement, transition to part-time)? The second objective pursued is the analysis of factors that may influence or have influenced this transition from work to retirement, as well as the determination of the factors that may postpone or that have postponed leaving the job market. These factors may be of different types: the general working conditions (planning of work schedules, better health conditions); other factors directly related to the job (training and obsolescence of skills); financial factors (financial incentives to stay in their job); personal factors (health, family reasons).

### **Inclusion criteria**

Individuals from 50 to 69 years of age on 31/12/2006 (individuals born between 1937 and 1956) who live in an ordinary household and whose household has been questioned for the 6th and final time by the Employment survey (6th outgoing)

## **- CROSS-SECTIONAL STUDY ON MIGRANTS' PATHS AND PROFILES**

Last update : 10/20/2017



Head :  
Lhommeau Bertrand, Bureau Jeunesse, famille

Responsible organization :  
DREES

## Type of database

Study databases

## Study databases (details)

Repeated cross-sectional studies (except case control studies)

## Main objective

This survey is intended to better understand the route, different trajectories (residential, professional, family) and use of social services of individuals who come to obtain a residence permit of at least one year in France and having a vocation to settle in France with stability. It also explores the vision and expectations that these migrants have for France and how these elements influence their trajectories in France. It specifically deals with understanding the migratory plan, its evolution over time, its individual or familial character, how it is modelled by institutions and the administrative categories with which the migrant is confronted. This allows for questions about the articulation between the migratory plan and the integration process. The two-part interview allows for introducing temporality into the understanding of this process, and to observe any connections between the evolution of the migratory plan and the living conditions encountered over the course of the first year with a residential permit.

## Inclusion criteria

Foreign individuals of legal age (18 years or older) accepted by the Agence nationale de l'accueil des étrangers et des migrations (ANAEM) between June and October 2006 who were asked to sign a reception and integration contract (CAI).

This survey deals with individuals eligible for a CAI, i.e.

- spouses of French citizens
- other relatives of French citizens
- beneficiaries of the family group
- refugees or members of their family
- individuals authorised due to personal and family connections in France (VPF 12 bis 7).

## - CROSS-SECTIONAL STUDY OF YEAR-10 STUDENTS IN THE LANGUEDOC-ROUSSILLON REGION

Last update : 01/01/2019

Head :  
Desmartin-Belarbi Valérie

Responsible organization :  
CREAI-ORS

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

### Main objective

The objectives are:

- to establish a collection of reproducible information in a school environment from representative samples of the enrolled population that will be able to provide better knowledge of the health condition of students, to identify risky behaviours of adolescents
- to optimise the practice of health care professionals of the national education system and to promote the homogenisation of practices.

### Inclusion criteria

Students in one of the high schools of the Montpellier school district.

## ECAAP - CROSS-SECTIONAL SURVEY ON THE ADDICTIVE BEHAVIOUR OF POLYNESIAN ADOLESCENTS

Last update : 01/01/2019

Head :

Renou Laurence  
Guignard Romain

Responsible organization :

INPES  
The French Polynesia Health Department

### Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

### Main objective

This is a school-based survey that provides an epidemiological and sociological assessment of the situation. A number of questions about their health, their living conditions, their everyday concerns, their level of well-being and their interests allow us to profile Polynesian adolescents.

### Inclusion criteria

Students from year 7 to the final year, Centres for Young Adolescents (CJA), Centre for Education for Appropriate Technologies for Development (CETAD), Development Education Centres (CED) and Family and Rural Homes (MFR) included (in French Polynesia).

## - CROSS-SECTIONAL STUDY OF PRECARIOUS SITUATIONS, PERCEIVED HEALTH AND MIXED ANXIETY-DEPRESSIVE DISORDERS

Last update : 10/20/2017

Head :

Royer Bernard

Gusto Gaelle

Tichet Jean

Responsible organization :

IRSA

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

The association between precarious situations and degradation of the state of health is widely accepted and links have been described for morbidity and mortality, and risk behaviours. In this context, our study aims to find a link between precarious situations, the quality of life experienced and the existence of mixed anxiety-depressive disorders

## **Inclusion criteria**

Subject being seen at one of the 12 health centres included in the study.

Insured subjects and beneficiaries of the general social security scheme, who have undergone a periodic health examination and are volunteering for the study

# **DIABÈTE - CROSS-SECTIONAL STUDY OF THE PREVALENCE OF TYPE 2 DIABETES IN PEOPLE FREQUENTING EMERGENCY SHELTERS**

Last update : 10/20/2017

Head :

Arnaud Amandine

Responsible organization :

Observatoire du Samu Social

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

As part of a systematic screening action proposed to people housed in CHUSI and CHUS, the main objective of the study is to estimate the prevalence of diabetes mellitus (known and unrecognised) in

these people.

The secondary objectives are:

- 1) Describe the characteristics of people with unrecognised diabetes mellitus,
- 2) Evaluate the management of known diabetes mellitus (insulin-treated or not).

### **Inclusion criteria**

The people interviewed are those accommodated in the aforementioned CHUSI and CHUS at the time of the survey and agreeing to participate.

Among the selected individuals, only the following were retained, those who:

-understand French

-have given their informed consent

People already knowing their diabetic condition will have samples of their blood taken to control their glycaemic balance and the medical part of the questionnaire will be filled in by the on-site physician

## **EPILEPSIE - CROSS-SECTIONAL STUDY OF EPILEPSY AMONG HOMELESS PEOPLE**

Last update : 10/20/2017

Head :

Arnaud Amandine

Responsible organization :

Samu Social

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

Estimate the prevalence of epilepsy in people frequenting SSP reception and accommodation facilities

Describe the types and characteristics of epilepsy

Assess the frequency of care and access to treatments

Assess the impact of the situation of exclusion on the frequency of seizures

### **Inclusion criteria**

All people going for medical consultations at the 3 CHUS and day centres (ESI) and present in the 3 CHUSI between 10 March and 18 April 2003

## **- PRIMINJECT: CROSS-SECTIONAL STUDY ON THE PATTERNS AND PRACTICES OF DRUG USE IN FRANCE**

Last update : 09/19/2017

Head :

Arwidson Pierre  
Guignard Romain  
Guichard Anne

Responsible organization :  
Santé publique France

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

Intravenous injection is the major mode of transmission of HCV among drug users, not only by needle sharing but also by the preparation equipment (container, filter, buffer, water), potentially contaminated by the blood of a contaminated person.

According to various sources, in recent years there have been significant changes in the patterns and practices of drug use in France, under the combined effect of the massive distribution of opiate substitution treatments (high-dose buprenorphine or BHD) from 1995 and the introduction of a wide variety of stimulants, including cocaine and synthetic drugs.

Often unplanned and practiced by a third party "initiator", the first injections present a high risk of contamination for the young initiate, often not well acquainted with injection techniques and the infectious risks and not in contact with first-line medical support structures.

In this context, a survey was put in place to identify the circumstances and practices of the first injection(s) among young drug users in France. An online methodology using the Internet has been developed to provide access and thus to discover user profiles that escape the more "classic" observation systems, especially those that are socially integrated or that are not in contact with support structures.

## **Inclusion criteria**

Be over 15 years old. Individuals who had never injected drugs were redirected to a short version of the questionnaire.

# **- LONGITUDINAL STUDY OF WOMEN WITHOUT DIABETES GIVING BIRTH: A STRATEGY FOR THE SELECTIVE DETECTION OF GESTATIONAL DIABETES**

Last update : 01/01/2020

Head :  
Cosson Emmanuel, Centre de Recherche en Nutrition Humaine d'Ile-de-France  
Carbillon Lionel, Department of Obstetrics and Gynecology

Responsible organization :  
Assistance Publique-Hôpitaux de Paris

## **Type of database**

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

### Main objective

The objectives are to evaluate:

- 1) the value of screening for risk factors for gestational diabetes
- 2) the prognostic value of risk factors for events related to gestational diabetes.

The study also shows changes in the prevalence of GDM risk factors

### Inclusion criteria

Woman without diabetes giving birth in Hôpital Jean Verdier between January 2002 and December 2010

## MFV - CROSS-SECTIONAL STUDY ON MIGRATIONS-FAMILY-AGEING IN FRENCH OVERSEAS DEPARTMENTS

Last update : 01/01/2020

Head :

Claude-Valentin Marie

Responsible organization :

INED

Insee

### Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

### Main objective

The MFV survey was carried out in a coordinated way in all the French overseas departments and focuses on the changes in family life, the sources and obstacles of social itineraries, the impact of migration and the evolution of intergenerational solidarity

### Inclusion criteria

Three main categories of key player were asked:

- \* Native people of the department
- \* Returning migrants: native people of the department who have had an emigration experience of more than six consecutive months and who were resettled there at the time of the survey
- \* Immigrants: the non-native people of the department who were settled there at the time of the survey, whether French or foreign

## PREVIPAP - CROSS-SECTIONAL STUDY OF SEXUALLY ACTIVE YOUNG FRENCH WOMEN ELIGIBLE FOR HPV VACCINATION

Last update : 01/01/2020

Head :

Mougin Christiane, Laboratory of Cellular and Molecular Biology, EA 3181, FED4234

Mercier Mariette, EA 3181, FED4234

Baudu Ariane, EA 3181, FED4234

Responsible organization :

CHU Jean Minjoz

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

The objective of this study was to assess the prevalence and risk factors for HPV infection in a cohort of sexually active young French women eligible for HPV vaccination

## **Inclusion criteria**

Sexually active women aged 15-23 years living in the geographical region of Franche-Comté in France, who attended a consultation in the gynaecology department of the Hôpital Universitaire de Besançon in France, between 1 January 1997 and 31 December 2007

# **- CROSS-SECTIONAL STUDY ON THE ELDERLY: ASSOCIATION OF BENZODIAZEPINE USE AND REPEATED FALLS**

Last update : 01/01/2018

Head :

Beauchet Olivier

Rossat Arnaud

Responsible organization :

CNAMTS

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

the objectives of this study were to examine whether benzodiazepine use has been associated with recurrent falls while taking into account the effect of potential confounders

## **Inclusion criteria**

Subjects aged 65 and over, volunteers, examined in one of the 3 health centres of Lorraine

---

# - CROSS-SECTIONAL STUDY OF PREGNANT WOMEN OR WOMEN WHO HAVE RECENTLY GIVEN BIRTH, ON EXPOSURE, KNOWLEDGE AND PERCEPTION OF THE RISKS OF TOBACCO

Last update : 10/20/2017

Head :

Dumas Agnès, Centre for Epidemiological and Population Health Research (CESP)

Responsible organization :

Inserm (CESP)

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

The objective of this study was to measure prenatal and postnatal tobacco exposure, assess women's perceptions and knowledge and identify risk factors.

## Inclusion criteria

Pregnant women or women who have recently given birth from the Family Service file

Women volunteering to participate in the study

# FRAMIG 3 - CROSS-SECTIONAL STUDY ON THE RECOGNITION AND THERAPEUTIC MANAGEMENT OF MIGRAINE

Last update : 10/20/2017

Head :

Radat Françoise

Lucas Christian

Responsible organization :

TNS SOFRES

Société Française d'Etudes des Migraines et Céphalées

## Type of database

Study databases

## Study databases (details)

Repeated cross-sectional studies (except case control studies)

## Main objective

Assess the proportion of migraine sufferers who are aware of their illness



Determine the factors (disability, quality of life, psychiatric comorbidities, medical consultations) that may lead to becoming aware of migraine

Assess the influence of these factors on migraine attacks.

### **Inclusion criteria**

Randomly selected subjects from the ACCESS-SANTE database

## **GRIM 2005 - CROSS-SECTIONAL STUDY IN GENERAL POPULATION: DETERMINANTS AND MOTIVATIONS OF A SEEING A PHYSICIAN FOLLOWING MIGRAINES**

Last update : 10/20/2017

Head :

Nachit-Ouinekh Fatima

Radat Françoise, CNRS UMR 5287 - INCIA Aquitaine Institute for Cognitive and Integrative Neuroscience

Responsible organization :

GSK laboratory

### **Type of database**

Study databases

### **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

### **Main objective**

The main objective of the study was to identify the determinants that lead subjects of the general population with migraine headaches to see a physician.

Study of the factors that determine whether subjects continued medical care for their headaches after an initial consultation.

### **Inclusion criteria**

Random selection of the sample

Over 18 years

## **SYMPHONIE - LONGITUDINAL STUDY OF METABOLIC SYNDROME IN FRANCE, OBSERVATION, INTERVENTION AND EPIDEMIOLOGY**

Last update : 10/20/2017

Head :

Thomas Frédérique

Responsible organization :

Inserm - Institut national de la santé et de la recherche médicale

Université Paris Descartes

Centre Investigations Préventives et Clinique

Laboratoire Fournier

---

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

Study clinical and biological characteristics, as well as cardiovascular risk factors, in men and women with metabolic syndrome.

## Inclusion criteria

- >- male and female
- aged between 18 and 80 years
- with metabolic syndrome

# - STUDY ON PHYSICAL ACTIVITY AND SPORTS PRACTICES IN FRANCE 2010

Last update : 01/01/2020

Head :

Mission des études, de l'observation et des statistiques MEOS

Responsible organization :

Ministère de la Ville, de la Jeunesse et des Sports (mission des études, de l'observation et des statistiques (MEOS))

## Type of database

Study databases

## Study databases (details)

Repeated cross-sectional studies (except case control studies)

## Main objective

The purpose of this survey is to describe the physical activity and sporting practices in a precise way, whether independent or supervised, and to show, as far as possible, the greatest diversity of practices, emerging practices, to measure the number of members of a sports federation and characterise the sociodemographic profiles of the respondents, from those who practised regularly to those who did not practise at all.

The purpose of the accident component is to provide an overview of the reported prevalence of accidents related to these practices in the population aged 15 years and over residing in France, and to identify certain risk factors for these accidents, as well as the people most exposed to them

## Inclusion criteria

Random drawing of numbers

For the 'accident' component, over 15 years

---

# - CROSS-SECTIONAL STUDY ON NURSERY SCHOOLS: INDIVIDUAL SOCIAL CHARACTERISTICS, RESIDENTIAL CONTEXT AND PREVALENCE OF WEIGHT PROBLEMS IN EARLY CHILDHOOD

Last update : 10/20/2017

Head :

Verger Pierre, Unité 379

Guagliardo Valérie , UMR912-SE4S

Responsible organization :

Observatoire régional de la santé, Provence-Alpes-Côte d'Azur (ORS PACA)

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

The objectives are:

?to analyse the association between the prevalence and social characteristics at an individual level;

?to study, independently of the individual characteristics, the links between the prevalence and the residential context approached by the municipalities. For the latter, two variables are distinguished, one linked to the level of precariousness, the other to the type of locality, urban, peri-urban or rural.

## Inclusion criteria

Children aged 3.5 to 4.5 enrolled in the first and second year of nursery school in the PACA region in 2002-2003

# - LONGITUDINAL STUDY OF NEWBORNS: SCREENING, PREVALENCE AND TREATMENT OF DEAFNESS

Last update : 01/01/2020

Head :

Rouillon Isabelle

Responsible organization :

hôpital Necker ? Enfants-Malades

## Type of database

Study databases

## Study databases (details)

Longitudinal study (except cohorts)

## Main objective

Evaluate hearing screening for newborns born in the Ile-de-France region

## Inclusion criteria

Screening conducted in four maternity clinics between 1 October 2005 and 31 December 2010, in conjunction with a Deafness Diagnosis and Guidance Centre (CDOS)

Positive screening in maternity clinics: lack of response to automated auditory evoked potential of 35 dB on both sides

For diagnostic confirmation: no response to standard auditory evoked potential of 30 dB at the CDOS

## **CALIPSO - CROSS-SECTIONAL STUDY ON FISH, SEAFOOD AND OMEGA-3 CONSUMPTION IN FRENCH COASTAL POPULATIONS**

Last update : 10/20/2017

Head :

Bemrah Nawel, Risk Assessment Department

Sirot Véronique

Leblanc Jean-Charles

Volatier Jean-Luc

Responsible organization :

ANSES

## Type of database

Study databases

## Study databases (details)

Not-repeated cross-sectional studies (except case control studies)

## Main objective

Provide data on seafood consumption in a French coastal population and look at n-3 polyunsaturated fatty acid intake

## Inclusion criteria

Person over 18 living in a coastal area and consuming fish and/or seafood at least 2 times per week

## **- CROSS-SECTIONAL STUDY IN THE GENERAL POPULATION: MENTAL HEALTH IN 4 FRENCH REGIONS (ILE-DE-FRANCE, LORRAINE, RHÔNE-ALPES, NORMANDY)**

Last update : 10/20/2017

Head :

Kovess-Masféty Viviane, METIS Department, Epidemiology and biostatistics for decision-making in public health / Health Process and Psychopathology Laboratory (EA 4057)

Fermanian Christophe, Métis

Responsible organization :  
EHESP

## **Type of database**

Study databases

## **Study databases (details)**

Not-repeated cross-sectional studies (except case control studies)

## **Main objective**

Provide indicators of mental health needs based on social indicators to enable better planning based on the needs of a "population pool"

## **Inclusion criteria**

Subjects drawn randomly