- PGRx: Graves' disease

Head: Grimaldi - Bensouda Lamiae

Last update: 07/10/2012 | Version: 1 | ID: 2634

General

Identification

Detailed name PGRx: Graves' disease

CNIL registration number, number and date of CPP agreement, AFSSAPS (French Health Products Safety Agency) authorisation CNIL 907.309

General Aspects

Medical area Endocrinology and metabolism

Internal medicine

Others (details) incident cases of Graves' disease

Keywords Graves, pharmaco-epidemiology

Scientific investigator(s)

(Contact)

Name of the director Grimaldi - Bensouda

Surname Lamiae

Address 10 place de Catalogne, 75014 Paris

Phone 01 55 42 53 00

Email contact@la-ser.com

Collaborations

Funding

Funding status Private

Details LA-SER

Governance of the database

Sponsor(s) or organisation(s)

responsible

LA-SER

Additional contact Main features Type of database Type of database Study databases Additional featith institutions and services No Satebase of Graves' disease are recruited by a network of endocrinology, internal medicine and pediatric centers, and the referents by a network of general practitioners, distributed all across the territory. Database objective Main objective Surveillance and assessment of the risk of Graves' disease associated with drug or vaccine use in real life situations. Inclusion criteria For the cases: patients, males and females aged 0 to 49 years, having clinical or biological signs or symptoms suggesting Graves' disease declared by the specialist, for which the first symptoms of the episode occurred less than 12 months ago. For the controls: patients, males and females aged 0 to 49 years and older, having consulted a general practitioner. Population type Age Newborns (birth to 28 days) Infant (28 days to 2 years) Early childhood (2 to 5 years) Childhood (2 to 5 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years) Adulthood (25 to 44 years) Adulthood (45 to 64 years)	Organisation status	Private
Type of database Type of database Study databases Case control study A selection of health care professionals out by an intermediary Database recruitment is carried out as part of an interventional study Additional information regarding sample selection. Cases of Graves' disease are recruited by a network of endocrinology, internal medicine and pediatric centers, and the referents by a network of general practitioners, distributed all across the territory. Database objective Main objective Surveillance and assessment of the risk of Graves' disease associated with drug or vaccine use in real life situations. Inclusion criteria For the cases: patients, males and females aged 0 to 49 years, having clinical or biological signs or symptoms suggesting Graves' disease declared by the specialist, for which the first symptoms of the episode occurred less than 12 months ago. For the controls: patients, males and females aged 0 to 49 years and older, having consulted a general practitioner. Population type Age Newborns (birth to 28 days) Infant (28 days to 2 years) Early childhood (2 to 5 years) Childhood (6 to 13 years) Adolescence (13 to 18 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years)	Additional contact	
Type of database Study databases Study databases Study databases Case control study A selection of health care professionals A selection of health institutions and services No Database recruitment is carried out as part of an interventional study Additional information regarding sample selection. Cases of Graves' disease are recruited by a network of endocrinology, internal medicine and pediatric centers, and the referents by a network of general practitioners, distributed all across the territory. Database objective Main objective Surveillance and assessment of the risk of Graves' disease associated with drug or vaccine use in real life situations. Inclusion criteria For the cases: patients, males and females aged 0 to 49 years, having clinical or biological signs or symptoms suggesting Graves' disease declared by the specialist, for which the first symptoms of the episode occurred less than 12 months ago. For the controls: patients, males and females aged 0 to 49 years and older, having consulted a general practitioner. Population type Age Newborns (birth to 28 days) Infant (28 days to 2 years) Early childhood (2 to 5 years) Childhood (6 to 13 years) Adolescence (13 to 18 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years)	Main features	
Database recruitment is carried out by an intermediary Database recruitment is carried out as part of an interventional study A selection of health care professionals A selection of health institutions and services No Cases of Graves' disease are recruited by a network of endocrinology, internal medicine and pediatric centers, and the referents by a network of general practitioners, distributed all across the territory. Database objective Main objective Surveillance and assessment of the risk of Graves' disease associated with drug or vaccine use in real life situations. Inclusion criteria For the cases: patients, males and females aged 0 to 49 years, having clinical or biological signs or symptoms suggesting Graves' disease declared by the specialist, for which the first symptoms of the episode occurred less than 12 months ago. For the controls: patients, males and females aged 0 to 49 years and older, having consulted a general practitioner. Population type Age Newborns (birth to 28 days) Infant (28 days to 2 years) Early childhood (2 to 5 years) Childhood (6 to 13 years) Adulthood (19 to 24 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years)	Type of database	
Database recruitment is carried out by an intermediary Database recruitment is carried out as part of an interventional study Additional information regarding sample selection. Cases of Graves' disease are recruited by a network of endocrinology, internal medicine and pediatric centers, and the referents by a network of general practitioners, distributed all across the territory. Database objective Main objective Surveillance and assessment of the risk of Graves' disease associated with drug or vaccine use in real life situations. Inclusion criteria For the cases: patients, males and females aged 0 to 49 years, having clinical or biological signs or symptoms suggesting Graves' disease declared by the specialist, for which the first symptoms of the episode occurred less than 12 months ago. For the controls: patients, males and females aged 0 to 49 years and older, having consulted a general practitioner. Population type Newborns (birth to 28 days) Infant (28 days to 2 years) Early childhood (2 to 5 years) Childhood (6 to 13 years) Adolescence (13 to 18 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years)	Type of database	Study databases
Database recruitment is carried out as part of an interventional study Additional information regarding sample selection. Cases of Graves' disease are recruited by a network of endocrinology, internal medicine and pediatric centers, and the referents by a network of general practitioners, distributed all across the territory. Database objective Main objective Surveillance and assessment of the risk of Graves' disease associated with drug or vaccine use in real life situations. Inclusion criteria For the cases: patients, males and females aged 0 to 49 years, having clinical or biological signs or symptoms suggesting Graves' disease declared by the specialist, for which the first symptoms of the episode occurred less than 12 months ago. For the controls: patients, males and females aged 0 to 49 years and older, having consulted a general practitioner. Population type Age Newborns (birth to 28 days) Infant (28 days to 2 years) Early childhood (2 to 5 years) Childhood (6 to 13 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years)	Study databases (details)	Case control study
Additional information regarding sample selection. Cases of Graves' disease are recruited by a network of endocrinology, internal medicine and pediatric centers, and the referents by a network of general practitioners, distributed all across the territory. Database objective Main objective Surveillance and assessment of the risk of Graves' disease associated with drug or vaccine use in real life situations. Inclusion criteria For the cases: patients, males and females aged 0 to 49 years, having clinical or biological signs or symptoms suggesting Graves' disease declared by the specialist, for which the first symptoms of the episode occurred less than 12 months ago. For the controls: patients, males and females aged 0 to 49 years and older, having consulted a general practitioner. Population type Age Newborns (birth to 28 days) Infant (28 days to 2 years) Early childhood (2 to 5 years) Childhood (6 to 13 years) Adolescence (13 to 18 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years)		
network of endocrinology, internal medicine and pediatric centers, and the referents by a network of general practitioners, distributed all across the territory. Database objective Main objective Surveillance and assessment of the risk of Graves' disease associated with drug or vaccine use in real life situations. Inclusion criteria For the cases: patients, males and females aged 0 to 49 years, having clinical or biological signs or symptoms suggesting Graves' disease declared by the specialist, for which the first symptoms of the episode occurred less than 12 months ago. For the controls: patients, males and females aged 0 to 49 years and older, having consulted a general practitioner. Population type Age Newborns (birth to 28 days) Infant (28 days to 2 years) Early childhood (2 to 5 years) Childhood (6 to 13 years) Adolescence (13 to 18 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years)	out as part of an interventional	No
Main objective Surveillance and assessment of the risk of Graves' disease associated with drug or vaccine use in real life situations. Inclusion criteria For the cases: patients, males and females aged 0 to 49 years, having clinical or biological signs or symptoms suggesting Graves' disease declared by the specialist, for which the first symptoms of the episode occurred less than 12 months ago. For the controls: patients, males and females aged 0 to 49 years and older, having consulted a general practitioner. Population type Age Newborns (birth to 28 days) Infant (28 days to 2 years) Early childhood (2 to 5 years) Childhood (6 to 13 years) Adolescence (13 to 18 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years)		network of endocrinology, internal medicine and pediatric centers, and the referents by a network of general practitioners, distributed all across the
disease associated with drug or vaccine use in real life situations. For the cases: patients, males and females aged 0 to 49 years, having clinical or biological signs or symptoms suggesting Graves' disease declared by the specialist, for which the first symptoms of the episode occurred less than 12 months ago. For the controls: patients, males and females aged 0 to 49 years and older, having consulted a general practitioner. Population type Age Newborns (birth to 28 days) Infant (28 days to 2 years) Early childhood (2 to 5 years) Childhood (6 to 13 years) Adolescence (13 to 18 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years)	Database objective	
to 49 years, having clinical or biological signs or symptoms suggesting Graves' disease declared by the specialist, for which the first symptoms of the episode occurred less than 12 months ago. For the controls: patients, males and females aged 0 to 49 years and older, having consulted a general practitioner. Population type Age Newborns (birth to 28 days) Infant (28 days to 2 years) Early childhood (2 to 5 years) Childhood (6 to 13 years) Adolescence (13 to 18 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years)	Main objective	disease associated with drug or vaccine use in real
Age Newborns (birth to 28 days) Infant (28 days to 2 years) Early childhood (2 to 5 years) Childhood (6 to 13 years) Adolescence (13 to 18 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years)	Inclusion criteria	to 49 years, having clinical or biological signs or symptoms suggesting Graves' disease declared by the specialist, for which the first symptoms of the episode occurred less than 12 months ago. For the controls: patients, males and females aged 0 to 49 years and older, having consulted a general
Infant (28 days to 2 years) Early childhood (2 to 5 years) Childhood (6 to 13 years) Adolescence (13 to 18 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years)	Population type	
	Age	Infant (28 days to 2 years) Early childhood (2 to 5 years) Childhood (6 to 13 years) Adolescence (13 to 18 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years)
Population covered General population	Population covered	General population

Gender	Male Woman
Geography area	International
Detail of the geography area	France, Italy, Spain, United Kingdom, Canada
Data collection	
Dates	
Date of first collection (YYYY or MM/YYYY)	04/2008
Size of the database	
Size of the database (number of individuals)	[10 000-20 000[individuals
Details of the number of individuals	- 153 cas/cases - 11125 référents/controls
Data	
Database activity	Current data collection
Type of data collected	Clinical data Declarative data
Clinical data (detail)	Direct physical measures
Declarative data (detail)	Phone interview
Presence of a biobank	No
Health parameters studied	Health event/morbidity
Procedures	
Participant monitoring	No
Links to administrative sources	No
Promotion and access	
Promotion	
Access	
Terms of data access (charter for data provision, format of data, availability delay)	Methods for accessing the database are currently being defined

Access to aggregated data	Access on specific project only
Access to individual data	Access on specific project only