

SURDIAGENE - Genetic Polymorphism in Subjects with Type 2 Diabetes and Longitudinal Analysis of Renal Function: ? SURDIAGENE? Study (SUivi Rénal , DIAbète de type 2 et GENEtique)

Head :Hadjadj Samy, CIC 1402

Last update : 08/07/2019 | Version : 4 | ID : 4071

General

Identification

Detailed name Genetic Polymorphism in Subjects with Type 2 Diabetes and Longitudinal Analysis of Renal Function: ?SURDIAGENE? Study (SUivi Rénal , DIAbète de type 2 et GENEtique)

Sign or acronym SURDIAGENE

CNIL registration number, number and date of CPP agreement, AFSSAPS (French Health Products Safety Agency) authorisation CPP : N°03.10.19 du 24/11/2003 ; DGS 2003/0530 du 15/12/2003 ; CNIL : en attente

General Aspects

Medical area Endocrinology and metabolism
Urology, andrology and nephrology

Health determinants Genetic
Medicine
Nutrition

Keywords Health episodes, renal function, cardiovascular, death, population, health system

Scientific investigator(s) (Contact)

Name of the director Hadjadj

Surname Samy

Address Centre d'investigation Clinique - Inserm 1402 ; 5 cour Est Jean Bernard ; BP 577 ; 86021 Poitiers Cedex

Phone + 33 (0)5 49 44 46 89

Email samy.hadjadj@gmail.com

Unit	CIC 1402
Organization	CHU Poitiers

Collaborations

Participation in projects, networks and consortia	Yes
---	-----

Funding

Funding status	Mixed
----------------	-------

Details	PHRC régional 2000 / PHRC interrégional 2004 / association GEMMS
---------	--

Governance of the database

Sponsor(s) or organisation(s) responsible	CHU de Poitiers
---	-----------------

Organisation status	Public
---------------------	--------

Additional contact

Name of the contact	GROSDENIER
---------------------	------------

Surname	Michele
---------	---------

Address	Centre d eRessources Biologiaues 5 cour Est Jean Bernard ; BP 577 ; CHU Poitiers 86021 Poitiers Cedex France
---------	--

Phone	+33549444689
-------	--------------

Email	michele.grosdenier-bruneau@chu-poitiers.fr
-------	--

Unit	Biobank
------	---------

Organization	CHU Poitiers
--------------	--------------

Main features

Type of database

Type of database	Study databases
------------------	-----------------

Study databases (details)	Cohort study
---------------------------	--------------

Database recruitment is carried	A selection of health institutions and services
---------------------------------	---

out by an intermediary

Database recruitment is carried out as part of an interventional study

No

Additional information regarding sample selection.

Prospective Inclusion cut-off date: 01/01/2009
Number of required subjects: [1,000-5,000] Details regarding this number: Initial calculation based on data from genotype frequency less relevant in 2009 compared to 2002 (date of conception).

Database objective

Main objective

MAIN: to investigate the genetic determinants associated with changes in renal function (changes in creatinine clearance) SECONDARY: 1. - to investigate the genetic determinants associated with changes in urinary albumin excretion 2 - to investigate the genetic determinants associated with other degenerative complications of diabetes (retinopathy, cardiovascular events)

Inclusion criteria

B1. Type 2 diabetes (T2D): Subjects from the SURDIAGENE study with T2D. The distinction between type 1 (TD1) and type 2 (T2D) diabetes is sometimes difficult. We define criteria for T2D diagnosis as: - age at diagnosis is 35 years old or over - apparent insulin-dependence after more than 2 years following diabetes diagnosis - absence of secondary diabetes - absence of ketonuria at diagnosis B2: Longitudinal follow-up: Subjects in the SURDIAGENE study with T2D who are being monitored for diabetes for at least one year, which includes a collection of clinical (blood pressure, weight) and biological data (glycated haemoglobin, serum creatinine, urinary albumin excretion) to analyse the progression of renal disease and other degenerative (retinal and cardiovascular) complications B3. Exclusion criteria is defined as: Subjects who are not permitted to participate in the SURDIAGENE study are subjects who a) fulfil medical criteria for exclusion - do not have diabetes or have T1D or secondary diabetes - are presenting renal disease caused by non-diabetic renal insufficiency (arterial hypertension is not considered as part of the exclusion criteria, except if hypertensive nephropathy precedes discovery of diabetes). - proteinuria at time of diabetes diagnosis - non-diabetic glomerular nephropathy confirmed by renal biopsy - pure vascular nephropathy - confirmed glomerular haematuria or malformed uropathy - chronic interstitial nephritis with history

of pyelonephritis or persistent urinary tract infections - follow-up is less than 1 year b) fulfilling legal exclusion criteria (amended law dated 20th December 1988) - minors - adults protected by law - deprived of liberty - brain-dead - who have not given their written participation consent - women who are pregnant, parturient or breastfeeding - not affiliated to social security

Population type

Age
Adulthood (25 to 44 years)
Adulthood (45 to 64 years)
Elderly (65 to 79 years)

Population covered
Sick population

Pathology
E11 - Type 2 diabetes mellitus

Gender
Male
Woman

Geography area
Local

Detail of the geography area
Poitiers (Vienne)

Data collection

Dates

Date of first collection (YYYY or MM/YYYY)
12/2003

Date of last collection (YYYY or MM/YYYY)
2020

Size of the database

Size of the database (number of individuals)
[1000-10 000[individuals

Details of the number of individuals
1200

Data

Database activity
Current data collection

Type of data collected
Clinical data
Biological data

Clinical data (detail)
Direct physical measures

Medical registration

Biological data (detail)	Renal function and glycaemic control: Renal function is based on a measurement of plasma creatinine and estimated glomerular filtration rate (GFR estimated by MDRD and CKD-EPI equations). Diabetic renal insufficiency (initial and during follow-up) is equally characterised by urinary albumin excretion and urinary albumin/urinary creatinine ratio. The level of renal impairment as well as GFR can be established as such. Glycaemic control rate is calculated by measuring glycated haemoglobin (HbA1c).
Presence of a biobank	Yes
Contents of biobank	Serum Plasma DNA Others
Details of biobank content	Other: urine, buffy coat
Health parameters studied	Health event/morbidity Health event/mortality

Procedures

Data collection method	Interview: input from a paper questionnaire (manual input) with double entry Clinical examinations: handwritten (manual input) with double data entry Biological analysis: direct input
Participant monitoring	Yes
Details on monitoring of participants	Follow-up duration: 10 years
Links to administrative sources	Yes
Linked administrative sources (detail)	CépiDC

Promotion and access

Promotion

Link to the document	http://www.ncbi.nlm.nih.gov/pubmed/?term=SURDIAGENE
Description	List of publications in Pubmed
Link to the document	Surdiagene.pdf

Access

Terms of data access (charter for data provision, format of data, availability delay)

Send request to scientist in charge

Access to aggregated data

Access on specific project only

Access to individual data

Access on specific project only