

ARCHIMED VILLE - Prevention of venous thromboembolic events with Arixtra 2.5mg for medical ill patients in general practice

Head :Leclerc-Zwirn Christel, Laboratoire GSK

Last update : 01/01/2020 | Version : 1 | ID : 142

General

Identification

Detailed name Prevention of venous thromboembolic events with Arixtra 2.5mg for medical ill patients in general practice

Sign or acronym ARCHIMED VILLE

CNIL registration number, number and date of CPP agreement, AFSSAPS (French Health Products Safety Agency) authorisation CNIL : 906318

General Aspects

Medical area General practice

Others (details) venous thromboembolic events

Keywords pharmaco-epidemiology, thrombosis, fondaparinux, arixtra

Scientific investigator(s) (Contact)

Name of the director Leclerc-Zwirn

Surname Christel

Phone +33 (0)1 39 17 86 96

Email christel.c.leclerc-zwirn@gsk.com

Unit Laboratoire GSK

Collaborations

Funding

Funding status Private

Details GSK laboratory

Governance of the database

Sponsor(s) or organisation(s) responsible LABORATOIRE GSK

Organisation status Private

Additional contact

Main features

Type of database

Type of database Study databases

Study databases (details) Longitudinal study (except cohorts)

Database recruitment is carried out by an intermediary A selection of health care professionals

Database recruitment is made on the basis of: Medication(s) taken

Database recruitment is carried out as part of an interventional study No

Additional information regarding sample selection. Investigators will be selected via random drawing using a complete sampling frame of the general practitioners exercising in metropolitan France. He will complete a specific inclusion questionnaire for the first 3 patients that meet the eligibility criteria.

Database objective

Main objective Evaluate the average duration of treatment via ARIXTRA® 2.5 mg with thromboprophylaxis in ambulatory general practice

Inclusion criteria Patient of at least 18 years of age,
-bedridden or with reduced mobility,
-for whom a treatment via ARIXTRA® 2.5 mg is initiated with thromboprophylaxis.

Population type

Age Adulthood (19 to 24 years)
Adulthood (25 to 44 years)
Adulthood (45 to 64 years)
Elderly (65 to 79 years)
Great age (80 years and more)

Population covered	Sick population
Gender	Male Woman
Geography area	National
Detail of the geography area	France
Data collection	
Dates	
Date of first collection (YYYY or MM/YYYY)	2007
Date of last collection (YYYY or MM/YYYY)	2010
Size of the database	
Size of the database (number of individuals)	[500-1000[individuals
Details of the number of individuals	834
Data	
Database activity	Data collection completed
Type of data collected	Clinical data Declarative data
Clinical data (detail)	Medical registration
Declarative data (detail)	Face to face interview
Presence of a biobank	No
Health parameters studied	Health care consumption and services
Care consumption (detail)	Medicines consumption
Procedures	
Data collection method	Throughout the entire duration of the study, the doctor will log the prescriptions of ARIXTRA® 2,.5 mg for thromboprophylaxis in a registry. This registry will make it possible to evaluate prescription volume and the representativeness of the patients effectively included in the study. The doctor will

complete for each patient included a follow-up questionnaire at the end of treatment.

Participant monitoring

Yes

Details on monitoring of participants

The data that will be collected in the follow-up questionnaire are: ? Premature interruption of the treatment, ? total duration of the administration of the treatment, ? prescriptions for prophylactic purposes after the treatment.

Links to administrative sources

No

Promotion and access

Promotion

Access

Terms of data access (charter for data provision, format of data, availability delay)

Publications in progress

Access to aggregated data

Access on specific project only

Access to individual data

Access on specific project only