- PrimInject: cross-sectional study on the patterns and practices of drug use in France

Head :Arwidson Pierre Guignard Romain Guichard Anne

Last update : 09/19/2017 | Version : 1 | ID : 73177

General	
Identification	
Detailed name	PrimInject: cross-sectional study on the patterns and practices of drug use in France
General Aspects	
Medical area	Infectious diseases
Health determinants	Addictions Lifestyle and behavior
Scientific investigator(s) (Contact)	
Name of the director	Arwidson
Surname	Pierre
Address	12, rue du Val d'Osne 94415 Saint-Maurice cedex
Email	pierre.arwidson@santepubliquefrance.fr
Organization	Santé publique France
Name of the director	Guignard
Surname	Romain
Address	12, rue du Val d'Osne 94415 Saint-Maurice cedex
Phone	0171801575
Email	romain.guignard@santepubliquefrance.fr
Organization	Santé publique France
Name of the director	Guichard

Surname	Anne
Organization	Université Laval, Québec
Collaborations	
Participation in projects, networks and consortia	No
Funding	
Funding status	Public
Details	Inpes
Governance of the database	
Sponsor(s) or organisation(s) responsible	Santé publique France
Organisation status	Public
Presence of scientific or steering committees	Yes
Additional contact	
Main features	
Type of database	
Type of database	Study databases
Study databases (details)	Not-repeated cross-sectional studies (except case control studies)
Database recruitment is carried out as part of an interventional study	No
Additional information regarding sample selection.	Online recruitment of participants
Database objective	
Main objective	Intravenous injection is the major mode of transmission of HCV among drug users, not only by needle sharing but also by the preparation equipment (container, filter, buffer, water), potentially contaminated by the blood of a contaminated person. According to various sources, in recent years there

	have been significant changes in the patterns and practices of drug use in France, under the combined effect of the massive distribution of opiate substitution treatments (high-dose buprenorphine or BHD) from 1995 and the introduction of a wide variety of stimulants, including cocaine and synthetic drugs. Often unplanned and practiced by a third party "initiator", the first injections present a high risk of contamination for the young initiate, often not well acquainted with injection techniques and the infectious risks and not in contact with first-line medical support structures. In this context, a survey was put in place to identify the circumstances and practices of the first injection(s) among young drug users in France. An online methodology using the Internet has been developed to provide access and thus to discover user profiles that escape the more "classic" observation systems, especially those that are socially integrated or that are not in contact with support structures.
Inclusion criteria	Be over 15 years old. Individuals who had never injected drugs were redirected to a short version of the questionnaire.
Population type	
Age	Adolescence (13 to 18 years) Adulthood (19 to 24 years) Adulthood (25 to 44 years) Adulthood (45 to 64 years)
Population covered	General population
Pathology	
Gender	Male Woman
Geography area	National
Detail of the geography area	France
Data collection	
Dates	
Date of first collection (YYYY or MM/YYYY)	October 2010
Date of last collection (YYYY or	March 2011

MM/YYYY)	
Size of the database	
Size of the database (number of individuals)	[500-1000[individuals
Details of the number of individuals	449 injectors and 455 non-injectors
Data	
Database activity	Data collection completed
Type of data collected	Declarative data
Declarative data (detail)	Internet self-questionnaire
Details of collected declarative data	Sociodemographic data, uses of psychoactive substances, practices and attitudes related to the first injection, HIV and HCV infection
Presence of a biobank	No
Health parameters studied	Health event/morbidity Others
Other (detail)	Uses of psychoactive substances, practices and attitudes related to the first injection
Procedures	
Data collection method	Internet. Promotion of the survey via a campaign of web banners, posters in cart-com format, articles and press releases.
Quality procedure(s) used	Pre-test of tools and questionnaire for intravenous drug users. Verification and cleaning of the database in Stata.
Participant monitoring	No
Followed pathology	
Links to administrative sources	No
Promotion and access	
Promotion	
Link to the document	http://www.ncbi.nlm.nih.gov/pubmed/23726980

Description	Practices related to the first injection according to the key periods of the risk reduction policy in France. Article published in Drug and Alcohol Dependence.
Link to the document	http://www.federationaddiction.fr/actal-n10- decembre-2011-hepatites-de-reelles-avancees/
Description	The first injection in the risk reduction prism. Article published in the Actal journal of the Fédération Addiction.
Access	
Presence of document that lists variables and coding procedures	No
Terms of data access (charter for data provision, format of data, availability delay)	On request from Pierre Arwidson, Director of Scientific Affairs at Inpes.
Access to aggregated data	Access on specific project only
Access to individual data	Access on specific project only